MBL/WHOI

0 0301 0013859 0

THE OEDOGONIACEAE

A MONOGRAPH

Including all the Known Species of the Genera
Bulbochaete, Oedocladium and Oedogonium

By

LEWIS HANFORD TIFFANY, M. Sc., Ph. D.

Associate Professor of Botany

The Ohio State University

1930

PUBLISHED BY THE AUTHOR
COLUMBUS, OHIO

THE SPAHR & GLENN COMPANY
PRINTERS
COLUMBUS, OHIO

FOREWORD.

For many years the only available keys to the Freshwater Algæ of North America were those published by Francis Wolle about 1887. These were in larger part transcriptions of European descriptions often based on a single collection. Moreover, at that time it was a common practice to name specimens on the basis of vegetative characters or of immature reproductive structures. Consequently many of the records are erroneous and many others are questionable.

In 1909 Frank S. Collins published a monograph of the Green Algæ of North America. He not only added many species and records to the North American algal flora, but also corrected many of the previous errors in nomenclature and in the locality records, and brought the taxonomy of the green algæ in America into harmony with the best usage in Europe. Collins' keys stimulated a wide interest in the green algæ in this country, and led to the publication of many local lists and papers describing new species.

In the determination of algæ, however, descriptions without plates are frequently difficult to interpret, and the greatest need of American students today is a series of illustrated monographs of the various families and orders.

The world wide distribution of many of the freshwater algæ, and the possibility that a species described from Australia may be found in Ohio has made it desirable that the monographs should attempt to cover all the described species. Too often the literature of a group is widely scattered and many of the papers are in obscure journals quite inaccessible to the average student.

In 1900 Karl E. Hirn of Helsingfors, Finland, published in Latin and German a monograph of the Oedogoniaceæ that has been a model and inspiration for all algologists, but this volume and its supplement are now difficult to secure. Moreover many new species have been described and the descriptions by Hirn in many cases have been emended. It has seemed therefore very desirable to publish descriptions and keys to all

the known species of Oedogoniaceæ in English. Dr. Tiffany's extensive acquaintance with American collections, and the species found in European exsiccati over a period of fifteen years have given him an excellent foundation for a critical view of the group.

It is the hope of the Department of Botany at The Ohio State University that this monograph will be followed in the near future by a monograph of the Zygnemales, and subsequently of other groups of the filamentous algæ.

EDGAR N. TRANSEAU.

TABLE OF CONTENTS

	PAGE
I. Acknowledgments.....	9
II. Introduction.....	11
III. Cell Structure, Reproduction, and Life History.....	12
IV. Periodicity and Distribution.....	22
V. Taxonomic and Systematic Treatment.....	24
1. The Family Oedogoniaceae.....	27
a. Description.....	27
b. Key to the Genera.....	27
2. The Genus Bulbochaete.....	27
a. Description.....	27
b. Key to the Species, Varieties, and Forms.....	28
c. The Species, Varieties, and Forms.....	30
3. The Genus Oedocladium.....	51
a. Description.....	51
b. Key to the Species.....	51
c. The Species.....	51
4. The Genus Oedogonium.....	53
a. Description.....	53
b. Key to the Species, Varieties, and Forms.....	53
c. The Species, Varieties, and Forms.....	65
d. Species Imperfectly Known.....	159
VI. Table of Synonyms: Bulbochaete.....	173
VII. Table of Synonyms: Oedogonium.....	174
VIII. Bibliography.....	177
IX. Exsiccatae.....	184
X. Index.....	185
XI. Plates.....	189

ACKNOWLEDGMENTS.

In attempting a complete report of a group as large as the Oedogoniaceæ, one encounters numerous difficulties. The literature is bulky and in many languages. Type specimens are unavailable in some cases. Thanks, however, to the very careful work of Hirn (1900), many forms are perpetuated by extremely accurate and representative drawings. Some of the difficulties are matched, on the other hand, by the relative constancy of the characteristics of the species as they occur in aquatic habitats the world over. Many species appear in the writer's own collections.

It is almost impossible to give full acknowledgment to all who have assisted in making this monographic report possible. The writer has examined the material distributed in the *Phycotheca Borealis-Americana* and important collections generously supplied by the following: Prof. E. N. Transeau, Ohio State University; the late Prof. O. Nordstedt; Prof. R. B. Wylie, Iowa State University; Prof. G. E. Nichols, Yale University; Prof. F. E. Lambert, Tufts College; Prof. W. R. Taylor, University of Pennsylvania; Prof. C. J. Chamberlain, University of Chicago; Prof. I. F. Lewis, University of Virginia; Mr. G. H. Wailes, British Columbia; Prof. Alma B. Ackley, College of the City of Detroit; Prof. B. B. McInteer, University of Kentucky.

Data on distribution of the genus outside of the United States other than that secured from the literature have been supplied by Mr. G. H. Wailes, British Columbia; Prof. O. Borge, Stockholm; Prof. K. M. Strom, Oslo; Prof. H. Skuja, Riga; Prof. N. N. Woronichin, Leningrad.

The section on *Bulbochaete* in this report has previously appeared, in substance, in the *TRANSACTIONS OF THE AMERICAN MICROSCOPICAL SOCIETY* (Tiffany 1928). The text and plates dealing with that genus are incorporated herewith with the permission of the Society, kindly extended through its Secretary, Professor H. J. Van Cleave. My thanks are due Professor I. F. Lewis, University of Virginia, for the loan of his unpublished manuscript on the genus *Oedocladium*. Professor Lewis is

responsible for the emended description of the genus and for three of the four species now known (two are previously undescribed).

I am particularly indebted to my colleague, Professor E. N. Transeau, who has very graciously placed at my disposal all his notes on Oedogoniaceæ recorded during his long study of freshwater algæ in the United States. He has offered many valuable suggestions during the fifteen years this study has been in progress.

L. H. T.

Department of Botany, The Ohio State University,
Columbus, Ohio, January 2, 1930.

THE OEDOGONIACEÆ

INTRODUCTION.

Perhaps the most sharply circumscribed and clearly delimited group among the green algæ is the family Oedogoniaceæ. The members of its three genera—Bulbochaete, Oedocladium, and Oedogonium—are not only easily separable from other filamentous algæ but also readily distinguishable from each other. The rather rigid vegetative cells, the prominent reproductive organs, the large zoospores each with a crown of cilia at the anterior end, the reticulate chloroplasts, and the attached habit are characteristics of the family.

The filaments of Bulbochaete are branched and composed of cells each bearing at its enlarged upper extremity a laterally placed hair or seta. The setæ are bulbous at the base, hollow, and frequently greatly attenuated, sometimes reaching a millimeter in length. Oedocladium is also branched but is devoid of hairs. Species of Oedogonium are unbranched and in common with Bulbochaete possess holdfast cells that provide attachment for the plants. *Oedocladium hazenii** and species of both Oedogonium and Bulbochaete are aquatic. The remaining three species of Oedocladium are terrestrial.

The family Oedogoniaceæ was established by De Bary in 1854; the genus Bulbochaete by C. A. Agardh in 1817; Oedocladium by Stahl in 1891; and Oedogonium by Link in 1820. The most noteworthy contributors to the taxonomy of the group, other than those above, have been Pringsheim, Wittrock, Nordstedt, and Hirn. Since 1900 a number of algologists have added to our knowledge of the family by descriptions of new species, further notes on life histories, and data on geographic distribution. A list of such workers includes Hirn and Hallas from Finland; William West, G. S. West, Fritsch, Carter, and Hodgetts from England; Borge from Sweden; Wille and Strom from Norway; Pascher and Heering from Germany; Skuja from Latvia; Collins, Transeau, Lewis, and Tiffany from the United States. There have been many other

*In the Index the name of each new species, variety, or form is preceded by an asterisk.

students of this algal family, and some of their names are found in the bibliography near the end of this monographic report.

Some idea of the relative sizes of the three genera of the Oedogoniaceæ may be gained by a numerical recital of the composition of each. The genus *Bulbochaete* is composed of 48 species, 15 varieties, and 7 forms, with 3 additional species having incomplete descriptions. In *Oedogonium* there are 195 species, 76 varieties, and 38 forms; in addition there are 41 species and 3 varieties incompletely known. *Oedocladium*, by far the smallest, has only 4 species recorded to date.

Although an *Oedogonium*, a *Bulbochaete*, or even an *Oedocladium* can be recognized on vegetative characters alone, the identification to species is impossible, except in a very few cases, without reproductive organs. Such unusual forms as *Oe.* undulatum*, *B. horrida*, *Oe. nodulosum*, or *Oe. reinschii* are perhaps always recognizable in vegetative states, but such species are few in number.

The comprehensive reports of Pringsheim (1858), Wittrock (1874), Hirn (1900, 1906), and Heering (1914) are not readily available to many students of the algæ, and their points of emphasis have been necessarily European. The early American accounts of the Oedogoniaceæ by Wood (1872) and by Wille (1887) are quite unreliable. The work of Collins (1909, *et seq.*) made available much European literature and was the first American publication on the green algæ that was really trustworthy. Perhaps to Collins should go the credit for the initiation of real interest in freshwater algæ in North America. Unfortunately his papers were inadequately illustrated. It is hoped that furnishing a complete English treatise of the family Oedogoniaceæ, including in addition to the taxonomic consideration, a discussion of the more important features of structure, reproduction, and distribution, will not only aid the student in the study of these common algæ but will stimulate added interest in algology generally.

CELL STRUCTURE AND REPRODUCTION.

The cell wall of the Oedogoniaceæ exhibits a rigidity not common to other filamentous Chlorophyceæ. The heavy cellulose layer is covered peripherally with pectose, and upon

*The abbreviation *Oe.* as used in this paper refers to the genus *Oedogonium*, and not to *Oedocladium*. When the latter is meant, the abbreviation is *O.*

this is found in varying amounts another compound, resembling chitin. In *Bulbochaete* and *Oedogonium* the latter substance does not appear on the holdfast cell or dwarf male at the points of attachment. The cell may frequently be so heavily incrustated with lime that identification is not possible in that condition. The addition of warm lactic acid removes the calcium carbonate without undue change to the filament. Epiphytic small algæ, including species of *Oedogonium* and *Bulbochaete*, are commonly found attached to the rigid filaments. Fungal parasites, particularly certain Chytridiales, attack the plants both internally and externally.

The vegetative cells are uninucleate with an elaborate parietal, reticulate chloroplast having one to many scattered pyrenoids. The chloroplast is quite variable in form, the network being considerably attenuated or hardly visible in the compact cylindrical mass (Carter, as reported by Fritsch and West, 1927).

A rather peculiar characteristic of the family is the method of division of the vegetative cell which results in the so-called "apical caps." Previous to the actual division there appears a ring-like thickening of the inner wall of the cell near its upper extremity. The growth of this ring ruptures the old wall circularly, and the outer layer of the ring becomes the new piece of cell wall. In the meantime nuclear division is followed by the formation of a cross wall at the middle of the cell, not connected at first with the cylinder. Later the septum assumes a permanent position near the lower part of the ruptured wall. The upper cell thus formed has a wall made up chiefly of the stretched ring, while the wall of the lower cell consists almost wholly of the lower part of the old cell membrane. The upper cell, however, has at its upper extremity a part of the old cell wall which projects slightly, forming the cap. This apical cap is an excellent diagnostic character of the genus *Oedogonium*, although in some of the smaller species it is difficultly visible; and the number of caps in each cell denotes the number of divisions it has undergone. (Cf. Pl. LXI, figs. 598-600).

In *Bulbochaete* there is rarely more than one apical cap to each cell, indicating, as will be explained later, that vegetative division in that genus occurs usually but once in a cell. (Cf. Pl. X, fig. 99).

In *Oedocladium* usually only the terminal cell undergoes division as outlined above for *Oedogonium*. Intercalary growth

is apparently uncommon. The cap (or successive caps) may remain at the top of the terminal cell (Pl. LXIII, figs. 616 and 617 and 621; Pl. LXI, fig. 602) or be pushed aside, adhering to the side of the filament (Pl. LXII, fig. 611; Pl. LXIII, fig. 622).

The nuclear phenomena accompanying cell division were early investigated by Strasburger (1880) and Klebahn (1892). More recently Tuttle (1910) has compared mitosis in *Oedogonium* with that seen in higher plants.

A single multiciliate zoospore is formed from the entire contents of a vegetative cell. Rarely a small amount of protoplasm is visible in the old cell after the emergence of the zoospore. The formation and liberation of the zoospore can readily be followed if the plants are brought into a warm room and observed under the microscope. The protoplast of the cell contracts slightly, and near one side appears a colorless area. A circular rent soon appears near the upper extremity of the cell in the vicinity of the apical cap. The mass then passes with a slow steady movement through the opening in the wall, being considerably constricted at the place of exit. Zoospores are formed from any cell of the filament except the holdfast.

Upon release from the cell, the zoospore assumes a spherical or ovoid shape. It is at first surrounded by a thin gelatinous envelope which rather rapidly increases in volume and transparency and finally disappears. At the anterior end of the forming zoospore appears a colorless area, around the base of which occurs the crown of cilia. The zoospore is active for a short time, its movements becoming progressively less and less until it comes to rest. The cilia are absorbed and the clear anterior end becomes the attached part of the holdfast cell (Cf. Pl. X, figs. 84-92). Sometimes growth of the zoospore begins before movement has ceased. Upon germination, repeated cell division occurs in the manner described above, except in those species where the first division does not result in the formation of a ring. In *Oedogonium* the sporeling thus formed develops into a filament by intercalary growth.

The development of the germinated zoospore in *Bulbochaete* involves variations that deserve separate mention. The basal cell formed from the germinated zoospore is usually the only one capable of division, as far as the main axis is concerned. In some of the forms with ellipsoid oospores, however, inter-

calary growth has been noted. The first division is of simple type, without a ring formation. The upper part of the cell is separated from the lower by a circular rent, but no new membrane develops between the two parts. As the new wall grows upward, the separated part is pushed to one side as a lid. The upper new cell grows out into a long tubular bristle with a swollen bulbous base. Subsequent divisions occur with the regular ring formation, and each new cell is intercalated between the basal cell and the next one above. The upper cell of each filament is therefore the oldest one. A lateral protrusion of each cell at its apex is cut off, developing into a hair. It is thus seen that the terminal cell bears two hairs, the other cells one each. Nearly every cell of *Bulbochaete* has two apical faces, one bearing a hair (or a cell of a side branch), the other the next upper cell of the main axis. Every cell of the principal axis of the filament, except the basal one, may cut off by an oblique division a lateral cell which becomes the basal cell for a side branch. By repeated divisions, first of the simple type and subsequent ones with ring formation, as noted above, this secondary basal cell (which of course is the primary basal cell of the side branch) forms the branch. Tertiary branches are formed similarly from the secondary ones, and so on through further branching. The mode of growth of the main axis is thus repeated by the side branches (See Plate X, figs. 93-99).

The germination of a zoospore of *Oedocladium* is in many respects unlike that of either *Oedogonium* or *Bulbochaete*. The first cell formed upon germination develops into (1) the green filament with the rhizoidal cell forming as a branch; or (2) the rhizoidal filament with the chlorophyllose cell as a side branch; or (3) the green filament with the branch cell also green; or (4) a green filament without early branching. (Cf. Pl. LXII, fig. 612; Pl. LXIII, fig. 623). The formation of a branch in *Oedocladium* occurs by the development of a circular rent in the upper extremity of a vegetative cell, forming an upper short "lid" and a lower, much longer cylindrical piece. Through this rent the protoplasm of the forthcoming daughter branch cell with its initially thin wall protrudes. A new wall separating the branch cell from the cell of the main filament is then developed. This wall is attached to the juncture of the upper extremity of the cylindrical piece of the primary cell and the lower end of the branch cell, on the one hand, and to the

original upper transverse wall of the primary cell, on the other. (Cf. Pl. LXII, figs. 606 and 612; Pl. LXIII, figs. 616 and 623).

Sexual reproduction of a rather advanced order has been observed in most species of the family. The position and relation of the oogonia and antheridia are used as classificatory items in all three genera. If oogonia and antheridia occur on the same filament the plant is *monoecious* (Pl. IX, fig. 69; Pl. XV, fig. 145); if on separate filaments, *dioecious* (Pl. XXXIII, figs. 291 and 292). *Oe. varians*, *Oe. trioicum*, and perhaps a few others are both dioecious and monoecious. If dioecious, the male filaments may be about equal in size to the female, and the species is *macrandrous* (Pl. XXX, figs. 262 and 263). If the male plants are few-celled, much smaller ("dwarf males" resulting from germinating androspores) and epiphytic on the female plant, the species is *nannandrous* (Pl. II, fig. 17; Pl. LII, fig. 494). If the dwarf males have arisen from the germination of androspores liberated from androsporangia occurring on the same filament with the oogonia, the alga is *gynandrosporous* (Pl. V, fig. 37; Pl. LIV, fig. 509; Pl. LXII, fig. 606); if the androsporangia are formed on separate filaments from the oogonia, the alga is *idioandrosporous* (Pl. VII, fig. 53; Pl. XLV, figs. 434 and 435). A few species among the nannandrous sections are both gynandrosporous and idioandrosporous. The dwarf males form on the suffultory cell (Pl. VII, fig. 52; Pl. L, fig. 482), or on the oogonium (Pl. V, fig. 37; Pl. LIV, fig. 508), or rarely scattered on vegetative cells (Pl. XLVII, fig. 458).

In *Oedogonium* and *Oedocladium* a single division of a vegetative cell results in an *oogonium* and a *suffultory* ("supporting") cell (Pl. LXIV, figs. 638–640). The latter (in *Oedogonium*) may by subsequent divisions form a series of oogonia (Pl. LIX, fig. 577). Flat discoid cells resulting from repeated transverse divisions of a vegetative cell are *antheridia* (in the monoecious and macrandrous species) and *androsporangia* (in the nannandrous species). No seriate oogonia are recorded for the species of *Oedocladium*.

In *Bulbochaete* oogonia are developed singly, never in series. An oogonium arises by a double division of a vegetative cell which thus results in two suffultory cells. After the first division the septum becomes fixed near the middle of the old sheath-like wall "so that the membrane of the upper segment is formed by a short cylindrical piece of the latter and above

that by the stretching thickening ring which is becoming bulged out, forming the oogonium. A second thickening ring then develops in the median region of the young oogonium followed by a transverse rupture of the wall at this point; the septum formed in connection with this second division lies on a level with the top of the first-formed sheath, so that the supporting-cells are included in the latter. This curious development is responsible for the fact that the wall of the mature *Bulbochaete* oogonium can nearly always be seen to consist of three pieces." (West and Fritch 1927, p. 219). (Cf. Pl. LXIV, figs. 641-645.)

If the lower of the two suffultory cells is the longer we speak of the division as supreme, superior, or supramedian, depending upon the relative lengths; if the upper is the longer of the two, the division becomes inframedian, inferior, or basal; if the two are equal in length, the division is median. In some species of *Bulbochaete* the position of the division is variable and cannot always be depended on as a distinctive diagnostic characteristic. If the two divisions are at right angles to the long axis of the vegetative cell, the oogonium is said to be *erect*; if one division is oblique, giving one suffultory cell a five-sided, or five-angled, appearance (viewed in optical section), the oogonium is *patent*. In members of the genus with globose (or nearly so) oogonia, the upper and not the lower suffultory cell is five-angled, while in those with more ellipsoid oogonia the reverse is true. In the latter case the upper cell is quite small, or may not be visible. A patent oogonium is always formed by a division of the basal cell of a side branch, with which may be associated a bulbous hair, an androsporangium, an antheridium, or rarely a vegetative cell. An erect oogonium on the other hand forms from a division of some vegetative cell other than the basal one. In the globose forms of the genus, patent oogonia are common while erect oogonia occur only in a few species. Among the ellipsoid forms both kinds are found. (Pl. LXIV, figs. 641-645).

The oogonium contains a single egg and varies considerably in shape, being largely ellipsoid, ovoid, globose, or some variation of these shapes. It is usually of noticeably larger diameter than the vegetative cell, although in *Oe. capillare* and in a few others the difference is not great. The oogonium wall is rarely internally ribbed (as in *Oe. acrosporum*), punctate (as in *Oe. minus*), plicate (as in *Oe. megaporum*, *Oe. platygynum*) or provided with a transversely disposed ring of conical warts (as in *Oe. itzigsohnii*); in most species it is smooth. The

suffultory cell may be noticeably larger than the other vegetative cells, or of the same diameter (Pl. LXIV, figs. 639-640). The opening in the wall of the oogonium may be a pore, either round and mouth-like, or rimiform, in which case the plant is designated as *poriferous*. An upper part of the oogonium may separate from a lower part, forming a lid; such species are *operculate*. The position of the pore as well as that of the operculum is quite constant for the species and is of considerable diagnostic value. The pore may be superior, supramedian, median, inframedian, or inferior. The division of the operculum

TABLE I.

Ready reference to illustrations of the kinds of ornamentations of oospore walls found in the Oedogoniaceae. It will readily be seen, upon examination of the figure, whether it is the middle or outer layer of the oospore wall with ornamentations.

OOSPORE WALL	SPECIES	PLATE	FIGURE
Angulate.....	<i>O. albemarlensis</i>	LXIII	614
Areolate.....	<i>Oe. areolatum</i>	XXVIII	248
Dentate.....	<i>Oe. cleveanum</i>	XLVI	443
Echinate.....	<i>Oe. hystricinum</i>	XLIII	414
Echinate.....	<i>Oe. hystrix</i> v. <i>canadense</i>	XLIII	418
Pitted.....	<i>Oe. giganteum</i>	XXIX	260
Reticulate.....	<i>Oe. dictyosporum</i>	XXVIII	246
Ribbed.....	<i>B. imperialis</i>	VIII	64
Ribbed.....	<i>B. bullardi</i>	VII	52
Ribbed.....	<i>Oe. paucocostatum</i>	XXXII	277
Ribbed.....	<i>Oe. crenulocostatum</i>	XXVIII	240
Ribbed.....	<i>Oe. exosporale</i>	XLIV	426
Smooth.....	<i>B. angulosa</i>	II	14
Smooth.....	<i>Oe. supremum</i>	LIX	573
Scrobiculate.....	<i>B. alabamensis</i>	V	38
Scrobiculate.....	<i>Oe. tiffanii</i>	XXX	261

may in addition to these positions be supreme or basal (*infimus*). (Pl. LXIV, figs. 624-635).

The oospore results from the development of the fertilized egg and is considered mature upon the formation of a distinct wall. It may or may not fill the oogonium and may be unlike it in shape. The oospore wall is usually three-layered, although occasionally of two layers. The middle or outer layer (it is invariably the latter in *Bulbochaete*) may be smooth or variously ornamented (See Table I). The kind of ornamentation is widely varied for the different species: pitted, scrobiculate, reticulate, angulate, dentate, areolate, echinate, or longitudinally or spirally ribbed. The ribs may be entire, crenulate, toothed,

and variously anastomosate. The color of the oospore changes with age from green to brown (or yellowish-brown) or red.

The oospore upon liberation from the rest of the filament by disintegration of the vegetative and oogonial cells sinks to the bottom of the pond or other body of water, or lodges on perennial aquatic macrophytes, or as in the terrestrial species of *Oedocladium* lies in the damp earth. After a period of rather prolonged dormancy, which apparently may vary from a few months to a few years, it begins to grow. Although the data have been secured for a few species only, it is the usual occurrence for the contents of the oospore to divide and form four zoospores at germination (Pl. X, figs. 78-83; Pl. LXIV, figs. 646-647). Upon their liberation from the old oospore wall, the zoospores swim about for a time, come to rest, and germinate into new filaments, precisely in the same manner as those produced directly from the vegetative cell. The reserve food in the oospore is a fatty oil dissolved in a reddish-brown pigment. The oospore is considered the diploid structure in the life history, the remainder of the plant being haploid.

The antheridium of the monoecious and dioecious-macrandrous species is derived from the division of a vegetative cell, the cross wall arising near the upper part of the cell. If no more divisions occur, we have a single antheridium, often designated as unicellular (Pl. XIV, fig. 136). Subsequent divisions result in a series of antheridia (2 to 45), termed multicellular (Pl. XIII, fig. 125). In the monoecious species the antheridia may be near the oogonium (epigynous, subepigynous, hypogynous, etc.) or scattered at some distance from the oogonium (Pl. XV). In the dioecious-macrandrous species the occurrence of seriate antheridia often results in considerable distortion of the filament, particularly if apically located (Pl. XVI, fig. 153). In *Bulbochaete* the antheridium may arise (as in the development of the oogonium) by a horizontal division of the vegetative cell, resulting in an *erect* antheridium (Pl. IX); if the division is oblique, the antheridium is part of a side branch terminated by a bristle: a *patent* antheridium (Pl. IX). Antheridia of *Oedocladium* usually occur on special branches, formed as in *Oedogonium*.

The contents of each antheridium produces one sperm, or as in most species, two sperms. The division of the antheridium when two sperms are formed is either *horizontal*, one lying superimposed on the other (Pl. XV, fig. 145); or *vertical*, the

sperms lying side by side (Pl. XX, fig. 182): In the dwarf males the division is always horizontal. The sperms look like small zoospores and are liberated in the same manner from the filament. After swimming actively for some time, the sperm enters the opening of the oogonium with subsequent fertilization of the egg and production of the oospore.

The formation of the antheridia in the dioecious-nannandrous species involves the production of *androsporangia*, the development of which is practically identical with that of the antheridia. The androsporangia may be developed singly or seriatly (Pl. V, fig. 37; Pl. LV, fig. 528; Pl. XLV, fig. 434). The entire contents of the androsporangium produces a single androspore (Pl. LXII, fig. 606), a spore that superficially resembles a sperm or zoospore in form, in manner of liberation from its sporangium (*androsporangium*), and in activity. Upon coming to rest, however, it germinates and produces a short filament known as the *dwarf male*, or *nannandrium* (Pl. LXIV, fig. 645D). This germination of the androspore occurs on the oogonium or suffultory cell, or rarely on scattered vegetative cells.

Division of the germinated androspore results in a lower non-reproductive *stipe* (holdfast cell) and an upper small cell, the *antheridium* (Pl. XLVI, fig. 443 A and S). Subsequent divisions of either stipe or antheridium produce additional antheridia (2 to 4) (Pl. XLVI, fig. 448). If merely an inner partition is formed, the antheridium is *internal* and single (Pl. LI, fig. 486); if the customary ring-formation takes place, the antheridium is termed *external* (Pl. LX, fig. 584). The dwarf males are rarely unicellular (Pl. LIII, fig. 505), and the stipe, though usually one-celled, may be few-celled (Pl. XLVII, fig. 459). Two sperms are normally formed in each antheridium by a transverse division of its contents. The basal cell of the dwarf male, formed as noted above, is usually an elongated, either simple or slightly lobed holdfast.

The basal cell of the filament, resulting as noted previously from the germinating zoospore, is most frequently elongated (Pl. LXIV, fig. 637). The attached lower end may be simple or much lobed. In some species the developing basal cell may be flattened into a subhemispherical cell or rarely appear nearly spherical (Pl. LXIV, fig. 636). The smaller species remain attached throughout almost their entire life history. The larger ones usually become detached and sometimes are

found in considerable mats on the surface of the water. There is no holdfast cell in *Oedocladium*.

In many of the larger species of the genus *Oedogonium*, in particular, and rarely in *Bulbochaete* the protoplast of the cell may form ovoid, ellipsoid, or irregularly shaped aplano-spores. In the terrestrial species of *Oedocladium* resting cells or "akinetes" occur on the rhizoidal filaments. They are in reality few-celled side branches with swollen cells containing reserve foods. Similar reserves sometimes accumulate in the vegetative cells and can thus endure prolonged desiccation (Stahl 1891). Such cells upon subsequent growth ("germination") may produce new plants.

There has been considerable speculation regarding the relationship between the macrandrous and nannandrous species of the Oedogoniaceæ. Hirn (1900) and West (1912a) are of the opinion that dwarf males of the nannandrous species (at least of *Oedogonium*) arose by the reduction and greater specialization of the zoospores normally producing the antheridial filaments of the macrandrous forms. They would assume as a beginning that the monoecious filaments as well as the zoospores are hermaphroditic. Differentiation of zoospores, some producing antheridial filaments only and others producing oogonial filaments only, gave rise to the dioecious, macrandrous species. Small male filaments of such plants as *Oe. fonticola* or *Oe. verrucosum* perhaps represent intermediate steps in further reduction and specialization leading to the dwarf male, epiphytic on or near the oogonium. Schaffner (1927) regards the androspores as having been derived phylogenetically from original sperms and as retaining sexuality to some degree; *i.e.*, they react toward the oogonium or suffultory cell sufficiently to lodge there, but not sufficiently to fuse with the egg. The dwarf male is thus formed parthenogenetically from an androspore. Pascher (1907) on the other hand states that the nannandrous species are not derived from the macrandrous species.

The numerical composition of the three genera of Oedogoniaceæ based on the nature of the sexual reproduction in the species presents an interesting comparison. In the genus *Bulbochaete* 42 species, 2 varieties, and 1 form are dioecious, nannandrous; 6 species, 2 varieties, and 1 form are monoecious; no dioecious, macrandrous species are now known.

In *Oedocladium* 3 species are monoecious and 1 dioecious, nannandrous. In the large genus *Oedogonium* 61 species, 15 varieties, and 8 forms are monoecious; 56 species, 25 varieties, and 11 forms are dioecious, macrandrous; and 77 species, 36 varieties, and 19 forms are dioecious, nannandrous. *Oe. varians* and *Oe. trioicum* are known to be both monoecious and dioecious. The above summary does not take into account any members of the Oedogoniaceæ with incomplete descriptions.

DISTRIBUTION AND PERIODICITY.

Species of *Bulbochaete* and of *Oedogonium* are widely distributed in freshwater, the world over. *Oedocladium* is so far reported only from the eastern United States and from

TABLE II.

Distribution by continents of the total number of species, varieties, and forms (grouped together) of the genera of the Oedogoniaceæ.

Genus	North America	South America	Europe	Asia	Africa	Australia
<i>Bulbochaete</i>	38	15	49	11	3	14
<i>Oedocladium</i>	3	0	1	0	0	0
<i>Oedogonium</i>	205	77	189	49	45	35

Germany; three of the species are terrestrial, growing on wet mud, and the fourth is aquatic. Hirn (1900) reports *Oe. capillare*, *Oe. oblongum*, *Oe. pluviale*, and *B. rhadinospora* var. *litoralis* from brackish waters of Europe. In a vegetative state *Oedogonium* is known as a part of the red and yellow snow of the Orkneys (Fritsch 1912). Species of both *Bulbochaete* and *Oedogonium* have been collected at considerable elevations in alpine regions, usually however not in fruiting condition.

Without distinguishing among species, varieties, and forms, the preceding table (Table II) shows the distribution of all members of the Oedogoniaceæ by continents. Purely as a matter of convenience, Australia is used to include not only that continent but also the nearby groups of islands of New Zealand, Melanesia, and Malaysia.

Data from the North Central States of the United States (Tiffany and Transeau 1927) indicate that most species of

Oedogonium and Bulbochaete grow in small permanent bodies of water rather than in lakes. The order of importance of algal habitats in this region of the United States for these two genera is: permanent ponds, lakes, temporary ponds, streams, and stream oxbows. Nearly 80% of the fruiting records for Oedogonium come from forms collected in the permanent ponds and the lakes. Streams are relatively poor habitats for fruiting species in spite of the prominent holdfast cells of the filaments. (See Table III).

TABLE III.

Data showing relative importance of Oedogonium habitats in the North Central States.

HABITAT	NUMBER OF RECORDS	PERCENTAGE
Permanent Ponds.....	603	54%
Lakes.....	254	23%
Temporary Ponds.....	172	15%
Streams.....	65	6%
Stream Oxbows.....	20	2%
Total.....	1114	100%

The above table gives the relative importance of Oedogonium habitats in the North Central States for the 1114 records. Nearly 100 records for the genus Bulbochaete show a very similar distribution among habitats in the same region.

Periodicity as applied to Oedogonium and Bulbochaete has been studied by Fritsch and Rich (1913), Hodgetts (1921), and Tiffany and Transeau (1927). It has been shown that the species of these genera produce reproductive structures at rather definite seasons of the year. The particular months of maximal sexual reproduction vary with the species, the geographical location, and the length of the growing season. The smaller forms as a rule have a short vegetative phase and fruit early in the growing season; the larger forms have usually a longer vegetative period and fruit later in the growing season. In north central United States maximal sexual reproduction is reached in May and in July (see text fig. 1) both in the annuals and perennials. Many species reach a second maximum in October, which is doubtless the development of a second generation.

An explanation of such periodicity in this algal group is not satisfactorily complete. It seems quite evident that a certain time—longer in some species than in others—must elapse, during which vegetative development occurs, before sexual reproduction takes place. When this vegetative phase is completed, it is almost impossible to prevent fruiting, except in extremes of temperature, very inadequate light, or toxic water. Hodgetts (1921) finds in observations on a small pond of Britain that there is some correlation between abundance of *Oedogonium* and (1) higher temperature and (2) competition with other algæ. We need data indicating the degree of

TEXT FIG. 1. Number of fruiting records each month for the genus *Oedogonium*, showing periods of maximum sexual reproduction.

relationship between algal growth and reproduction, on the one hand, and intensity and quality of light, mineral composition of the water, and hydrogen ion concentration, on the other.

Zoospore production may occur at practically any time before the formation of oospores. The germination of the zoospores produced directly from the vegetative cells accounts in large measure for the number of individual filaments of a given species found occupying a particular habitat.

TAXONOMIC CONSIDERATION.

In working through a group as large as the *Oedogoniaceæ* one finds it quite difficult to pass on the validity of some species. Such decisions, while always influenced to some extent by

personal interpretations, must rest upon one's knowledge of all species and their variations.

The ultimate criterion must be the constancy of such characteristics as methods of reproduction, size (within limits), shape of cells, opening of oogonia. How much are all these altered by ecological conditions of the body of water in which the algæ are growing? Unfortunately our data here are far from complete. When one finds a combination of the same characteristics in a given alga year after year in various habitats, one can be reasonably sure of the constancy of the species.

If an alga occurs year after year in the same habitat with a known species, differing in one or more morphological characteristics but evidently closely related to the type, it should be classed as a "variety." As an example, the robust variety (*majus*) of *Oe. grande* has been collected with the type in nearly all collections from the United States. Such differences can hardly be accounted for on the basis of environmental influences.

The category "form" should be regarded as a temporary disposition only. If an alga varies from the description of a given species and has been seen only once or twice, it is hardly possible to know whether the variation is constant or merely an ecological variation. If the form occurs with the type and is found on several occasions to maintain its morphological variation, it should be raised to varietal rank. A "form" then should be retained only when the data are insufficient to pass on the constancy of the variations noted in the description.

In the identification of members of the three genera—*Bulbochaete*, *Oedogonium*, and *Oedocladium*—and in the use of the keys to the species certain terminology is necessary to be able to differentiate among the species. The preceding pages have been written with the hope of elucidating such terminology and at the same time giving a brief introduction to the life history of species of the genera of the Oedogoniaceæ.

Keys to plant genera and species have not always been distinguished from synopses. It seems to the author that a key is merely an arrangement of characteristics, by which through proper manouvering, an "unknown" becomes a "known." A key has sometimes been defined as a scientific puzzle made by one expert to disturb another expert. Such keys may be interesting contributions if one already knows the field covered by them, but it is needless to say that the student, or any beginner for that matter, will find scant encouragement in them

to proceed further with the group concerned. On the other hand, of course, no key to plant forms can be so simple as to preclude the necessity of having a modicum of intelligence regarding scientific terms. Some species are more closely related than are others, and one must exercise judgment in the evaluation of characteristics. One measurement will not suffice, a single oogonium may be a monstrosity, one oospore may be parasitized, and a single vegetative cell may be far from typical. The keys of the following pages have been constructed in as simple a fashion as possible with the particular aim of enabling the student of the group to name his collection.

It is very convenient to use the group name *Stephanokontæ* proposed by Blackman and Tansley (1902) as a division of the class *Chlorophyceæ* and including those forms whose motile reproductive cells have a crown of cilia round the clear anterior end. Quite recently Fritsch (1927) has elevated the *Isokontæ* to be synonymous with and to replace *Chlorophyceæ* (i.e., the *Chlorophyceæ* exclusive of the *Heterokontæ*). *Oedogoniales* with its single family *Oedogoniaceæ* is further considered by Fritsch as a group division of *Isokontæ*.

There is no question but that the *Oedogoniales* have characteristics of cell wall, cell division, and reproductive structures that separate them as a quite readily distinguishable group. Even Fritsch remarks that they are a "sharply circumscribed group affording but few points of affinity with the other *Isokontæ*" (*loc. cit.*, p. 212). It appears to the writer to be to the best interests of taxonomic nomenclature to retain the group name *Stephanokontæ* as a division of the *Chlorophyceæ*, as noted above.

There is but one order, OEDOGONIALES. The simple or branched filaments are almost invariably attached, at least during a part of their life history, except in *Oedocladium*.

The single family, OEDOGONIACEÆ, may be characterized as follows:

OEDOGONIACEAE De Bary

1854, p. 94; Pringsheim 1858, p. 68; Wittrock 1874, p. 6; Hirn 1900, p. 71; Collins 1909, p. 222.

Plants aquatic or terrestrial; filaments simple or branching; cells uninucleate; chloroplast parietal, generally in the form of a reticulum, with one or more pyrenoids; cell division by the rupture of a ring-like thickening on the inner wall of each cell near the apex. Reproduction asexually by multiciliate zoospores, formed singly by the vegetative cells, germinating almost immediately. Reproduction sexually by eggs and sperms in oogonia and antheridia; oogonia single or in groups, arising as a result of division of a vegetative cell, opening by a pore or a lid, through which the sperm may pass; the egg when fertilized becomes the oospore, with a wall of one to three layers, which after a period of rest produces four zoospores. Antheridia either in the same filament as the oogonia or in separate filaments; male filaments approximately the same size as the female, or much smaller ("dwarf males"); dwarf males, arising from the germination of androspores, are epiphytic usually on or near the oogonia, sometimes scattered; sperms one or two in each antheridium.

KEY TO THE GENERA.

1. Filaments unbranched.....III. **Oedogonium** (p. 53)
1. Filaments branched..... 2
 2. Vegetative cells with laterally placed setae.... I. **Bulbochaete** (p. 27)
 2. Vegetative cells devoid of setae..... II. **Oedocladium** (p. 51)

Genus I. **Bulbochaete** Agardh

1817, p. XXIX; Pringsheim 1858, p. 71; Wittrock 1874, p. 43; Hirn 1900, p. 321; Collins 1909, p. 266; Tiffany 1928, p. 129.

Filaments usually attached, branched, the branches unilateral; vegetative cells normally widening upwards; basal cell ordinarily the only one capable of division in formation of main axis, first new cell forming a long tubular bristle with swollen bulb-like base, subsequent cells intercalated between basal cell and next one above; rarely division is intercalary; other vegetative cells potential origin for side branches; terminal cell of each branch always furnished with a bristle; oogonium arising by a double division of a vegetative cell.

KEY TO THE SPECIES, VARIETIES, AND FORMS OF
BULBOCHAETE.

1. Oogonium 24–27 μ in length.....8. *B. minuta*
1. Oogonium 27–40 μ in length.....2
1. Oogonium 40–65 (–68) μ in length.....21
1. Oogonium 65–95 μ in length.....52
1. Oogonium 104–108 μ in length.....47. *B. imperialis*
1. Oogonium not, or imperfectly, known.....59
 2. Monoecious.....3
 2. Dioecious, nannandrous.....4
3. Vegetative cells longer than broad.....1. *B. nana*
3. Vegetative cells not longer than broad.....2. *B. monile*
 4. Suffultory cells without division.....5
 4. Suffultory cells with division.....6
5. Oospore wall smooth.....33. *B. doliiformis*
5. Oospore wall ribbed.....34. *B. pygmaea*
 6. Division of suffultory cell basal.....7
 6. Division of suffultory cell inferior.....10. *B. furberae*
 6. Division of suffultory cell inframedian, median, or supramedian†.....11
 6. Division of suffultory cell superior.....16
 6. Division of suffultory cell supreme.....17a. *B. polyandria* f. *notabilis*
7. Oospore wall smooth.....8
7. Oospore wall scrobiculate.....9
 8. Oogonium 34–44 $\times$ 31–38 μ7. *B. elatior*
 8. Oogonium 31–37 $\times$ 28–37 μ7a. *B. elatior* f. *pumila*
9. Vegetative cells 13–17 μ10
9. Vegetative cells 17–20 μ9. *B. brebissonii*
 10. Antheridia exterior.....7b. *B. elatior* var. *scrobiculata*
 10. Antheridia interior.....9a. *B. brebissonii* var. *minor*
11. Oospore wall smooth.....12
11. Oospore wall scrobiculate.....13
 12. Diameter of oogonium 33–38 μ14a. *B. angulosa* f. *picta*
 12. Diameter of oogonium 37–42 μ14. *B. angulosa*
13. Vegetative cells 10–15 μ in diameter.....10. *B. furberae*
13. Vegetative cells 15–20 μ in diameter.....14
 14. Oospore wall nearly smooth to scrobiculate.....15
 14. Oospore wall scrobiculate to nearly crenulate.....11. *B. crenulata*
15. Vegetative cells 2–3 diameters long.....12. *B. intermedia*
15. Vegetative cells 3–5 diameters long.....12a. *B. intermedia* var. *depressa*
 16. Diameter of vegetative cells 10–12 μ16b. *B. nordstedtii* var. *minor*
 16. Diameter of vegetative cells 14–22 μ17
17. Vegetative cells 1 $\frac{1}{4}$ –2 diameters long.....15. *B. borealis*
17. Vegetative cells 2–5 diameters long.....18
 18. Gynandrosporous.....19
 18. Idioandrosporous.....20
19. Diameter of oogonium 28–32 μ16a. *B. nordstedtii* f. *suberecta*
19. Diameter of oogonium 36–43 μ16. *B. nordstedtii*
 20. Vegetative cells 15–20 $\times$ 45–100 μ17. *B. polyandria*
 20. Vegetative cells 17–22 $\times$ 54–88 μ17a. *B. polyandria* f. *notabilis*
21. Vegetative cells less than 12 μ in diameter.....48. *B. spirogranulata*
21. Vegetative cells more than 12 μ in diameter.....22
 22. Monoecious.....23
 22. Dioecious, nannandrous.....29
 22. Reproductive structures imperfectly known.....49. *B. brevifulta*
23. Pore of oogonium less than 5 μ wide.....24
23. Pore of oogonium more than 5 μ wide.....3. *B. megastoma*
24. Vegetative cells 1–1 $\frac{1}{4}$ diameters long.....25
24. Vegetative cells 1 $\frac{1}{4}$ –2 diameters long.....26
25. Oogonium 24–28 μ in diameter.....4. *B. basispora*
25. Oogonium 28–34 μ in diameter.....2a. *B. monile* var. *robusta*

† Some variations of *B. nordstedtii* and *B. polyandria* may be sought here.

26. Oogonium regularly ellipsoid.....5. *B. debaryana* 27
 26. Oogonium cylindric to oblong-ellipsoid.....27
 27. Length of oogonium 40-48 μ6a. *B. mirabilis* f. *immersa* 28
 27. Length of oogonium 47-58 μ28
 28. Diameter of vegetative cells 13-16 μ6b. *B. mirabilis* var. *gracilis* 28
 28. Diameter of vegetative cells 15-20 μ6. *B. mirabilis* 28
 29. Oospore wall smooth.....30
 29. Oospore wall scrobiculate.....32
 29. Oospore wall reticulate-scrobiculate.....18. *B. gigantea* 41
 29. Oospore wall ribbed.....41
 30. Division of suffultory cell basal.....19. *B. diamesandria* 31
 30. Division of suffultory cell median or nearly so.....31
 30. Division of suffultory cell superior.....22a. *B. sessilis* f. *glabra* 31
 31. Oogonium about 46 $\times$ 44 μ20. *B. elachistandria* 31
 31. Oogonium 55-64 $\times$ 43-51 μ21. *B. obliqua* 33
 32. Diameter of oogonium 40-60 μ33
 32. Diameter of oogonium 60-80 μ38
 33. Division of suffultory cell supreme.....25a. *B. dispar* var. *ripariana* 34
 33. Division of suffultory cell superior.....34
 33. Division of suffultory cell supramedian, median, or inframedian.....35
 33. Division of suffultory cell basal.....9. *B. brebissonii* 35
 34. Oogonium subquadrangular-globose.....22. *B. sessilis* 36
 34. Oogonium subdepressed-globose.....25. *B. dispar* 36
 35. Gynandrosporous.....31. *B. congener* 37
 35. Idioandrosporous.....37
 36. Diameter of oogonium 40-51 μ28. *B. crassa* 37
 36. Diameter of oogonium 52-60 μ30. *B. quadrata* 39
 37. Vegetative cells 1-2 diameters long.....13. *B. subintermedia* 39
 37. Vegetative cells 2-5 diameters long.....39
 38. Gynandrosporous.....27. *B. crassiuscula* 40
 38. Idioandrosporous.....40
 39. Diameter of oogonium 59-70 μ23. *B. setigera* 42
 39. Diameter of oogonium 70-80 μ29. *B. valida* 43
 40. Length of oogonium 48-56 μ26. *B. punctulata* 43
 40. Length of oogonium 57-60 μ35. *B. tenuis* 44
 41. Oogonium 22-27 μ in diameter.....42
 41. Oogonium 27-41 μ in diameter.....43
 42. Vegetative cells rectangular.....36. *B. repanda* 44
 42. Vegetative cells repand.....44
 42. Vegetative cells neither rectangular nor repand.....44
 43. Diameter of oogonium 28-32 μ37a. *B. rectangularis* var. *hiloensis* 45
 43. Diameter of oogonium 32-39 μ37. *B. rectangularis* 45
 44. Oogonium 39-56 μ in length.....48
 44. Oogonium 58-80 μ in length.....46
 45. Vegetative cells 13-18 μ in diameter.....38. *B. varians* 47
 45. Vegetative cells 17-22 μ in diameter.....47
 46. Dwarf male stipes 11-14 $\times$ 15-24 μ38b. *B. varians* var. *hawaiiensis* 47
 46. Dwarf male stipes 17-18 $\times$ 28-31 μ38a. *B. varians* var. *subsimpler* 49
 47. Length of oogonium 39-46 μ35a. *B. tenuis* var. *norvegica* 49
 47. Length of oogonium 47-53 μ49
 48. Oogonium fusiformly ellipsoid.....50
 48. Oogonium broadly ellipsoid to ovoid.....40. *B. rhadinospora* 50
 49. Ribs of oospore smooth.....40a. *B. rhadinospora* f. *antiqua* 51
 49. Ribs of oospore crenulate.....39. *B. lagoensis* 51
 50. Ribs of oospore coarsely serrate or crenulate.....51
 50. Ribs of oospore finely serrate or smooth.....41. *B. minor* 53
 51. Vegetative cells 18-25 $\times$ 27-50 μ41a. *B. minor* var. *germanica* 53
 51. Vegetative cells 15-22 $\times$ 22-66 μ40b. *B. rhadinospora* var. *litoralis* 54
 52. Oogonium 29-37 μ in diameter.....54
 52. Oogonium 38-45 μ in diameter.....42. *B. bullardi* 54
 52. Oogonium 41-56 μ in diameter.....58
 52. Oogonium 56-66 μ in diameter.....58
 52. Oogonium 68-92 μ in diameter.....58

53. Ribs of oospore smooth.....40. *B. rhadinospora*
 53. Ribs of oospore crenulate.....40a. *B. rhadinospora* f. *antiqua*
 54. Vegetative cells 19–27 μ in diameter.....55
 54. Vegetative cells 27–30 μ in diameter.....43. *B. denticulata*
 55. Oogonium 41–46 μ in diameter.....44. *B. affinis*
 55. Oogonium 46–56 μ in diameter.....56
 56. Vegetative cells 19–25 $\times$ 40–100 μ57
 56. Vegetative cells about 27 $\times$ 54 μ45. *B. anomala*
 57. Ribs of oospores broadly denticulate.....46. *B. insignis*
 57. Ribs of oospores reticulate-denticulate.....46a. *B. insignis* var. *reticulata*
 58. Oogonium globose.....24. *B. alabamensis*
 58. Oogonium pyriform.....32. *B. pyrum*
 58. Oogonium ellipsoid.....47a. *B. imperialis* var. *regalis*
 59. Vegetative cells blood-red in color.....50. *B. sanguinea**
 59. Vegetative cells medianly plicate.....51. *B. horrida**

1. *B. nana* Wittrock.

(Pl. IX, fig. 75.)

1872, p. 7, Pl. 1, fig. 9; 1874, p. 50; *B. nana* Witt. var. *subbasispora* Witt. 1874, p. 50; Hirn 1900, p. 349, Pl. LVII, fig. 362; Collins 1909, p. 272; Heering 1914, p. 229, fig. 349; West 1916, p. 396, fig. 251c; Tiffany 1928, p. 132, Pl. XXII, fig. 75.

Monoecious; oogonia ellipsoid, patent, below terminal setae or vegetative cells; outer wall of oospore longitudinally ribbed; antheridia 1–2†, erect (rarely patent) subepigynous or scattered; vegetative cells 10–16 $\times$ 10–22 μ ; oogonia 20–25 $\times$ 33–40 μ ; oospores 18–23 $\times$ 30–38 μ ; antheridia 7–9 $\times$ 5–9 μ .

Greenland, Alaska, Ceylon, India, Austria, England, Finland, Germany, Latvia, Sweden, Norway, Turkestan.

2. *B. monile* Wittrock and Lundell.

(Pl. IX, fig. 72.)

In Wittrock 1874, p. 50; (?) *B. nana* Witt. in Wolle 1887, p. 100; Hirn 1900, p. 348, Pl. LVII, fig. 360; Collins 1909, p. 272; Heering 1914, p. 229, fig. 348; Tiffany 1928, p. 132, Pl. XXII, fig. 72; P. B. A. No. 1432.

Monoecious; usually few celled; vegetative cells short and sometimes globose; oogonia ellipsoid, patent (rarely erect), below terminal setae or vegetative cells; outer wall of oospore longitudinally ribbed; antheridia 1–2, erect or patent, subepigynous or scattered; vegetative cells 11–16 $\times$ 10–16 μ ; oogonia 22–25 $\times$ 30–38 μ ; oospores 20–23 $\times$ 28–35 μ ; antheridia 8–10 $\times$ 6–8 μ .

United States: Massachusetts, New Jersey; Sweden, Finland, Latvia; British Columbia.

2a. Var. *robusta* Hirn.

(Pl. IX, fig. 73.)

1900, p. 349, Pl. LVII, fig. 361; Heering 1914, p. 229; Tiffany 1928, p. 133, Pl. XXII, fig. 73.

*Species of incomplete descriptions and included in the key with such characters as are available.

†See foot note, page 65.

Oogonia considerably larger and broadly ellipsoid; vegetative cells $14-19 \times 14-19 \mu$; oogonia $28-34 \times 39-45 \mu$; oospores $26-32 \times 37-42 \mu$; antheridia $8-10 \times 5-8 \mu$.

Germany.

B. nana and *B. monile*, as far as specific dimensions are concerned, are quite similar plants. If one has a mature plant of the former, however, he will find vegetative cells longer than their width, and this is almost never the case in the latter. The variety *robusta* is easily separated by its longer oogonia.

3. *B. megastoma* Wittrock and Lundell.

(Pl. IX, fig. 76.)

In Wittrock 1874, p. 51, Pl. 1, fig. 21; Hirn 1900, p. 354, Pl. LIX, fig. 370; Heering 1914, p. 231, fig. 353; Tiffany 1928, p. 133, Pl. XXII, fig. 76.

Monoecious; oogonia ellipsoid to subcylindric ellipsoid, patent (rarely erect), below terminal setae or vegetative cells, with large mouth-like pore; outer oospore wall longitudinally ribbed; antheridia 1-2, erect (rarely patent), scattered or subepigynous; vegetative cells $18-24 \times 18-35 \mu$; oogonia $32-36 \times 48-54 \mu$; oospores $30-34 \times 46-52 \mu$; antheridial cells $10-13 \times 7-9 \mu$.

Sweden, Denmark.

The large mouth-like pore of the oogonium rather readily separates mature plants from any others of the genus.

4. *B. basispora* Wittrock and Lundell.

(Pl. IX, fig. 74.)

In Wittrock 1874, p. 50; Hirn 1900, p. 350, Pl. LVIII, fig. 364; Heering 1914, p. 229, fig. 350; Tiffany 1928, p. 133, Pl. XXII, fig. 74.

Monoecious; oogonia ellipsoid or subcylindric-ellipsoid, patent or erect, below vegetative cells; outer oospore wall longitudinally ribbed; antheridia 1-3, patent or erect, scattered or subepigynous; vegetative cells $15-19 \times 15-23 \mu$; oogonia $24-28 \times 40-45 \mu$; oospores $22-25 \times 38-43 \mu$; antheridia $8-11 \times 6-8 \mu$.

United States: Ohio; Sweden.

Among the monoecious forms *B. basispora* bears some resemblance to *B. debaryana* and *B. mirabilis*, but can usually be distinguished by its relatively shorter vegetative cells and its oogonia.

5. *B. debaryana* Wittrock and Lundell.

(Pl. IX, fig. 77.)

In Wittrock 1874, p. 51; Hirn 1900, p. 353, Pl. LIX, fig. 369; Heering 1914, p. 231, fig. 352; Tiffany 1928, p. 133, Pl. XXII, fig. 77.

Monoecious; oogonia ellipsoid, patent, below vegetative cells or terminal setae; outer wall of oospore longitudinally ribbed; antheridia

1-?, patent or seldom erect, subepigynous or scattered; vegetative cells $18-22 \times 23-44\mu$; oogonia $27-33 \times 43-50\mu$; oospores $25-30 \times 40-48\mu$; antheridia $10-12 \times 8-9\mu$.

Sweden, Latvia, Canada.

6. **B. mirabilis** Wittrock.

(Pl. IX, fig. 69.)

1870, p. 137, Pl. 1, figs. 8 and 9; 1874, p. 50; *B. sp.* Reinsch 1875, p. 81; *B. mirabilis* Wittr. var. *lapponica* Wittr. and Lund. in Wittrock 1874, p. 51; Hirn 1900, p. 351, Pl. LVIII, fig. 365; Hirn 1906, p. 57; Collins 1909, p. 272; Heering 1914, p. 229, fig. 351; Tiffany 1928, p. 134, Pl. XXII, fig. 69; P. B. A. No. 1431.

Monoecious; oogonia cylindric-ellipsoid to suboblong-ellipsoid, patent (rarely erect), below terminal setae or vegetative cells; outer wall of oospore longitudinally ribbed; antheridia 1-4, erect or patent, subepigynous or scattered; vegetative cells $15-20 \times 20-40\mu$; oogonia $26-33 \times 46-58\mu$; oospores $25-31 \times 44-56\mu$; antheridia $9-12 \times 6-9\mu$.

United States: New Jersey, Michigan, Illinois, Minnesota; Greenland, Africa, Europe, Australia, Siberia, Mongolia.

6a. Form **immersa** (Wittrock) Hirn.

(Pl. IX, fig. 70.)

1900, p. 352, Pl. LVIII, fig. 367; *B. mirabilis* Wittr. var. *immersa* Wittr. 1874, p. 51; Heering 1914, p. 230; Tiffany 1928, p. 134, Pl. XXII, fig. 70.

Vegetative cells $13-17 \times 16-30\mu$; oogonia $25-33 \times 40-48\mu$; oospores $23-30 \times 38-46\mu$; antheridia $7-11 \times 6-8\mu$.

Norway, Germany.

6b. Var. **gracilis** (Pringsheim) Hirn.

(Pl. IX, fig. 71.)

1900, p. 353, Pl. LIX, fig. 368; *B. gracilis* Pringsh. 1858, p. 74; Wittrock 1874, p. 57; Heering 1914, p. 230; Tiffany 1928, p. 134, Pl. XXII, fig. 71.

Vegetative cells $13-16 \times 17-32\mu$; oogonia $20-28 \times 47-55\mu$; oospores $18-26 \times 45-53\mu$; antheridia $8-11 \times 6-8\mu$.

Germany, Sweden, Denmark.

7. **B. elatior** Pringsheim.

(Pl. I, fig. 1.)

1858, p. 73, Pl. 6, fig. 5; Wittrock 1874, p. 49; Hirn 1900, p. 321, Pl. LI, fig. 327; Hirn 1906, p. 56; Heering 1914, p. 231, fig. 355; Tiffany 1928, p. 134, Pl. XIV, fig. 1.

Diocious, nannandrous, gynandrosporous; oogonia depressed globose to depressed subquadrangular-globose, erect or seldom patent, below androsporangia; division of suffultory cell basal; oospore smooth; androsporangia epigynous or rarely scattered, 1-2; dwarf males usually on suffultory cells, antheridia exterior, unicellular, stipe scarcely longer than the antheridia; vegetative cells $13-18 \times 20-63\mu$; oogonia $34-44 \times 31-38\mu$; oospores $32-42 \times 29-36\mu$; androsporangia $10-13 \times 8-11\mu$; dwarf male stipe $8-10 \times 18-35\mu$; antheridia $6-8 \times 9-11\mu$.

United States: Massachusetts; India, Ceylon, Austria, Australia, Germany, Sweden, New Caledonia.

7a. Form **pumila** Hirn.

(Pl. I, fig. 2.)

1900, p. 322, Pl. LI, fig. 328; Tiffany 1928, p. 135, Pl. XIV, fig. 2.

Smaller; vegetative cells $10-15 \times 25-60\mu$; oogonia $31-37 \times 28-37\mu$; androsporangia $10-11 \times 9-11\mu$; dwarf male stipe $8-10 \times 19-24\mu$; antheridia $7-9 \times 6-8\mu$.

Brazil.

7b. Var. **scrobiculata** Tiffany.

(Pl. I, fig. 3.)

1928, p. 135, Pl. XIV, fig. 3.

Similar to the type except outer oospore wall is minutely scrobiculate; antheridia 1-2; vegetative cells $13-16 \times 28-40\mu$; oogonia $28-44 \times 28-36\mu$; oospores $26-42 \times 26-34\mu$; androsporangia $10-13 \times 10-16\mu$; antheridia $6-8 \times 4-7\mu$; dwarf male stipe $11-13 \times 16-20\mu$.

United States: Alabama.

This variety clearly belongs with *B. elatior* because of its exterior antheridia, although the oospore markings are similar to those of *B. brebissonii*. It is barely possible, of course, that *B. elatior* sometimes has scrobiculate oospores, although there is no evidence for it. Until such time as more is known about the variations, if any, in the oospore of the species, it seems best to catalogue the separate variety, as above.

8. **B. minuta** West and West.

(Pl. I, fig. 9.)

1902, p. 126, Pl. 17, fig. 10; Hirn 1906, p. 25, Pl. IV, fig. 28; West 1916, p. 388, fig. 242; Tiffany 1928, p. 135, Pl. XIV, fig. 9.

Dioecious, nannandrous, gynandrosporous; oogonia depressed-globose, patent or erect, below the androsporangia; division of oogonium median or nearly so; suffultory cells without division, or with division basal; oospore wall smooth; androsporangia unicellular, epigynous; dwarf males on suffultory cells, antheridia exterior, unicellular, stipe twice as long as antheridia, curved; vegetative cells $9-12 \times 18-35\mu$; oogonia $29-35 \times 24-27\mu$; oospores $27-33 \times 22-25\mu$; androsporangia $9-11 \times 6-9\mu$; dwarf males $6-7 \times 18-20\mu$.

Heneratgodha, Ceylon.

B. minuta is among the smallest of the genus, the length of its oogonium being less than that of any other known species.

9. **B. brebissonii** Kuetzing.

(Pl. I, fig. 7.)

1854, p. 19, Pl. 86, figs. B and f; *B. tumida* Witt. 1870, p. 139; 1874, p. 46; Hirn 1900, p. 323, Pl. LI, fig. 330; Collins 1909, p. 268; Heering 1914, p. 231, fig. 356; Tiffany 1928, p. 135, Pl. XIV, fig. 7.

Dioecious, nannandrous gynandrosporous; oogonia depressed-sub-quadrangular-globose, erect, below terminal setae or androsporangia;

division of the suffultory cell basal; outer oospore wall scrobiculate; androsporangia scattered or epigynous, 1-3; dwarf males on oogonia (rarely on adjacent cells), antheridia interior, stipe slightly curved, shorter than antheridia; vegetative cells $17-20 \times 50-90\mu$; oogonia $42-50 \times 37-45\mu$; oospores $40-48 \times 35-43\mu$; androsporangia $11-15 \times 12-18\mu$; dwarf males $10-12 \times 28-33\mu$.

United States: Massachusetts, Illinois, Alaska; Europe.

This species is rather readily distinguished by its usually basal division of the suffultory cells, its interior antheridium, and its erect oogonia. It is distinguished from the variety *minor* by its larger size.

9a. Var. **minor** Woronichin.

1923, p. 99; (?) *B. brebissonii* Kuetz. f. in Borge 1923, p. 30; Tiffany 1928, p. 136.

Smaller than the type; vegetative cells $13-17 \times 20-100\mu$; oogonia $35-38 \times 31-38\mu$; oospores $33-36 \times 29-36\mu$; dwarf males $8-9 \times 26-27\mu$.

Tiflis, Caucasus.

10. **B. furberae** Collins.

(Pl. I, fig. 8.)

1918a, p. 142, Pl. 124, figs. 1-5; Tiffany 1928, p. 136, Pl. XIV, fig. 8.

Dioecious, nannandrous, gynandrosporous; oogonia depressed globose, below terminal setae, patent (rarely erect); division of suffultory cell inframedian; outer wall of oospore scrobiculate; androsporangia scattered, 1-9; dwarf males on oogonia, stipe strongly curved; antheridia interior; vegetative cells $10-15 \times 30-75\mu$; oogonia $36-43 \times 27-34\mu$; oospores $34-41 \times 25-32\mu$; androsporangia $9-10 \times 8-9\mu$; dwarf males $7-8 \times 20-25\mu$.

United States: Massachusetts (Woods Hole).

This species differs from *B. elatior* in its scrobiculate spore wall and its division of suffultory cell. It is smaller than *B. brebissonii*.

11. **B. crenulata** Pringsheim.

(Pl. I, fig. 4.)

1858, p. 72, Pl. 6, fig. 4; *B. crenulata* Pringsh. var. *plena* Wittr. 1872, p. 19; Wittrock 1874, p. 45; Hirn 1900, p. 331, Pl. LIII, fig. 337; Collins 1909, p. 269; Heering 1914, p. 233, fig. 359; Tiffany 1928, p. 136, Pl. XIV, fig. 4.

Dioecious, nannandrous, gynandrosporous; oogonia subdepressed-globose, patent, below terminal setae or androsporangia, or rarely vegetative cells; division of suffultory cells median or slightly below; spore wall scrobiculate to crenulate; androsporangia epigynous or scattered, 1-5; dwarf males on or near oogonia, antheridia interior, stipe slightly curved, shorter than antheridium; vegetative cells $16-20 \times 32-70\mu$; oogonia $43-48 \times 35-43\mu$; oospores $40-46 \times 33-40\mu$; antheridia $10-15 \times 7-10\mu$; dwarf males $9-10 \times 24-26\mu$.

United States: Michigan, Illinois; Australia, Finland, Germany, Norway, Sweden, Denmark, Patagonia.

B. crenulata has often been confused with forms of *B. intermedia*, since the "crenulations" of the oospore walls are not always distinctly "crenulate." It seems best, however, to retain the two as distinct species because (1) the latter normally has longer cells in proportion to diameter and (2) the two species normally represent extremes of a series of gradations in oospore wall ornamentation: *B. intermedia* may have nearly smooth to scrobiculate oospores, while those of *B. crenulata* vary from distinctly scrobiculate to crenulate.

12. *B. intermedia* De Bary.

(Pl. I, fig. 5.)

1854, p. 72, Pl. 4, figs. 1-7; Wittrock 1874, p. 44; *B. crenulata* Pringsh. var. *supremediana* Wittr. 1883 in Wittr. and Nordst. Alg. Exs. Fasc. 11, No. 509; (?) *B. elachistandria* Wittr. in Borge 1896, p. 3; Hirn 1900, p. 326, Pl. LII, fig. 333; *B. intermedia* De Bary f. *americana* Hirn 1900, p. 328, Pl. LII, fig. 334; *B. intermedia* De Bary f. *supramediana* (Wittr.) Hirn 1900, p. 328, Pl. LII, fig. 335; Hirn 1906, p. 56; Collins 1909, p. 268, fig. 82; Heering 1914, p. 233, fig. 358; Tiffany 1926, p. 109, Pl. X, fig. 107; 1928, p. 137, Pl. XIV, fig. 5; P. B. A. No. 973.

Diocious, nannandrous, gynandrosporous; oogonia subdepressed-globose, patent, below androsporangia; division of suffultory cells nearly median; outer wall of oospore scrobiculate, rarely apparently smooth; androsporangia 1-2, epigynous or rarely scattered; dwarf males on the oogonium; antheridium interior, stipe slightly curved, shorter than the antheridium; vegetative cells $17-20 \times 35-70\mu$; oogonia $40-48 \times 31-40\mu$; oospores $38-46 \times 30-38\mu$; androsporangial cells $11-13 \times 7-12\mu$; dwarf males $9-10 \times 21-26\mu$.

United States: Iowa, Illinois, Ohio, Michigan, Connecticut, Pennsylvania, Massachusetts, Alaska; Canada, Australia, Europe, Greenland, Africa.

12a. Var. *depressa* Wittrock.

(Pl. I, fig. 6.)

1874, p. 44, Pl. 1, fig. 18; Hirn 1900, p. 329, Pl. LII, fig. 336; *B. intermedia* De Bary var. *depressa* Hirn in Heering 1914, p. 233; Tiffany 1926, p. 109; 1928, p. 137, Pl. XIV, fig. 6.

Vegetative cells a little longer than in the type; oogonia depressed-globose; outer oospore wall scrobiculate, often nearly smooth; division of suffultory cell a little above the median position (rarely a little below median); vegetative cells $14-19 \times 35-88\mu$; oogonia $42-46 \times 30-40\mu$; oospores $40-44 \times 28-38\mu$; androsporangial cells $11-13 \times 9-12\mu$; dwarf males $9-11 \times 22-25\mu$.

United States: Iowa; Finland, Germany, Norway, Sweden; British Columbia.

Hirn recognized with *B. intermedia* its variety *depressa* and two forms, *americana* Hirn and *supramediana* (Wittrock)

Hirn. I have seen a great number of *intermedia* plants from the North Central states, and judging from this American material and from original descriptions, it seems almost impossible to recognize with certainty either of the two forms above; the variety is probably distinct. Occasionally individual plants and more often parts of filaments are found with slight differences and variations in form and shape of vegetative cells and oogonia. Frequently all variations between the type and the two forms can be noted in one plant. The variety *depressa* with relatively longer cells in proportion to diameter is perhaps sufficiently distinct to recognize. It does not appear to be to the best interests of taxonomic nomenclature in this species to do little more than recognize its variability. I have, therefore, included with the species proper the two forms *americana* and *supramediana*. The relationship to *B. crenulata* is noted above, under that species.

13. **B. subintermedia** Elfving.

(Pl. II, fig. 15.)

In Hirn 1895, p. 8; Hirn 1900, p. 332, Pl. LIII, fig. 338; Heering 1914, p. 233, fig. 360; West 1916, p. 396, fig. 251A; Tiffany 1928, p. 138, Pl. XV, fig. 15.

Dioecious, nannandrous, gynandrosporous; oogonia subdepressed-globose, patent, below terminal setae, androsporangia, or rarely vegetative cells; division of suffultory cell submedian; outer spore wall scrobiculate; androsporangia 1-?, epigynous or scattered; dwarf males on oogonia, stipe curved or nearly erect, shorter than the oogonia; vegetative cells $17-22 \times 34-100\mu$; oogonia $44-51 \times 39-44\mu$; oospores $42-48 \times 37-41\mu$; androsporangia $13-16 \times 9-10\mu$; dwarf males $10-12 \times 25-30\mu$.

Finland, Great Britain.

14. **B. angulosa** Wittrock and Lundell.

(Pl. II, fig. 14.)

In Wittrock 1874, p. 45; (?) *B. elachistandria* Wittr. in Wolle 1887, p. 97; Hirn 1900, p. 336, Pl. LIV, fig. 346; Heering 1914, p. 235, fig. 364; Tiffany 1928, p. 138, Pl. XV, fig. 14.

Dioecious, nannandrous, gynandrosporous; oogonia biconically and angularly globose, apically truncate, below terminal setae or rarely androsporangia; division of suffultory cells a little above median position; oospore wall smooth; androsporangia scattered or epigynous, 1-3; dwarf males on oogonia, antheridia interior, stipe somewhat curved; vegetative cells $13-18 \times 20-45\mu$; oogonia $36-42 \times 33-39\mu$; oospores $34-40 \times 30-36\mu$; androsporangia $10-11 \times 9-10\mu$; dwarf males $8-9 \times 18-21\mu$.

United States: Pennsylvania; England, Sweden, Finland, Africa.

14a. Form **picta** Hirn.

1900, p. 337.

Of smaller dimensions; vegetative cells $13-16 \times 19-40\mu$; oogonia $33-38 \times 31-37\mu$.

Brazil.

15. **B. borealis** Wittrock.

(Pl. II, fig. 10.)

1870, p. 138; 1874, p. 46; Hirn 1900, p. 324, Pl. LI, fig. 331; Heering 1914, p. 233, fig. 357; Tiffany 1928, p. 139, Pl. XV, fig. 10.

Dioecious, nannandrous, gynandrosporous; oogonia subdepressed-globose, patent, below vegetative cells, wall thick; division of suffultory cell superior (occasionally nearly supreme); outer oospore wall scrobiculate or sometimes nearly smooth; androsporangia 1-3, subepigynous, patent; dwarf males on oogonia, antheridia interior, stipe a little curved, shorter than antheridia; vegetative cells $16-21 \times 20-42\mu$; oogonia $40-48 \times 35-40\mu$; oospores $38-46 \times 33-38\mu$; androsporangia $12-13 \times 8-9\mu$; dwarf males $9-10 \times 18-21\mu$.

Norway, Sweden, Finland.

16. **B. nordstedtii** Wittrock.

(Pl. II, fig. 17.)

1874, p. 44; Hirn 1900, p. 332, Pl. LIII, fig. 339; Collins 1909, p. 269; Heering 1914, p. 234, fig. 361; West 1916, p. 396, fig. 251B; Tiffany 1928, p. 139, Pl. XV, fig. 17; P. B. A. No. 717.

Dioecious, nannandrous, gynandrosporous; oogonia depressed-globose or subquadrangularly so, patent, below androsporangia or rarely terminal setae; division of suffultory cell superior (occasionally submedian); oospore wall finely scrobiculate or nearly smooth; androsporangia unicellular, epigynous; dwarf males on oogonia, stipe slightly curved, shorter than antheridia; vegetative cells $14-18 \times 28-85\mu$; oogonia $36-43 \times 29-36\mu$; oospores $34-41 \times 27-34\mu$; androsporangia $10-12 \times 9-12\mu$; dwarf males $9-19 \times 23-25\mu$.

United States: Ohio, Illinois, Connecticut, Alaska; Greenland, Australia, Finland, Germany, Spain, Sweden, Austria, Ireland.

16a. Form **suberecta** Collins.

(Pl. II, fig. 18.)

1912, p. 88; Tiffany 1928, p. 139, Pl. XV, fig. 18.

Smaller; oogonia $28-32 \times 34-38\mu$, depressed-globose, often erect.

United States: Massachusetts.

The suberect habit of this form appears to be very inconstant, but the smaller dimensions warrant a separate form.

16b. Var. **minor** Woronichin.

1923, p. 99.

Vegetative cells $10-12 \times 20-25\mu$; oospores about $36 \times 33\mu$; dwarf males $8-10 \times 19-20\mu$.

Tiflis (Caucasus).

17. **B. polyandria** Cleve.

(Pl. II, figs. 11 and 12.)

In Wittrock 1870, p. 140; *B. polyandra* Cleve in Wittrock 1874, p. 46; Hirn 1900, p. 334, Pl. LIV, fig. 342; Collins 1909, p. 270; Heering 1914, p. 234, fig. 362; Tiffany 1928, p. 140, Pl. XV, figs. 11 and 12; P. B. A. No. 1682.

Dioecious, nannandrous, idioandrosporous; oogonia subdepressed-globose, patent, below terminal setae or vegetative cells; division of suffultory cell superior (rarely submedian); outer oospore wall scrobiculate or nearly smooth; androsporangia to 10-seriate; dwarf males on oogonia, antheridia interior, stipe slightly curved, shorter than antheridia; vegetative cells $15-20 \times 45-100\mu$; oogonia $39-46 \times 32-42\mu$; oospores $37-44 \times 30-40\mu$; androsporangia $12-14 \times 11-15\mu$; dwarf males $8-9 \times 23-26\mu$.

United States: Massachusetts, Florida; Brazil, Europe.

17a. Form **notabilis** Hirn.

(Pl. II, fig. 13.)

1900, p. 334, Pl. LIV, fig. 343; Tiffany 1928, p. 140, Pl. XV, fig. 13.

Division of suffultory cell often supreme; vegetative cells $17-22 \times 51-88\mu$; oogonia $41-46 \times 36-43\mu$; oospores $39-44 \times 34-41\mu$; androsporangia $12-14 \times 12-15\mu$; dwarf males $8-10 \times 23-25\mu$.

Great Britain.

18. **B. gigantea** Pringsheim.

(Pl. V, figs. 35 and 36.)

1858, p. 71, Pl. 6, fig. 1; Wittrock 1874, p. 48; Hirn 1900, p. 347, Pl. LVII, fig. 359; Collins 1909, p. 271; Heering 1914, p. 239, fig. 373; Tiffany 1926, p. 109, Pl. X, figs. 108-110; Tiffany 1928, p. 140, Pl. XVIII, figs. 35 and 36.

Dioecious, nannandrous, idioandrosporous; oogonia subdepressed globose or rarely depressed oboviform-globose, patent, below terminal setae, rarely below vegetative cells; division of suffultory cell slightly below median; outer wall of oospore reticulate-scrobiculate; androsporangia 1-5; dwarf males slightly longer than the oogonia on which they develop; antheridia interior; stipe about twice as long as antheridium, curved; vegetative cells $24-32 \times 50-112\mu$; oogonia $60-70 \times 50-58\mu$; oospores $58-68 \times 48-56\mu$; androsporangial cells $18-20 \times 10-14\mu$; dwarf male stipe $10-13 \times 28-45\mu$; antheridial cell $13-14 \times 20-30\mu$.

United States: Michigan, Florida, Iowa, Pennsylvania; Australia, Europe.

19. **B. diamesandria** Nordstedt and Hirn.

(Pl. II, fig. 16.)

In Hirn 1900, p. 323, Pl. LI, fig. 329; Tiffany 1928, p. 140, Pl. XV, fig. 16.

Dioecious, nannandrous, gynandrosporous; oogonia subdepressed-globose, erect, below terminal setae or vegetative cells; division of suffultory cell basal; oospore wall smooth; androsporangia scattered or subepigynous, 1-?; dwarf males on suffultory cells (rarely on oogonia), antheridia unicellular; vegetative cells $17-23 \times 42-80\mu$; oogonia $48-54 \times 45-52\mu$; oospores $45-53 \times 43-50\mu$; androsporangia $14-17 \times 10-15\mu$; dwarf male stipe $11-13 \times 15-21\mu$; antheridia $10-12 \times 14-18\mu$.

Bengal, Asia.

20. **B. elachistandria** Wittrock.

(Pl. III, fig. 21.)

1874, p. 43; *B. intermedia* Pringsh. 1858, p. 72, Pl. IV, figs. 15-23; Hirn 1900, p. 342, Pl. LVI, fig. 354; Heering 1914, p. 231, fig. 354; Tiffany 1928, p. 141, Pl. XVI, fig. 21.

Dioecious, nannandrous, gynandrosporous; oogonia globose, patent, below androsporangia; division of suffultory cell nearly median; oospore wall smooth (?); androsporangia epigynous or scattered, 1-2; dwarf males on oogonia, small and unicellular; vegetative cells $20-24 \times 30-60\mu$; oogonia about $46 \times 44\mu$; androsporangia $14-16 \times 10-11\mu$; dwarf males $9 \times 19\mu$.

Germany.

21. **B. obliqua** Lundell.

(Pl. III, fig. 26.)

In Hirn 1900, p. 344, Pl. LVI, fig. 356; Heering 1914, p. 238, fig. 371; Tiffany 1928, p. 141, Pl. XVI, fig. 26.

Dioecious, nannandrous, gynandrosporous and idioandrosporous; oogonia depressed-globose, patent, below terminal setae or (rarely) androsporangia; division of suffultory cell median or nearly so; oospore wall smooth; androsporangia 1-?, epigynous; dwarf males on oogonia, antheridia interior; vegetative cells $21-27 \times 42-108\mu$; oogonia $55-64 \times 43-51\mu$; oospores $53-62 \times 40-49\mu$; androsporangia $16-21 \times 7-10\mu$; dwarf male stipes $9-10 \times 25-36\mu$; antheridia $10-12 \times 15-21\mu$.

United States: Michigan; Austria, Sweden.

22. **B. sessilis** Wittrock.

(Pl. III, fig. 19.)

1872, p. 18, Pl. I, fig. 2; 1874, p. 47; Hirn 1900, p. 338, Pl. LV, fig. 349; Heering 1914, p. 236, fig. 366; Tiffany 1928, p. 141, Pl. XVI, fig. 19.

Dioecious, nannandrous, gynandrosporous; oogonia subquad-rangularly globose, patent, below terminal setae or occasionally below vegetative cells; division of suffultory cell superior; outer oospore wall scrobiculate; androsporangia scattered, 1-3; dwarf males on oogonia or rarely suffultory cells, antheridia interior, stipe a little curved, shorter than antheridia; vegetative cells $19-22 \times 38-72\mu$; oogonia $51-56 \times 44-50\mu$; oospores $48-54 \times 42-48\mu$; androsporangia $10-13 \times 10-12\mu$; dwarf males $9-11 \times 26-29\mu$.

France, Sweden, Canada.

22a. Form **glabra** Hirn.

(Pl. III, fig. 20.)

1900, p. 339, Pl. LV, fig. 350; Heering 1914, p. 236; Tiffany 1928, p. 141, Pl. XVI, fig. 20.

Smaller, with oospore wall smooth; vegetative cells $16-21 \times 32-75\mu$; oogonia $44-51 \times 40-45\mu$; oospores $42-48 \times 38-43\mu$; androsporangia $10-12 \times 10-12\mu$; dwarf males $9-11 \times 23-27\mu$.

Sweden.

23. **B. setigera** (Roth) C. A. Agardh.

(Pl. V, fig. 37.)

1817, p. 71; *Conferva setigera* Roth 1806, p. 283; *B. setigera* C. A. Agardh 1817, p. 71; Wittrock 1874, p. 47; (?) *B. setigera* var. *Canbyii* Wood in Wolle 1887, p. 99; (?) *B. Canbyii*, Wood 1872, p. 202; Hirn 1900, p. 339, Pl. LV, fig. 351; Collins 1909, p. 271; Heering 1914, p. 236, fig. 367; Tiffany 1928, p. 142, Pl. XVIII, fig. 37.

Dioecious, nannandrous, gynandrosporous; oogonia subdepressed or depressed (quadrangularly) globose, patent, generally below terminal setae, more rarely below androsporangia or vegetative cells; division of suffultory cell slightly above median or rarely superior; outer oospore wall scrobiculate; androsporangia scattered or more rarely epigynous, 1-3; dwarf males on oogonia or near it; antheridia interior, stipe curved, shorter than antheridium; vegetative cells $25-28 \times 62-140\mu$; oogonia $70-80 \times 56-65\mu$; oospores $67-77 \times 53-62\mu$; androsporangia $16-20 \times 10-18\mu$; dwarf males $11-14 \times 30-36\mu$.

United States: Connecticut, New Jersey, Florida, Alabama, South Carolina, Michigan; Europe, South America; Africa.

24. **B. alabamensis** Transeau and Brown.

(Pl. V, fig. 38.)

In Tiffany 1928, p. 142, Pl. XVIII, fig. 38.

Dioecious, nannandrous, gynandrosporous; oogonia globose, depressed globose, or rarely angularly globose, patent; division of suffultory cells superior; outer spore wall scrobiculate; androsporangia hypogynous or epigynous or rarely scattered; dwarf males, a little shorter than oogonia, on vegetative cells, antheridia interior, stipe a little curved; vegetative cells $22-40 \times 62-111\mu$; oogonia $77-92 \times 70-88\mu$; oospores $75-90 \times 68-86\mu$; dwarf male stipes $14-18 \times 35-38\mu$; antheridia $14-15 \times 15-16\mu$.

United States: Alabama.

In form and general appearance this species is near *B. setigera*. It is larger, however, in all dimensions and is thus the largest globose species of the genus.

25. **B. dispar** Wittrock.

(Pl. III, fig. 23.)

In Wittrock and Nordstedt Exs. 1882; Hirn 1900, p. 335, Pl. LIV, fig. 344; Collins 1909, p. 270; Heering 1914, p. 234, fig. 363; Tiffany 1928, p. 142, Pl. XVI, fig. 23.

Dioecious, nannandrous, gynandrosporous and idioandrosporous; oogonia subdepressed globose, patent, below terminal setae or rarely vegetative cells; division of suffultory cell superior (rarely submedian); outer spore wall thick, finely scrobiculate; androsporangia scattered, 1-2; dwarf males on oogonia, antheridia interior, stipe a little curved, shorter than antheridia; vegetative cells $16-21 \times 32-95\mu$; oogonia $44-56 \times 42-51\mu$; oospores $40-54 \times 38-48\mu$; androsporangia $12-16 \times 10-12\mu$; dwarf males $9-11 \times 23-26\mu$.

Sweden, Greenland.

25a. *Var. ripartiana* Wittrock.

(Pl. III, figs. 24 and 25.)

In Hirn 1900, p. 335, Pl. LIV, fig. 345; Tiffany 1928, p. 143, Pl. XVI, figs. 24 and 25.

Idioandrosporous; division of suffultory cell supreme; dwarf males on or near oogonia; vegetative cells $15-21 \times 45-105\mu$; oogonia $48-54 \times 41-50\mu$; oospores $46-53 \times 39-48\mu$; androsporangia $12-16 \times 11-14\mu$; dwarf males $10-12 \times 23-25\mu$.

France.

26. *B. punctulata* (Nordstedt) Hirn.

(Pl. IV, fig. 31.)

1900, p. 326, Pl. LI, fig. 332; *B. setigera* (Roth) C. A. Agardh var. *punctulata* Nordstedt 1888, p. 9; Tiffany 1928, p. 143, Pl. XVII, fig. 31.

Dioecious, nannandrous, gynandrosporous; oogonia subdepressed globose, patent, below terminal setae or androsporangia, wall thickened, division median, narrow but distinct; division of suffultory cell superior; outer oospore wall scrobiculate; androsporangia epigynous, 1-?; dwarf males on oogonia, antheridia interior, stipe a little curved, shorter than the antheridium; vegetative cells $22-26 \times 66-104\mu$; oogonia $64-68 \times 57-60\mu$; oospores $60-66 \times 53-58\mu$; androsporangia $18 \times 11\mu$; dwarf males $11-12 \times 30-34\mu$.

New Zealand.

27. *B. crassiuscula* Nordstedt.

(Pl. IV, figs. 28 and 29.)

1877, p. 30, Pl. 3, figs. 14 and 15; *B. setigera* (Roth) C. A. Agardh *in* Moebius 1894, p. 315; (?) *B. ellipsospora* West 1899, p. 54; Hirn 1900, p. 341, Pl. LV, fig. 352; Collins 1909, p. 271; Heering 1914, p. 236, fig. 368; Tiffany 1928, p. 143, Pl. XVII, figs. 28 and 29.

Dioecious, nannandrous, idioandrosporous; oogonia depressed (sub-quadrangularly) globose, patent, below terminal setae or occasionally vegetative cells; division of suffultory cells superior (rarely submedian); outer oospore wall scrobiculate; androsporangia 1-4; dwarf males on oogonium or near it; antheridia interior, stipe slightly curved, shorter than antheridium; vegetative cells $22-27 \times 55-148\mu$; oogonia $60-78 \times 50-62\mu$; oospores $58-76 \times 48-60\mu$; antheridia $16-19 \times 10-13\mu$; dwarf males $12-14 \times 30-34\mu$.

United States: Illinois, Massachusetts, Michigan; Canada, Greenland, Europe, Australia, Ceylon.

28. *B. crassa* Pringsheim.

(Pl. IV, fig. 32.)

1855, Pl. 1, fig. 29; Wittrock 1874, p. 48; Hirn 1900, p. 343, Pl. LVI, fig. 355; Heering 1914, p. 238, fig. 370; Tiffany 1928, p. 143, Pl. XVII, fig. 32.

Dioecious, nannandrous, gynandrosporous; oogonia subdepressed globose to globose, patent, below terminal setae or rarely vegetative cells; division of suffultory cell median or a little above; outer oospore wall scrobiculate; androsporangia scattered, up to 4-seriate; dwarf males a little longer than oogonia, antheridia exterior, unicellular, stipe about twice as long as antheridium; vegetative cells $22-26 \times 44-60\mu$;

oogonia $52-60 \times 42-51 \mu$; oospores $50-58 \times 40-48 \mu$; androsporangia $13-15 \times 14-16 \mu$; dwarf male stipes $9-11 \times 32-44 \mu$; antheridia $8-9 \times 21-24 \mu$.

United States: Ohio; Germany, Austria, Switzerland.

The outer oospore wall of this species is recorded by Hirn as doubtfully smooth. Material collected in Ohio in 1920 showed the wall to be scrobiculate, and the description is emended accordingly.

29. **B. valida** Wittrock.

(Pl. IV, fig. 27.)

1872, p. 17, Pl. 1, fig. 1; 1874, p. 48; Hirn 1900, p. 346, Pl. LVI, fig. 357; Heering 1914, p. 238, fig. 372; Tiffany 1928, p. 144, Pl. XVII, fig. 27.

Dioecious, nannandrous, gynandrosporous; oogonia depressed globose, patent, below terminal setae; division of suffultory cell about median; outer oospore wall scrobiculate; androsporangia scattered, 1-3; dwarf males on oogonia, antheridia interior, stipe a little longer than antheridium, curved; vegetative cells $23-27 \times 46-95 \mu$; oogonia $59-70 \times 48-56 \mu$; oospores $57-68 \times 46-54 \mu$; androsporangia $18-20 \times 10-14 \mu$; dwarf male stipes $9-10 \times 23-29$ (-40) μ ; antheridia $11-12 \times 20-22 \mu$.

United States: Michigan; Finland, Sweden.

30. **B. quadrata** Wittrock.

(Pl. III, fig. 22.)

1872, p. 19, Pl. 1, fig. 3; 1874, p. 45; Hirn 1900, p. 338, Pl. LV, fig. 348; Heering 1914, p. 236, fig. 365; Tiffany 1928, p. 144, Pl. XVI, fig. 22.

Dioecious, nannandrous, gynandrosporous; oogonia subdepressed (subquadrangularly) globose, patent, below androsporangia; division of suffultory cell inframedian (rarely nearly median); outer oospore wall scrobiculate; androsporangia 1-2, epigynous, rarely scattered; dwarf males on oogonia, antheridia interior, stipe a little curved, shorter than antheridium; vegetative cells $19-25 \times 30-50 \mu$; oogonia $40-50 \times 40-45 \mu$; oospores $38-48 \times 38-43 \mu$; androsporangia $15-17 \times 11-12 \mu$; dwarf males $10-12 \times 27-32 \mu$.

United States: Ohio; Finland, Sweden.

31. **B. congener** Hirn.

(Pl. V, figs. 33 and 34.)

1900, p. 346, Pl. LVII, fig. 358; Tiffany 1926, p. 108, Pl. X, figs. 111 and 112; Tiffany 1928, p. 144, Pl. XVIII, figs. 33 and 34.

Dioecious, nannandrous, idioandrosporous; oogonia depressed-globose or more rarely oboviform-globose, patent, below terminal setae; division of suffultory cell inframedian; outer oospore wall scrobiculate; androsporangia 1-4; dwarf males a little shorter than the oogonia on which they develop; antheridia interior, stipe about twice as long as the antheridial cell, curved; vegetative cells $21-24 \times 40-75 \mu$; oogonia $44-54 \times 40-48 \mu$; oospores $42-52 \times 38-46 \mu$; androsporangial cells $13-15 \times 10-13 \mu$; dwarf males $9-10 \times 29-34 \mu$; antheridial cells $10-11 \times 14-17 \mu$.

United States: Iowa; South America.

32. *B. pyrulum* Lundell.

(Pl. IV, fig. 30.)

In Hirn 1900, p. 342, Pl. LV and LVI, fig. 353; Heering 1914, p. 238, fig. 369; Tiffany 1928, p. 145, Pl. XVII, fig. 30.

Dioecious, nannandrous, (?) idioandrosporous; oogonium pyriform, patent, below terminal setae, wall thickened; division of suffultory cells submedian; outer oospore wall finely scrobiculate; dwarf males near or on oogonia, antheridia interior, stipe a little curved, shorter than antheridium; vegetative cells $23-29 \times 64-178\mu$; oogonia $(58-)$ $68-85 \times 64-79\mu$; oospores $65-83 \times 61-77\mu$; dwarf males $11-14 \times 29-33\mu$.

Austria, Sweden, Finland.

33. *B. doliiformis* Borge.

(Pl. VI, figs. 39 and 40.)

1925, p. 11, Pl. I, fig. 41; Tiffany 1928, p. 145, Pl. XIX, figs. 39 and 40.

Dioecious, nannandrous; oogonia ellipsoid, erect, below terminal setae; suffultory cell without division; oospores ellipsoid, not quite filling oogonia, wall smooth; dwarf males on vegetative cells, antheridia exterior, 2; vegetative cells $11-15 \times 11-16\mu$; oogonia $23-25 \times 26-33\mu$; oospores $22-23 \times 27-28\mu$; dwarf male stipe $10-12 \times 10-12\mu$; antheridia $7-8 \times 7-8\mu$.

Argentina.

This species is near *B. pygmaea* in size and shape of cells and in having suffultory cells without division. The plant is few celled with the vegetative cells convex, as seen laterally, sometimes with a broad median constriction, and the oospore wall is smooth.

34. *B. pygmaea* Pringsheim; Wittrock.

(Pl. VI, fig. 41.)

In Wittrock 1870, p. 141; *B. pygmaea* b. *minor* Pringsh. 1858, p. 74; *B. pygmaea* Wittr. 1870, p. 141; Wittrock 1874, p. 52; Hirn 1900, p. 356, Pl. LIX, fig. 372; Collins 1909, p. 273; Heering 1914, p. 239, fig. 374; Tiffany 1928, p. 145, Pl. XIX, fig. 41.

Dioecious, nannandrous, gynandrosporous; oogonia ellipsoid, patent, below terminal setae or vegetative cells; outer wall of oospore longitudinally ribbed; suffultory cells without division; androsporangia subepigynous or scattered, 1-? celled; dwarf males near oogonia, antheridia exterior, 1-3; vegetative cells $11-15 \times 8-15\mu$; oogonia $22-25 \times 32-40\mu$; oospores $20-23 \times 30-38\mu$; androsporangia $7-10 \times 6-9\mu$; dwarf male stipes $11-12 \times 15-19\mu$; antheridia $7-8 \times 7-8\mu$.

United States: Massachusetts; Europe, Brazil.

35. *B. tenuis* (Wittrock) Hirn.

(Pl. VII, fig. 59.)

1900, p. 368, Pl. LXIII, fig. 388; *B. rectangularis* Wittr. var. *tenuis* Wittr. 1874, p. 56; *B. rectangularis* Wittr. var. *norvegica* Wittr. 1874, p. 56; *B. tenuis* (Wittr.) Hirn var. *norvegica* (Wittr.) Hirn 1900, p. 369 (not 1906, p. 60); Collins 1912, p. 89; Heering 1914, p. 242, fig. 381; Tiffany 1928, p. 146, Pl. XX, fig. 59.

Dioecious, nannandrous, gynandrosporous; oogonia suboblong-ellipsoid, erect or patent, below terminal setae or androsporangia; outer oospore wall longitudinally ribbed; androsporangia epigynous or more rarely scattered, 1-?; dwarf males on or near oogonia; antheridia exterior, 1-3; vegetative cells $13-16 \times 20-40\mu$; oogonia $22-26 \times 42-48\mu$; oospores $20-24 \times 40-46\mu$; androsporangia $10-12 \times 13-19\mu$; dwarf male stipes $12-14 \times 18-24\mu$; antheridia $7-9 \times 6-7\mu$.

United States: Massachusetts; Northern Europe.

35a. Var. **norvegica** (Wittrock) Hirn.

(Pl. VII, fig. 60.)

1906, p. 60; Tiffany 1928, p. 146, Pl. XX, fig. 60.

Larger vegetative cells and oogonia; vegetative cells $15-18 \times 22-36\mu$; oogonia $27-34 \times 47-53\mu$; oospores $25-32 \times 45-51\mu$; dwarf male stipes $12-14 \times 19-23\mu$; antheridia $8-9 \times 8\mu$.

Finland.

The dimensions of the variety *norvegica* as given by Hirn in his monograph (1900, p. 369) are so nearly the same as those of the species proper as to make its varietal position scarcely tenable. Material collected by Silfvenius in Finland in 1903 is labelled variety *norvegica* by Hirn (1906) having the dimensions given above. It seems best, therefore, to consider the variety as having the dimensions as recorded for the Finnish material, and to regard the former description as included in the species proper.

36. **B. repanda** Wittrock.

(Pl. VI, fig. 47.)

1874, p. 55; *B. rectangularis* Wittr. var. *Lundellii* Wittr. 1874, p. 56; (?) *B. rhadinospora* Wittr. in Wolle 1887, p. 103; Hirn 1900, p. 363, Pl. LXI, fig. 380; Collins 1909, p. 274; Heering 1914, p. 240, fig. 378; Tiffany 1928, p. 146, Pl. XIX, fig. 47.

Dioecious, nannandrous, gynandrosporous; oogonia suboblong-ellipsoid, patent or erect, below androsporangia, terminal setae, or vegetative cells; outer oospore wall longitudinally ribbed; androsporangia epigynous or subepigynous, 1-?; dwarf males near or on oogonia; antheridia exterior, 1-3; vegetative cells, frequently repand, $12-17 \times 24-60\mu$; oogonia $26-36 \times 43-58\mu$; oospores $21-33 \times 40-50\mu$; androsporangia $13-15 \times 16-21\mu$; dwarf male stipes $11-15 \times 21-27\mu$; antheridia $7-10 \times 5-7\mu$.

United States: Massachusetts, Maine, New Jersey, Florida; Greenland, Finland, Norway, Sweden, Germany.

The peculiar repand appearance of the vegetative cells usually makes the identification of this species easy. Occasionally a single plant may lack this vegetative character and may thus be confused with *B. rectangularis* or *B. varians*.

37. *B. rectangularis* Wittrock.

(Pl. VI, fig. 43.)

1870, p. 142; 1874, p. 55; (?) *B. ignota* Wood 1872, p. 201; Hirn 1900, p. 359, Pl. LX, fig. 376; Collins 1909, p. 273; Heering 1914, p. 239, fig. 376; Tiffany 1928, p. 147, Pl. XIX, fig. 43; P. B. A. No. 516.

Diocious, nannandrous, gynandrosporous; oogonia ellipsoid, patent or more rarely erect, below terminal setae or androsporangia or more rarely vegetative cells; outer oospore wall longitudinally ribbed; androsporangia scattered or epigynous, 1-?; dwarf males near or occasionally on oogonia; antheridia exterior, 1-4; vegetative cells, subrectangular in cross section, $16-23 \times 20-46\mu$; oogonia $32-39 \times 45-63\mu$; oospores $29-37 \times 43-61\mu$; androsporangia $13-16 \times 10-27\mu$; dwarf male stipes $14-18 \times 22-27\mu$; antheridia $8-10 \times 5-7\mu$.

United States: Pennsylvania, Ohio, Illinois, Michigan, Massachusetts, Rhode Island, Connecticut; Europe, Patagonia, Mongolia.

This species is usually characterized by its long and few branches and its nearly rectangular cells in cross section.

37a. Var. *hiloensis* Nordstedt.

(Pl. VI, fig. 44.)

1878, p. 22; Hirn 1900, p. 361, Pl. LX, fig. 377; Collins 1909, p. 274; Tiffany 1928, p. 147, Pl. XIX, fig. 44.

Smaller, androsporangia generally epigynous; vegetative cells $14-19 \times 24-48\mu$; oogonia $28-32 \times 47-51\mu$; androsporangia $12-14 \times 13-16\mu$; dwarf male stipes $13-14 \times 22-24\mu$; antheridia $8-9 \times 5-7\mu$.

Australia; (?) Pennsylvania (U. S. A.). The only American record is that of Wolle.

38. *B. varians* Wittrock.

(Pl. VI, fig. 48.)

1870, p. 143; 1874, p. 53; Hirn 1900, p. 357, Pl. LIX, fig. 373; Heering 1914, p. 239, fig. 375; Tiffany 1926, p. 109, Pl. X, fig. 113; Tiffany 1928, p. 147, Pl. XIX, fig. 48.

Diocious, nannandrous, gynandrosporous; oogonia ovoid, patent or erect, below terminal setae or below androsporangial cells; outer oospore wall longitudinally ribbed, ribs serrate; androsporangia scattered, epigynous or hypogynous, 1-2; dwarf males on or near the oogonia, antheridia exterior, 1-3; vegetative cells $17-22 \times 22-33\mu$; oogonia $30-36 \times 44-54\mu$; oospores $28-34 \times 42-52\mu$; androsporangial cells $14-17 \times 14-18\mu$; dwarf male stipes $14-16 \times 24-27\mu$; antheridial cells $8-10 \times 6-7\mu$.

United States: Iowa, Ohio, Illinois, Michigan, Alabama, Mississippi, Indiana; Canada, Europe.

38a. Var. *subsimplex* (Wittrock) Hirn.

(Pl. VI, fig. 49.)

1900, p. 357, Pl. LIX and LX, fig. 374; *B. pygmaea* a. *major* Pringsh. 1858, p. 74; *B. subsimplex* Wittr. 1870, p. 142; 1874, p. 52; *B. varians* Wittr. var. *alpina* Wittr. and Lund. in Wittrock 1874, p. 53; *B. dumosa* Wood 1872, p. 202; *B. reticulata* Nordst. var. *minor* Lemmermann 1895, p. 25; Collins 1909, p. 273; Heering 1914, p. 239; Tiffany 1926, p. 110; 1928, p. 147, Pl. XIX, fig. 49.

Smaller in nearly all parts; ribs of oospore serrulate or smooth; vegetative cells $13-18 \times 16-34\mu$; oogonia $26-30 \times 39-46\mu$; oospores $24-28 \times 37-44\mu$; androsporangial cells $10-14 \times 7-16\mu$; dwarf male stipes $11-14 \times 15-24\mu$; antheridial cells $7-8 \times 5-7\mu$.

United States: Iowa, Ohio, Mississippi, Illinois, Pennsylvania; Brazil, India, Europe, Australia; British Columbia.

38b. Var. **hawaiiensis** Nordstedt.

(Pl. VI, fig. 50.)

1878, p. 21; Hirn 1900, p. 358, Pl. LX, fig. 375; Tiffany 1928, p. 148, Pl. XIX, fig. 50.

Vegetative cells $13-18 \times 17-36\mu$; oogonia $27-30 \times 44-54\mu$; androsporangia $12-16 \times 13-14\mu$; dwarf male stipes $17-18 \times 28-31\mu$; antheridia $10-11 \times 6-7\mu$.

United States: Ohio, Illinois; Australia.

39. **B. lagoensis** Wittrock.

(Pl. VI, fig. 46.)

1874, p. 53; Hirn 1900, p. 361, Pl. LXI, fig. 378; Tiffany 1928, p. 148, Pl. XIX, fig. 46.

Diocious, nannandrous; oogonia ellipsoid, erect or rarely patent, below vegetative cells; outer oospore wall longitudinally ribbed, the ribs dentate and connected by transverse lines; dwarf males near oogonia; antheridia exterior, 1-2; vegetative cells $18-23 \times 23-35\mu$; oogonia $37-41 \times 56-63\mu$; oospores $35-39 \times 54-61\mu$; dwarf male stipes $15-17 \times 24-25\mu$; antheridia $9-10 \times 7-8\mu$.

Brazil.

40. **B. rhadinospora** Wittrock.

(Pl. VIII, fig. 66.)

1874, p. 53; Hirn 1900, p. 372, Pl. LXIII and LXIV, fig. 393; Heering 1914, p. 242, fig. 383; Tiffany 1928, p. 148, Pl. XXI, fig. 66.

Diocious, nannandrous, gynandrosporous; oogonia fusiformly ellipsoid, patent or rarely erect, below androsporangia or terminal setae or rarely vegetative cells; outer wall of oospore longitudinally ribbed, ribs smooth; androsporangia epigynous or rarely scattered, 1-?; dwarf males on or near oogonia; antheridia exterior, 1-4; vegetative cells $15-22 \times 20-44\mu$; oogonia $29-37 \times 61-80\mu$; oospores $27-35 \times 59-78\mu$; androsporangia $13-17 \times 16-23\mu$; dwarf male stipes $13-16 \times 23-25\mu$; antheridia $7-9 \times 6-8\mu$.

Sweden.

40a. Forma **antiqua** (Nordstedt) Hirn.

(Pl. VIII, fig. 67.)

1900, p. 372, Pl. LXIV, fig. 394; Nordstedt 1883, p. 154; *B. varians* Wittr. var. *antiqua* Nordstedt in Borge 1896, p. 4; Tiffany 1928, p. 148, Pl. XXI, fig. 67.

Ribs of oospore crenulate; vegetative cells $15-21 \times 20-42\mu$; oogonia $32-37 \times 58-78\mu$; oospores $30-35 \times 56-76\mu$; androsporangia $15-18 \times 23-26\mu$; dwarf male stipes $14-17 \times 23-26\mu$; antheridia $9-11 \times 6-10\mu$.

Australia.

40b. Var. *litoralis* Hirn.

(Pl. VIII, fig. 68.)

1900, p. 373, Pl. LXIV, fig. 395; Tiffany 1928, p. 149, Pl. XXI, fig. 68.

Larger; vegetative cells $22-28 \times 28-56\mu$; oogonia $38-45 \times 75-87\mu$; oospores $36-43 \times 73-85\mu$; androsporangia $17-20 \times 18-20\mu$; dwarf male stipes $15-17 \times 28-30\mu$; antheridia $9-13 \times 6-9\mu$.

Sweden.

This variety is the only *Bulbochaete* known to inhabit brackish waters.

41. *B. minor* Al. Braun.

(Pl. VII, fig. 54.)

In Kuetzing 1849, p. 422; Wittrock 1874, p. 54; Hirn 1900, p. 369, Pl. LXIII, fig. 390; Collins 1909, p. 275; Heering, p. 242; Tiffany 1926, Pl. X, fig. 114; 1928, p. 149, Pl. XX, fig. 54.

Dioecious, nannandrous, gynandrosporous; oogonia ovoid, erect or rarely patent, below terminal setae, androsporangial cells, or more rarely vegetative cells; outer oospore wall longitudinally ribbed; androsporangia epigynous, subepigynous, or scattered, 1-?; dwarf males on or near oogonia; antheridium exterior, 1-4; vegetative cells $18-25 \times 27-50\mu$; oogonia $32-42 \times 59-69\mu$; oospores $30-40 \times 57-67\mu$; androsporangia $15-16 \times 16-21\mu$; dwarf male stipes $12-15 \times 22-24\mu$; antheridia $6-10 \times 6-7\mu$.

United States: Iowa, Indiana, Illinois, New Jersey; Germany, Sweden, Austria, England.

41a. Var. *germanica* Hirn.

(Pl. VII, fig. 55.)

1900, p. 370, Pl. LXIII, fig. 391; (?) *B. minor* Al. Br. in De Bary 1854, p. 72; *B. minor* Al. Br. in Pringsheim 1858, p. 74; Heering 1914, p. 242, fig. 382; Tiffany 1928, p. 149, Pl. XX, fig. 55.

Vegetative cells $15-22 \times 23-56\mu$; oogonia $30-35 \times 56-65\mu$; oospores $28-33 \times 54-63\mu$; dwarf male stipes $10-12 \times 21-24\mu$; antheridia $6-9 \times 6-7\mu$.

Germany.

This variety on the basis of dimensions and habit seems scarcely tenable. I have not been able to examine the original material, however, and have followed Hirn's disposition of it.

42. *B. bullardi* Transeau and Tiffany.

(Pl. VII, figs. 52 and 53.)

1919, p. 241, Pl. XIV, figs. a-e; Tiffany 1928, p. 149, Pl. XX, figs. 52 and 53.

Dioecious, nannandrous, idioandrosporous; oogonia ovoid-ellipsoid or ellipsoid, erect, terminated by long setae, pore evident and supra-median; oospores filling or nearly filling oogonia, ribs of outer wall of oospore dentate, the teeth connected by transverse lines, longitudinal ribs uniting irregularly; androsporangia 1-6; dwarf males slightly curved, on oogonia or suffultory cells or vegetative cells; antheridia exterior, 1-3; vegetative cells $20-32 \times 60-165\mu$; oogonia $56-66 \times 70-96\mu$;

oospores $54-64 \times 67-94\mu$; androsporangia $18-21 \times 15-33\mu$; dwarf male stipes $18-21 \times 30-33\mu$; antheridia $9-14 \times 6-10\mu$.

United States: Massachusetts.

The combination of dimensions and oospore markings rather readily separates this species from *B. imperialis* and *B. insignis*, its nearest relatives.

43. *B. denticulata* Wittrock.

(Pl. VI, fig. 51.)

1874, p. 54; Hirn 1900, p. 362, Pl. LXI, fig. 379; Heering 1914, p. 240, fig. 377; Tiffany 1928, p. 150, Pl. XIX, fig. 51.

Dioecious, nannandrous; oogonia broadly ellipsoid, patent, below terminal setae or rarely vegetative cells; outer oospore wall longitudinally ribbed, ribs often anastomosing; dwarf males on or near oogonia; antheridia exterior, 1-2; vegetative cells $27-30 \times 34-45\mu$; oogonia $48-54 \times 69-76\mu$; oospores $46-52 \times 67-74\mu$; dwarf male stipes $18-21 \times 30-31\mu$; antheridia $13-14 \times 8-10\mu$.

Norway.

44. *B. affinis* Hirn.

(Pl. VII, fig. 61.)

1900, p. 371, Pl. LXIII, fig. 392; *B. minor* Al. Br. in Borge 1896, p. 4; Tiffany 1928, p. 150, Pl. XX, fig. 61.

Dioecious, nannandrous, gynandrosporous; oogonia suboblong-ellipsoid to ellipsoid, patent or rarely erect, below terminal setae or androsporangia; outer oospore wall longitudinally ribbed, androsporangia epigynous, 1-?; dwarf males near or on oogonia, antheridia exterior, 1-3; vegetative cells $22-27 \times 27-47\mu$; oogonia $40-46 \times 67-78\mu$; oospores $38-44 \times 65-76\mu$; androsporangia $18-19 \times 23-30\mu$; dwarf male stipes $17-19 \times 22-28\mu$; antheridia $9-12 \times 9-13\mu$.

Australia.

45. *B. anomala* Pringsheim.

(Pl. VI, fig. 45.)

1858, p. 73, Pl. 6, fig. 6; Wittrock 1874, p. 56; Hirn 1900, p. 366, Pl. LXII, fig. 385; Heering 1914, p. 240, fig. 380; Tiffany 1928, p. 150, Pl. XIX, fig. 45.

Dioecious, nannandrous; oogonia ellipsoid, erect, below vegetative cells; dwarf males near oogonia; antheridia exterior, 1-2; vegetative cells $26-27 \times 52-54\mu$; oogonia $49-51 \times 75-80\mu$; oospores $47-48 \times 73-78\mu$; dwarf male stipes $18 \times 37\mu$; antheridia $13 \times 13\mu$.

Germany.

46. *B. insignis* Pringsheim.

(Pl. VIII, fig. 62.)

1858, p. 73, Pl. 6, fig. 7; (?) *B. pringsheimiana* Archer 1866, p. 121; *B. speciosa* Wittr. 1870, p. 143; Wittrock 1874, p. 55; (?) *B. pachyderma* Reinsch 1875, p. 82; (?) *B. anomala* Pringsheim in Nordstedt 1888, p. 8; Hirn 1900, p. 364, Pl. LXII, fig. 383; Collins 1909, p. 274; Heering 1914, p. 240, fig. 379; Tiffany 1928, p. 150, Pl. XXI, fig. 62; P. B. A. Nos. 1332 and 1430.

Dioecious, nannandrous, gynandrosporous; oogonia ellipsoid, erect or patent, below androsporangia or terminal setae or vegetative cells; oospores ellipsoid with broadly denticulate longitudinal ribs on the

outer oospore wall; androsporangia epigynous or scattered, 1-?; dwarf males near or on oogonia; antheridia exterior, 1-3; vegetative cells $19-25 \times 48-88\mu$; oogonia $46-56 \times 70-90\mu$; oospores $44-54 \times 68-88\mu$; androsporangia $16-20 \times 9-25\mu$; dwarf male stipes $16-19 \times 29-33\mu$; antheridia $10-13 \times 7-10\mu$.

United States: Massachusetts, Michigan, Illinois, New York, New Jersey, Alaska; Europe, Australia.

46a. Var. **reticulata** (Nordstedt) Hirn.

(Pl. VIII, fig. 63.)

1900, p. 365, Pl. LXII, fig. 384; *B. reticulata* Nördst. 1877, p. 32; Collins 1909, p. 275; Tiffany 1928, p. 151, Pl. XXI, fig. 63.

Outer wall of oospore reticulate-dentate, with doubly dentate, occasionally anastomosing, longitudinal ridges, the teeth united to each other by transverse ridges; vegetative cells $20-25 \times 40-87\mu$; oogonia $44-52 \times 66-85\mu$; oospores $42-50 \times 64-83\mu$; androsporangia $16-19 \times 13-23\mu$; dwarf male stipes $17-20 \times 30-33\mu$; antheridia $11-13 \times 6-9\mu$.

United States: Massachusetts, Illinois; Sweden, Denmark.

47. **B. imperialis** Wittrock.

(Pl. VIII, fig. 64.)

1874, p. 54; Hirn 1900, p. 367, Pl. LXII, fig. 386; Tiffany 1928, p. 151, Pl. XXI, fig. 64.

Dioecious, nannandrous, gynandrosporous; oogonia broadly ellipsoid, erect, terminated by setae; outer oospore wall longitudinally ribbed, the ribs irregularly anastomosing or connected by transverse ribs; androsporangia scattered, 1-?; dwarf males on or near oogonia; antheridia exterior, 1-2 celled; vegetative cells $26-31 \times 32-106\mu$; oogonia $81-83 \times 104-108\mu$; oospores $79-81 \times 102-106\mu$; androsporangia $18-20 \times 25-26\mu$; dwarf male stipes $19-21 \times 30-35\mu$; antheridia $13-16 \times 7-9\mu$.

Brazil.

47a. Var. **regalis** Wittrock.

(Pl. VIII, fig. 65.)

1874, p. 55; Hirn 1900, p. 367, Pl. LXII, fig. 387; Tiffany 1928, p. 151, Pl. XXI, fig. 65.

Smaller than the type; vegetative cells $24-26 \times 72-117\mu$; oogonia $68-70 \times 88-90\mu$; dwarf male stipe $20 \times 39\mu$; antheridia $15-16 \times 6-7\mu$.

Brazil.

48. **B. spirogranulata** West and West.

(Pl. VII, figs. 57 and 58.)

1902, p. 126, Pl. 17, figs. 8 and 9; Hirn 1906, p. 27, Pl. IV, fig. 29.

Oogonia oblong-ellipsoid, subpatent; outer wall of oospore thick with longitudinal, crenulate ribs; androsporangia scattered; suffultory cells without division; setae long; vegetative cells, spirally granulate, $9-12 \times 31-57\mu$; oogonia $20-24 \times 48-49\mu$; androsporangia $7-8 \times 9-10\mu$.

Heneratgodha, Ceylon.

The vegetative cells of this species are minutely granulate, with the markings spirally arranged. This character is rather unique among the ellipsoid members of the genus.

49. **B. brevifulta** Wittrock.

(Pl. VII, fig. 56.)

1874, p. 57; Hirn 1900, p. 373, Pl. LXI, fig. 382; Tiffany 1928, p. 152, Pl. XX, fig. 56.

Oogonia broadly ellipsoid, erect, terminated by setae; oospores with outer wall longitudinally and irregularly ribbed; vegetative cells $19-24 \times 38-72\mu$; oogonia $34-42 \times 50-56\mu$; oospores $32-40 \times 48-54\mu$.

Brazil.

This description is quite incomplete and the position of the species must be considered temporary until more data can be secured.

50. **B. sanguinea** Hansgirg.

1905, p. 437; Hirn 1906, p. 26; Heering 1914, p. 242; Tiffany 1928, p. 152.

Few celled and little branched; cell walls $2-3\mu$ thick; vegetative cells $26-35 \times 24-50\mu$, with contents blood-red in color; reproductive cells unknown.

Austria.

Hansgirg gives no figures for this interesting form, but compares it in habit with *B. minor*. A somewhat similar species, *B. purpurea* Corda (Hansgirg *loc. cit.*) with purple-red vegetative cells alternating with colorless empty cells, was described from Bohemia. Cleve (1895) also observed a red *Bulbochaete*, collected in October, 1894. Until more is known of the reproductive structures of these forms, their specific position must be merely a provisional record.

51. **B. horrida** Nordstedt.

(Pl. VI, fig. 42.)

In Hirn 1900, p. 355, Pl. LIX, fig. 371; Tiffany 1928, p. 152, Pl. XIX, fig. 42.

Dioecious, nannandrous; vegetative cells medianly plicate; oogonia ellipsoid; antheridia of dwarf males 1-?; vegetative cells $16-25 \times 12-25\mu$; basal cells $17-20 \times 10-20\mu$.

South America.

Little is known of the reproduction of this interesting species, and figures are available only for the vegetative cells. The median plications of the latter, however, readily separate it from any other species of the genus known at the present time. Hirn's record of Lagerheim's collections in the Guianas in which dwarf males and ellipsoid oogonia were seen is our only information regarding the fructification of the species.

Genus II. *Oedocladium* Stahl.

1891, p. 339; Hirn 1900, p. 374; Collins 1918, p. 71; Lewis 1930.

Terrestrial or aquatic, branching, rhizoidal; if terrestrial, partly subterranean and subhyaline, partly above ground and green; vegetative cell cylindrical or nearly so, growth chiefly by division of terminal cell of a filament or branch; asexual reproduction by "akinetes" and zoospore; oogonium arising by a simple division of a vegetative cell; sexual reproduction both monoecious and dioecious, nannandrous.

KEY TO THE SPECIES OF OEDOCLADIUM.

1. Monoecious, terrestrial.....	2
1. Dioecious-nannandrous, aquatic.....	4. <i>O. hazenii</i>
2. Oospore smooth.....	3
2. Oospore angulate.....	4
3. Vegetative cell $7 \times 20\mu$	1. <i>O. protonema</i>
3. Vegetative cell $12-18 \times 56-78\mu$	3. <i>O. media</i>
4. Oogonium $52 \times 52\mu$	3. <i>O. media</i>
4. Oogonium $90 \times 95\mu$	2. <i>O. albemarlensis</i>

1. *O. protonema* Stahl.

(Pl. LXIII, figs. 619-623.)

1891, p. 347, Pl. XVI, figs. 1-6, Pl. XVII, figs. 1-12; Hirn 1900, p. 374, Pl. LXIV, fig. 396.

Monoecious; oogonium single, subglobose, rarely terminal, pore median; oospore globose, filling oogonium, spore wall smooth; antheridium scattered, 1-6; vegetative cell $7 \times 20\mu$, rhizoidal $3 \times$ (up to) 300μ ; oogonium $50-76 \times 58-76\mu$; oospore $45-60 \times 45-60\mu$.

Germany (Strassburg).

2. *O. albemarlensis* Lewis.

(Pl. LXIII, figs. 613-617.)

In Collins (as *O. albemarlense* Lewis) 1918, p. 71, Pl. III, figs. 22a and 22b.

Monoecious; oogonium single, mostly terminal, subglobose with conical apex (if terminal), pore inferior; oospore globose, not filling oogonium, spore wall of three layers: outer layer smooth, middle layer angulate (in optical section undulate), inner layer smooth; antheridium to 9; vegetative cell subcylindrical; terminal cell with conical apex; vegetative cell $25-40 \times 50-150\mu$, rhizoidal $4-10 \times$ (up to) 350μ ; oogonium $90 \times 95\mu$; oospore $69 \times 69\mu$; antheridium $20-25 \times 8-19\mu$; zoospore $32-50\mu$.

United States: Virginia (on sandy loam, bank of the Rivanna, near Charlottesville, Albemarle County) growing with *Vaucheria* and *Riccia*. Collected by Professor I. F. Lewis.

3. **O. media** Lewis nov. sp. mss.

(Pl. LXI, figs. 601-603.)

Monoecious; oogonium single, mostly terminal, globose to subglobose, with conical apex (if terminal), pore median; oospore globose, not quite filling oogonium, spore wall with middle layer angulate or smooth; antheridium 1-2; terminal cell with conical apex; vegetative cell $12-18 \times 56-78\mu$, rhizoidal $10 \times$ (up to) 270μ ; oogonium $52 \times 52\mu$; oospore $44 \times 44\mu$; antheridium $15 \times 9-15\mu$.

United States: Massachusetts (Woods Hole). Collected by Professor I. F. Lewis.

4. **O. hazenii** Lewis nov. sp. mss.

(Pl. LXII, figs. 604-612.)

Dioecious, nannandrous, gynandrosporous; oogonium globose to subglobose, terminal or intercalary, pore inferior; oospore globose, not or nearly filling oogonium, spore wall with middle layer angulate; androsporangium 1-2, epigynous, subepigynous, or scattered; dwarf male curved or erect, on suffultory cell or more rarely on oogonium; antheridium exterior, single; vegetative cell $15-26 \times 47-132\mu$, rhizoidal $4 \times$ (up to) 425μ ; oogonium $50-75 \times 50-75\mu$; oospore $45-55 \times 45-55\mu$; androsporangium $16 \times 7-12\mu$; dwarf male stipe $18-22 \times 7-10\mu$; antheridium $6-8 \times 6-7\mu$.

United States: New Jersey. Collected by Professor T. E. Hazen.

O. hazenii is the only dioecious, nannandrous species of the genus as well as the only aquatic member, so far recorded.

Data for descriptions and plates of *Oedocladium media* and *O. hazenii* were supplied by Professor I. F. Lewis of the University of Virginia. Lewis' *O. hazenii* makes it necessary to emend Stahl's original description of the genus to include this nannandrous species growing in an aquatic habitat.

Genus III. *Oedogonium* Link.

1820, p. 5; Pringsheim 1858, p. 68; Wittrock 1874, p. 6; Hirn 1900, p. 72; Collins 1909, p. 223; *Prolifera* Vaucher 1803, p. 14; *Vesiculifera* Hassall 1845, p. 195; *Cymatonema* Kuetzing 1849, p. 375; *Androgynia* Wood 1872, p. 196; *Pringsheimia* Wood 1872, p. 195; *Conferva* (early authors).

Filaments single, unbranched; vegetative cells cylindrical, or sometimes capitellate, nodulose or undulate; basal cell with holdfast; terminal cell obtuse, apiculate, or hyaline; all vegetative cells, except the basal one, capable of division; oogonia and antheridia produced by direct division of vegetative cells.

KEY TO THE SPECIES, VARIETIES, AND FORMS OF
OEDOGONIUM

1. Vegetative cell undulate or nodulose.....	2
1. Vegetative cell punctate or granulate.....	6
1. Vegetative cell distinctly capitellate.....	7
1. Vegetative cell cylindrical.....	35
1. Vegetative cell subhexagonal or subellipsoid.....	236. <i>Oe. reinschii</i> *
2. Without dwarf male.....	3
2. With dwarf male.....	4
3. Diameter of oogonium 18-23 μ	105. <i>Oe. sphaerandrium</i>
3. Diameter of oogonium 48-57 μ	96. <i>Oe. nodulosum</i>
3. Diameter of oogonium 64-74 μ	96a. <i>Oe. nodulosum</i> var. <i>commune</i>
4. Diameter of oogonium 44-56 μ	5
4. Diameter of oogonium 58-68 μ	119a. <i>Oe. undulatum</i> var. <i>americanum</i>
5. Dwarf male 36-46 μ in length.....	119b. <i>Oe. undulatum</i> f. <i>senegalense</i>
5. Dwarf male 48-70 μ in length.....	119. <i>Oe. undulatum</i>
6. Diameter of vegetative cell 6-8 μ	155. <i>Oe. elegans</i>
6. Diameter of vegetative cell 9-13 μ	84. <i>Oe. minus</i>
6. Diameter of vegetative cell 16-22 μ	85. <i>Oe. punctatostriatum</i>
7. Without dwarf male.....	8
7. With dwarf male.....	17
7. Reproductive structures imperfectly known.....	34
8. Dioecious.....	9
8. Monoecious.....	11
9. Division of oogonium basal.....	78. <i>Oe. infimum</i>
9. Division of oogonium inferior.....	77. <i>Oe. inversum</i>
9. Division of oogonium median.....	10
9. Division of oogonium suprmedian or superior.....	89. <i>Oe. mitratum</i>
10. Oogonium 26-29 μ in diameter.....	79a. <i>Oe. howardii</i> var. <i>minus</i>
10. Oogonium 29-32 μ in diameter.....	79. <i>Oe. howardii</i>
10. Oogonium 32-36 μ in diameter.....	81. <i>Oe. latiusculum</i>
11. Oogonium opening by a pore.....	20a. <i>Oe. hirnii</i> var. <i>africanum</i>
11. Oogonium opening by a lid.....	12
12. Diameter of oogonium 15-25 μ	13
12. Diameter of oogonium 34-46 μ	16
13. Division of oogonium median.....	14
13. Division of oogonium suprmedian.....	15
14. Oogonium subdepressed-globose.....	86. <i>Oe. capitellatum</i>
14. Oogonium angular-globose.....	87. <i>Oe. quadratum</i>
15. Diameter of oogonium 15-20 μ	210. <i>Oe. virceburgense</i> *
15. Diameter of oogonium 18-23 μ	105. <i>Oe. sphaerandrium</i>
15. Diameter of oogonium 26-29 μ	211. <i>Oe. spurium</i> *

*Species so marked have incomplete descriptions and are included in what is perhaps their probable position in the genus, using such characters as are available.

16.	Vegetative cell punctate.....	84.	<i>Oe. minus</i>	
16.	Vegetative cell not punctate.....	88.	<i>Oe. bohemicum</i>	
17.	Poriferous.....	120.	<i>Oe. nebraskense</i>	
17.	Operculate.....			18
	18. Division of oogonium median.....			19
	18. Division of oogonium supramedian.....			24
	18. Division of oogonium inframedian.....			26
	18. Division of oogonium superior.....	156.	<i>Oe. rigidum</i>	
	18. Division of oogonium supreme.....			29
19.	Division of oogonium narrow.....			20
19.	Division of oogonium wide.....			21
	20. Oogonium $28-35 \times 23-38\mu$	171a.	<i>Oe. decipiens</i> f. <i>dissimile</i>	
	20. Oogonium $46-54 \times 44-54\mu$	157.	<i>Oe. bengalense</i>	
	20. Oogonium $57-64 \times 48-53\mu$	158.	<i>Oe. indicum</i>	
21.	Diameter of oospore $22-26\mu$			22
21.	Diameter of oospore $30-32\mu$	160b.	<i>Oe. areschougii</i> f. <i>robustum</i>	
21.	Diameter of oospore $42-48\mu$			23
21.	Diameter of oospore $48-53\mu$	159.	<i>Oe. brasiliense</i>	
	22. Gynandrosporous.....	160.	<i>Oe. areschougii</i>	
	22. Idioandrosporous.....	160a.	<i>Oe. areschougii</i> var. <i>americanum</i>	
23.	Vegetative cell $14-19\mu$ in diameter.....	157.	<i>Oe. bengalense</i>	
23.	Vegetative cell $19-26\mu$ in diameter.....	161.	<i>Oe. confertum</i>	
	24. Diameter of oogonium $14-26\mu$	163.	<i>Oe. clavatum</i>	
	24. Diameter of oogonium $27-42\mu$			25
	24. Diameter of oogonium $50-55\mu$	173.	<i>Oe. costatum</i>	
	24. Diameter of oogonium $58-65\mu$	174.	<i>Oe. boreale</i>	
25.	Oogonium $19-25\mu$ in length.....	164a.	<i>Oe. oelandicum</i> f. <i>minus</i>	
25.	Oogonium $25-32\mu$ in length.....	164.	<i>Oe. oelandicum</i>	
25.	Oogonium $40-45\mu$ in length.....	165.	<i>Oe. megaporum</i>	
	26. Suffultory cell enlarged.....	167b.	<i>Oe. platygynum</i> var. <i>continuum</i>	
	26. Suffultory cell not enlarged.....			27
27.	Diameter of vegetative cell $6-10\mu$			28
27.	Diameter of vegetative cell $12-15\mu$	166.	<i>Oe. bahusiense</i>	
	28. Gynandrosporous only.....	167c.	<i>Oe. platygynum</i> var. <i>novaezelandiae</i>	
	28. Idioandrosporous only.....	167a.	<i>Oe. platygynum</i> f. <i>obtusum</i>	
	28. Gynandrosporous and idioandrosporous.....	167.	<i>Oe. platygynum</i>	
29.	Oospore wall smooth.....			30
29.	Oospore wall longitudinally ribbed.....	194.	<i>Oe. michiganense</i>	
	30. Idioandrosporous.....			31
	30. Gynandrosporous.....			32
31.	Oogonium $48-60 \times 62-74\mu$	190.	<i>Oe. praticolum</i>	
31.	Oogonium $66-78 \times 72-90\mu$	191.	<i>Oe. supremum</i>	
	32. Diameter of oogonium $36-42\mu$	192.	<i>Oe. wabashense</i>	
	32. Diameter of oogonium $42-55\mu$			33
	32. Diameter of oogonium $55-58\mu$	189a.	<i>Oe. obtruncatum</i> var. <i>completum</i>	
33.	Length of oogonium $56-68\mu$	189.	<i>Oe. obtruncatum</i>	
33.	Length of oogonium $68-75\mu$	189b.	<i>Oe. obtruncatum</i> var. <i>ellipsoideum</i>	
	34. Vegetative cell $2-3\mu$ in diameter.....	203.	<i>Oe. fusus</i> *	
	34. Vegetative cell $4-6\mu$ in diameter.....	210.	<i>Oe. virceburgense</i> *	
	34. Vegetative cell $6-13\mu$ in diameter.....	211.	<i>Oe. spurium</i> *	
35.	Diameter of vegetative cell not more than 2μ	197.	<i>Oe. angustissimum</i> *	
35.	Diameter of vegetative cell $2-60\mu$			36
35.	Diameter of vegetative cell ($56-$) $60-93\mu$			268
	36. Without dwarf males.....			37
	36. With dwarf males.....			186
37.	Oogonium opening by a pore.....			38
37.	Oogonium opening by a lid.....			139
	38. Pore median.....			39
	38. Pore supramedian (rarely varying to superior).....			51
	38. Pore superior†.....			63

*See footnote, page 53.

†Some variations of *Oe. varians* may be sought here.

39.	Wall of oospore smooth.....	40
39.	Wall of oospore scrobiculate.....	49
39.	Wall of oospore echinate.....	50
	40. Monoecious.....	41
	40. Dioecious.....	44
41.	Diameter of oogonium 18-28 μ	42
41.	Diameter of oogonium 32-38 μ	1. <i>Oe. laeve</i>
	42. Filament irregularly curved.....	2. <i>Oe. curvum</i>
	42. Filament straight.....	43
43.	Oogonium 23-28 $\times$ 26-31 μ	3. <i>Oe. cryptoporum</i>
43.	Oogonium 18-25 $\times$ 18-26 μ	3a. <i>Oe. cryptoporum</i> var. <i>vulgare</i>
	44. Diameter of oogonium 22-27 μ	45
	44. Diameter of oogonium 27-32 μ	47
	44. Diameter of oogonium 30-39 μ	48
45.	Diameter of vegetative cell 5-10 μ	46
45.	Diameter of vegetative cell 8-13 μ	4b. <i>Oe. rufescens</i> var. <i>lundellii</i>
	46. Diameter of vegetative cell 5-9 μ	4a. <i>Oe. rufescens</i> var. <i>exiguum</i>
	46. Diameter of vegetative cell 8-10 μ	4. <i>Oe. rufescens</i>
47.	Diameter of vegetative cell 9-11 μ	218a. <i>Oe. inerme</i> var. <i>mentiens</i> *
47.	Diameter of vegetative cell 11-14 μ	5. <i>Oe. calcareum</i>
	48. Oospore 28-35 $\times$ 28-35 μ	6. <i>Oe. sociale</i>
	48. Oospore 33-38 $\times$ 28-32 μ	218. <i>Oe. inerme</i> *
49.	Monoecious.....	7. <i>Oe. cymalosporum</i>
49.	Dioecious (very rarely monoecious).....	8. <i>Oe. magnusii</i>
	50. Diameter of vegetative cell 9-14 μ	9. <i>Oe. suecicum</i>
	50. Diameter of vegetative cell 14-16 μ ,.....	9a. <i>Oe. suecicum</i> f. <i>australe</i>
51.	Oospore scrobiculate, 22-27 μ in diameter.....	217. <i>Oe. moniliforme</i> *
51.	Oospore scrobiculate, 43-48 μ in diameter,.....	66a. <i>Oe. argenteum</i> f. <i>michiganense</i>
51.	Oospore smooth.....	52
	52. Monoecious.....	53
	52. Dioecious.....	58
	52. Both monoecious and dioecious.....	10. <i>Oe. varians</i>
53.	Pore a little above median.....	54
53.	Pore nearly superior.....	56
53.	Pore variable between these two positions.....	57
	54. Diameter of oogonium 34-45 μ	55
	54. Diameter of oogonium 46-55 μ	11. <i>Oe. urbicum</i>
55.	Oospore depressed-globose, 30-34 $\times$ 28-32 μ	12. <i>Oe. obsoletum</i>
55.	Oospore regularly globose, 30-39 $\times$ 30-39 μ	13. <i>Oe. plusiosporum</i>
	56. Oospore 31-41 $\times$ 30-41 μ	10. <i>Oe. varians</i>
	56. Oospore 40-48 $\times$ 40-48 μ	14. <i>Oe. tyrolicum</i>
57.	Terminal cell narrowed and often setiferous.....	225. <i>Oe. inflatum</i> *
57.	Terminal cell neither narrowed nor setiferous.....	11. <i>Oe. urbicum</i>
	58. Pore a little above median.....	59
	58. Pore nearly superior.....	60
	58. Pore variable between these two positions.....	62
59.	Vegetative cell 1-3 diameters long.....	15b. <i>Oe. cardiacum</i> f. <i>pulchellum</i>
59.	Vegetative cell 3-7 diameters long.....	15. <i>Oe. cardiacum</i>
	60. Diameter of vegetative cell 8-12 μ	16. <i>Oe. franklinianum</i>
	60. Diameter of vegetative cell 12-16 μ	10. <i>Oe. varians</i>
	60. Diameter of vegetative cell 14-30 μ	61
61.	Oospore depressed-globose, 31-42 μ in diameter,.....	15d. <i>Oe. cardiacum</i> var. <i>minus</i>
61.	Oospore subglobose, 42-49 μ in diameter.....	17. <i>Oe. glabrum</i>
61.	Oospore globose, 43-58 μ in diameter.....	15a. <i>Oe. cardiacum</i> f. <i>interjectum</i>
61.	Oospore ellipsoid-globose, 40-52 μ in diameter,.....	15c. <i>Oe. cardiacum</i> var. <i>carbonicum</i>
	62. Oospore not filling oogonium.....	225. <i>Oe. inflatum</i> *
	62. Oospore completely filling oogonium.....	18. <i>Oe. lautumnarium</i>

*See footnote, page 53.

63.	Wall of oospore smooth.....	64
63.	Wall of oospore longitudinally ribbed.....	116
63.	Wall of oospore areolate.....	129
63.	Wall of oospore reticulate.....	131
63.	Wall of oospore pitted.....	132
63.	Wall of oospore scrobiculate.....	133
64.	Monoeocious.....	65
64.	Dioecious.....	83
64.	Reproductive structures imperfectly known.....	113
65.	Diameter of vegetative cell 8-34 μ	66
65.	Diameter of vegetative cell 33-54 μ	80
66.	Diameter of oogonium 32-36 (-40) μ	67
66.	Diameter of oogonium 36-63 μ	70
66.	Diameter of oogonium 63-68 μ ... 29a. <i>Oe. upsaliense</i> var. <i>fennicum</i>	
67.	Vegetative cell 8-14 μ in diameter.....	68
67.	Vegetative cell 15-18 μ in diameter..... 19. <i>Oe. intermedium</i>	
68.	Length of oogonium 32-46 μ	69
68.	Length of oogonium about 53 μ 226. <i>Oe. warmingianum</i> *.....	
69.	Oospore 28-31 μ in diameter..... 20. <i>Oe. hirnii</i>	
69.	Oospore 30-37 (-40) μ in diameter..... 21. <i>Oe. globosum</i>	
70.	Plant few celled when mature.....	71
70.	Plant many celled when mature.....	72
71.	Antheridium single, alternating with vegetative cell. 22. <i>Oe. zigzag</i>	
71.	Antheridium 1-4, in series..... 23. <i>Oe. curtum</i>	
72.	Oospore globose to subglobose.....	73
72.	Oospore obovoid to ellipsoid.....	77
73.	Diameter of vegetative cell 12-19 μ	74
73.	Diameter of vegetative cell 19-34 μ	76
74.	Oospore completely filling oogonium.....	75
74.	Oospore not filling oogonium... 19a. <i>Oe. intermedium</i> var. <i>fennicum</i>	
75.	Antheridium 1-3..... 24. <i>Oe. fragile</i>	
75.	Antheridium 3-6..... 24a. <i>Oe. fragile</i> var. <i>abyssinicum</i>	
76.	Oogonium usually globose..... 24b. <i>Oe. fragile</i> var. <i>robustum</i>	
76.	Oogonium usually ovoid..... 25. <i>Oe. vaucherii</i>	
77.	Division of antheridium horizontal.....	26. <i>Oe. richterianum</i>
77.	Division of antheridium vertical.....	78
78.	Vegetative cell 8-14 μ in diameter..... 27. <i>Oe. pseudoboscii</i>	
78.	Vegetative cell 14-24 μ in diameter.....	79
79.	Oogonium 40-44 $\times$ 70-80 μ	28. <i>Oe. sodiroanum</i>
79.	Oogonium 45-50 $\times$ 66-100 μ	29. <i>Oe. upsaliense</i>
79.	Oogonium 48-55 $\times$ 65-80 μ	30. <i>Oe. oviforme</i>
80.	Diameter of oogonium 53-67 μ	81
80.	Diameter of oogonium 68-95 μ	82
81.	Diameter of vegetative cell 37-48 μ	31. <i>Oe. geniculatum</i>
81.	Diameter of vegetative cell 50-54 μ	32. <i>Oe. suboctangulare</i>
82.	Diameter of vegetative cell 33-37 μ	33. <i>Oe. martinicense</i>
82.	Diameter of vegetative cell 44-52 μ	34. <i>Oe. kurzii</i>
83.	Diameter of vegetative cell 11-13 μ 223. <i>Oe. lageniforme</i> *.....	
83.	Diameter of vegetative cell 14-56 μ	84
84.	Oogonium scarcely exceeding vegetative cell in diameter.....	85
84.	Oogonium noticeably exceeding vegetative cell in diameter.....	86
85.	Oospore globose to cylindric-globose.....	35. <i>Oe. capillare</i>
85.	Oospore cylindric-globose to subcylindrical, 35a. <i>Oe. capillare</i> f. <i>stagnale</i>	
85.	Oospore subrectangular-ellipsoid..... 32. <i>Oe. suboctangulare</i>	
86.	Oospore globose, subglobose, or cylindric-globose.....	87
86.	Oospore ellipsoid, obovoid, subcylindrical, or subellipsoid.....	95
87.	Male filament of same diameter as female.....	88
87.	Male filament smaller than female.....	90
88.	Oogonium 36-42 μ in diameter... 36a. <i>Oe. plagiostomum</i> var. <i>gracilius</i>	
88.	Oogonium 42-49 μ in diameter..... 36. <i>Oe. plagiostomum</i>	
88.	Oogonium 54-63 μ in diameter.....	89

*See footnote, page 53.

89.	Vegetative cell 16-26 × 112-250μ.....	17.	<i>Oe. glabrum</i>	
89.	Vegetative cell 33-42 × 42-130μ.....	38.	<i>Oe. princeps</i>	
	90. Diameter of male vegetative cell 18-33μ.....			91
	90. Diameter of male vegetative cell 34-50μ.....			94
91.	Oospore usually ellipsoid to cylindric-globose.....	37.	<i>Oe. capilliforme</i>	
91.	Oospore usually globose to subglobose.....			92
	92. Division of antheridium horizontal.....			93
	92. Division of antheridium vertical.....	39.	<i>Oe. biforme</i>	
93.	Oogonium 40-53 × 40-65μ.....	37b.	<i>Oe. capilliforme</i> f. <i>lorentzii</i>	
93.	Oogonium 46-57 × 52-64 (-80)μ.....	37a.	<i>Oe. capilliforme</i> f. <i>debaryanum</i>	
93.	Oogonium 36-53 × 43-60μ.....	37c.	<i>Oe. capilliforme</i> var. <i>australe</i>	
	94. Oogonium 46-56μ in diameter.....	37d.	<i>Oe. capilliforme</i> var. <i>diversum</i>	
	94. Oogonium 54-65μ in diameter.....	40.	<i>Oe. anomalum</i>	
	94. Oogonium 70-85μ in diameter.....	42.	<i>Oe. rivulare</i>	
95.	Diameter of oogonium 35-63 (-68)μ.....			96
95.	Diameter of oogonium (60-) 63-90μ.....			106
	96. Male filament larger than female.....	43.	<i>Oe. pachyandrium</i>	
	96. Male filament not larger than female.....			97
97.	Vegetative cell 1-3 diameters long.....			98
97.	Vegetative cell 2-6 diameters long.....			100
97.	Vegetative cell 3-11 diameters long.....			102
	98. Oospore 40-60μ long.....	35a.	<i>Oe. capillare</i> f. <i>stagnale</i>	
	98. Oospore 60-80μ long.....			99
99.	Diameter of oospore 41-51μ.....	209.	<i>Oe. oryzae</i> *	
99.	Diameter of oospore 40-56μ.....	209a.	<i>Oe. oryzae</i> var. <i>seriosporum</i> *	
99.	Diameter of oospore 51-60μ.....	44.	<i>Oe. mexicanum</i>	
	100. Cell diameter of male filament 18-28μ.....			101
	100. Cell diameter of male filament 28-33μ.....	45.	<i>Oe. grande</i>	
	100. Cell diameter of male filament 32-42μ.....			
		45d.	<i>Oe. grande</i> var. <i>majus</i>	
101.	Oogonium usually obovoid.....	45b.	<i>Oe. grande</i> var. <i>aequatoriale</i>	
101.	Oogonium usually subobovoid.....	39.	<i>Oe. biforme</i>	
102.	Division of antheridium horizontal.....			
		15c.	<i>Oe. cardiacum</i> var. <i>carbonicum</i>	
	102. Division of antheridium vertical.....			103
103.	Oogonium usually obovoid.....	46.	<i>Oe. oboviforme</i>	
103.	Oogonium usually cylindric-obovoid.....	39.	<i>Oe. biforme</i>	
103.	Oogonium usually subovoid or ellipsoid-ovoid.....			104
	104. Diameter of oogonium 42-52μ.....	45c.	<i>Oe. grande</i> var. <i>angustum</i>	
	104. Diameter of oogonium 49-68μ.....			105
105.	Diameter of vegetative cell 20-27μ.....	45a.	<i>Oe. grande</i> f. <i>gemelliparum</i>	
105.	Diameter of vegetative cell 28-37μ.....	45.	<i>Oe. grande</i>	
105.	Diameter of vegetative cell 36-46μ.....	45d.	<i>Oe. grande</i> var. <i>majus</i>	
	106. Oospore conspicuously not filling oogonium.....	42.	<i>Oe. rivulare</i>	
	106. Oospore nearly or quite filling oogonium.....			107
107.	Vegetative cell 1-6 diameters long.....			108
107.	Vegetative cell 6-10 diameters long.....	47.	<i>Oe. subrectum</i>	
	108. Cell diameter of male filament 30-37μ.....			109
	108. Cell diameter of male filament 36-50μ.....			110
109.	Oospore obovoid to ellipsoid.....	48.	<i>Oe. landsboroughi</i>	
109.	Oospore ellipsoid to globose-ellipsoid.....	49.	<i>Oe. crassum</i>	
	110. Length of oospore 77-100μ.....			111
	110. Length of oospore 100-125μ.....	49b.	<i>Oe. crassum</i> var. <i>longum</i>	
111.	Diameter of oogonium 64-73μ.....	48a.	<i>Oe. landsboroughi</i> var. <i>norvegicum</i>	
111.	Diameter of oogonium 75-90μ.....			112
	112. Vegetative cell 1-3 diameters long.....			
		49a.	<i>Oe. crassum</i> f. <i>amplum</i>	
	112. Vegetative cell 3-5 diameters long.....			
		49c.	<i>Oe. crassum</i> var. <i>subtumidum</i>	
113.	Diameter of vegetative cell 6-24 (-27)μ.....			114
113.	Diameter of vegetative cell 24-42μ.....			115

*See footnote, page 53.

114.	Oogonium about $35 \times 53\mu$	226.	<i>Oe. warmingianum</i> *
114.	Oogonium $33-36 \times 48-63\mu$	223.	<i>Oe. lageniforme</i> *
114.	Oogonium $37-49 \times 54-76\mu$	225.	<i>Oe. inflatum</i> *
115.	Oogonium $43-55 \times 45-95\mu$	209.	<i>Oe. oryzae</i> *
115.	Oogonium $42-60 \times 43-99\mu$	209a.	<i>Oe. oryzae</i> var. <i>seriosporum</i> *
116.	Outer oospore wall ribbed externally.....	50.	<i>Oe. exocostatum</i>
116.	Middle oospore wall ribbed externally.....	117	
117.	Ribs continuous and entire.....	118	
117.	Ribs granulate, crenate, crenulate, or pitted.....	126	
117.	Ribs dentate.....	57b.	<i>Oe. crenulatocostatum</i> var. <i>aureum</i>
118.	Ribs 15-22 in number.....	119	
118.	Ribs 26-35 in number.....	120	
118.	Ribs 35-45 in number.....	51.	<i>Oe. kjellmanii</i>
119.	Vegetative cell $10-15\mu$ in diameter.....	52.	<i>Oe. paulense</i>
119.	Vegetative cell $24-30\mu$ in diameter.....	220.	<i>Oe. urceolatum</i> *
120.	Monoecious.....	121	
120.	Dioecious.....	123	
121.	Oospore completely filling oogonium.....	122	
121.	Oospore not filling oogonium.....	53b.	<i>Oe. paludosum</i> var. <i>parvisporum</i>
122.	Oogonium $39-48\mu$ in diameter.....	53.	<i>Oe. paludosum</i>
122.	Oogonium $54-63\mu$ in diameter.....	53a.	<i>Oe. paludosum</i> var. <i>americanum</i>
123.	Oospore usually completely filling oogonium.....	54.	<i>Oe. leiopleurum</i>
123.	Oospore usually not filling oogonium.....	124	
124.	Oospore ellipsoid.....	125	
124.	Oospore ellipsoid-globose, rarely subglobose,		
	55a. <i>Oe. boscii</i> f. <i>dispar</i>		
124.	Oospore ellipsoid-obovoid or obovoid,		
	55b. <i>Oe. boscii</i> var. <i>notabile</i>		
125.	Diameter of oogonium $33-38\mu$	55c.	<i>Oe. boscii</i> var. <i>occidentale</i>
125.	Diameter of oogonium $39-51\mu$	55.	<i>Oe. boscii</i>
126.	Ribs 14-20 in number.....	127	
126.	Ribs 30-35 in number.....	56.	<i>Oe. margaritifera</i>
126.	Ribs 35-45 in number.....	51.	<i>Oe. kjellmanii</i>
127.	Ribs crenulate.....	128	
127.	Ribs crenate.....	57c.	<i>Oe. crenulatocostatum</i> var. <i>longiarticulatum</i>
128.	Ribs distinctly crenulate.....	57.	<i>Oe. crenulatocostatum</i>
128.	Ribs scarcely crenulate.....	57a.	<i>Oe. crenulatocostatum</i> f. <i>cylindricum</i>
129.	Dioecious.....	130	
129.	Monoecious.....	58.	<i>Oe. arcyosporum</i>
130.	Diameter of oogonium $30-35\mu$	57b.	<i>Oe. crenulatocostatum</i> var. <i>aureum</i>
130.	Diameter of oogonium $48-60\mu$	59.	<i>Oe. areolatum</i>
130.	Diameter of oogonium $66-72\mu$	214.	<i>Oe. hoecknei</i> *
131.	Oogonium $33-40 \times 38-46\mu$	60.	<i>Oe. dictyosporum</i>
131.	Oogonium about $42 \times 52\mu$	60a.	<i>Oe. dictyosporum</i> f. <i>westii</i>
132.	Oogonium $35-38 \times 53-58\mu$	216.	<i>Oe. capense</i> *
132.	Oogonium $53-69 \times 67-106\mu$	215.	<i>Oe. giganteum</i> *
133.	Monoecious.....	61.	<i>Oe. foveolatum</i>
133.	Dioecious.....	134	
134.	Diameter of vegetative cell $11-25\mu$	135	
134.	Diameter of vegetative cell $25-48\mu$	138	
135.	Suffultory cell enlarged.....	136	
135.	Suffultory cell not enlarged.....	137	
136.	Diameter of oogonium $40-48\mu$	62.	<i>Oe. scrobiculatum</i>
136.	Diameter of oogonium $56-58\mu$	63.	<i>Oe. verrucosum</i>
136.	Diameter of oogonium $64-76\mu$	64.	<i>Oe. tiffanii</i>
137.	Diameter of oogonium $38-45\mu$	65.	<i>Oe. punctatum</i>
137.	Diameter of oogonium $45-53\mu$	66.	<i>Oe. argenteum</i>
137.	Diameter of oogonium $51-64\mu$	67.	<i>Oe. wylicii</i>
138.	Length of oogonium $48-70\mu$	68.	<i>Oe. americanum</i>
138.	Length of oogonium $78-113\mu$	69.	<i>Oe. taphrosporum</i>

*See footnote, page 53.

WITHOUT DWARF MALES; OPERCULATE OOGONIUM, CYLINDRICAL
VEGETATIVE CELLS.

139.	Wall of oospore echinate.....	212.	<i>Oe. velatum</i> *	
139.	Wall of oospore punctate-granulate....	109a.	<i>Oe. crispum</i> f. <i>granulosum</i>	
139.	Wall of oospore longitudinally ribbed.....			140
139.	Wall of oospore smooth.....			145
140.	Division of oogonium superior.....			141
140.	Division of oogonium supreme.....	70.	<i>Oe. pseudacrosporum</i>	
141.	Ribs 15-23 in number.....			142
141.	Ribs 30-50 in number.....			143
	142. Diameter of oogonium 48-52 μ	71a.	<i>Oe. paucocostatum</i> var. <i>gracilis</i>	
	142. Diameter of oogonium 54-60 μ	71.	<i>Oe. paucocostatum</i>	
	142. Diameter of oogonium 59-74 μ	72.	<i>Oe. australianum</i>	
143.	Diocious.....	73.	<i>Oe. tumidulum</i>	
143.	Monoecious.....			144
143.	Reproductive structures imperfectly known.....	219.	<i>Oe. sol</i> *	
	144. Oospore 48-55 $\times$ 50-58 μ	74.	<i>Oe. nobile</i>	
	144. Oospore 58-59 $\times$ 72-80 μ	74a.	<i>Oe. nobile</i> var. <i>minus</i>	
	144. Oospore 65-75 $\times$ 88-104 μ	75.	<i>Oe. insigne</i>	
145.	Oogonium without projections.....			146
145.	Oogonium with projections.....			184
	146. Oospore with a median constriction.....			147
	146. Oospore without a median constriction.....			148
	Oogonium medianly plicate.....	76.	<i>Oe. excisum</i>	
147.	Oogonium not medianly plicate.....	200.	<i>Oe. pusillum</i> *	
	148. Division of oogonium median.....			149
	148. Division of oogonium supramedian.....			155
	148. Division of oogonium superior.....			157
	148. Division of oogonium inferior.....	77.	<i>Oe. inversum</i>	
	148. Division of oogonium basal.....	78.	<i>Oe. infimum</i>	
149.	Diameter of oogonium 13-20 μ			150
149.	Diameter of oogonium 20-30 μ			151
149.	Diameter of oogonium 30-40 μ			152
	150. Oogonium 13-15 $\times$ 18-23 μ	198.	<i>Oe. inconspicuum</i> *	
	150. Oogonium 14-15 $\times$ 31-35 μ	203.	<i>Oe. fusus</i> *	
	150. Oogonium 15-19 $\times$ 18-23 μ	196.	<i>Oe. tapeinosporum</i> *	
151.	Division of oogonium narrow.....	101.	<i>Oe. gunnii</i>	
151.	Division of oogonium wide.....	224.	<i>Oe. poecilosporum</i> *	
	152. Dioecious.....			153
	152. Monoecious.....			154
153.	Diameter of oogonium 29-33 μ	79.	<i>Oe. howardii</i>	
153.	Diameter of oogonium 33-40 μ	80.	<i>Oe. pratense</i>	
	154. Oogonium subglobose.....	82.	<i>Oe. acmandrium</i>	
	154. Oogonium pyriform-globose.....	83.	<i>Oe. psaeognatosporum</i>	
155.	Dioecious.....	89.	<i>Oe. mitratum</i>	
155.	Monoecious.....	90.	<i>Oe. petri</i>	
155.	Antheridium not known.....			156
	156. Oogonium 15-20 $\times$ 15-23 μ	210.	<i>Oe. virceburgense</i> *	
	156. Oogonium 27-28 $\times$ 26-28 μ	222.	<i>Oe. consociatum</i> *	
157.	Dioecious.....			158
157.	Monoecious.....			165
157.	Both dioecious and monoecious.....	91.	<i>Oe. trioicum</i>	
157.	Reproductive structures imperfectly known.....			183
	158. Diameter of oogonium 13-16 μ	91.	<i>Oe. trioicum</i>	
	158. Diameter of oogonium 16-43 μ			159
	158. Diameter of oogonium 43-50 μ	92.	<i>Oe. welwitschii</i>	
	158. Diameter of oogonium 52-60 μ	93.	<i>Oe. iowense</i>	
	158. Diameter of oogonium about 82 μ	94.	<i>Oe. dioicum</i>	
159.	Oospore globose to subglobose.....			160
159.	Oospore ellipsoid, ovoid, or oblong.....			162

*See footnote, page 53.

160. Vegetative cell 1-3 diameters long, 95a. *Oe. pringsheimii* var. *abbreviatum*
160. Vegetative cell 2-5 diameters long, 161
161. Female vegetative cell 10-15 μ in diameter, 95b. *Oe. pringsheimii* var. *nordstedtii*
161. Female vegetative cell 14-20 μ in diameter, 95. *Oe. pringsheimii*
162. Diameter of oogonium 16-20 μ , 99a. *Oe. pisanum* var. *gracilis*
162. Diameter of oogonium 23-29 μ , 163
162. Diameter of oogonium 28-33 μ , 202. *Oe. sancti thomæ**
163. Oospore filling oogonium longitudinally, 164
163. Oospore not filling oogonium longitudinally, 97. *Oe. porrectum*
164. Basal cell elongate, 99. *Oe. pisanum*
164. Basal cell subhemispherical, 98. *Oe. nanum*
165. Oogonium pyriform to subpyriform, 166
165. Oogonium obovoid-globose to subglobose, 169
165. Oogonium ellipsoid to oblong, 177
166. Diameter of vegetative cell 8-11 μ , 167
166. Diameter of vegetative cell 11-13 μ , 102. *Oe. simplex*
166. Diameter of vegetative cell 13-16 μ , 100. *Oe. pyriforme*
167. Diameter of oogonium 23-30 μ , 168
167. Diameter of oogonium 30-33 μ , 103. *Oe. pyrulum*
167. Diameter of oogonium 34-40 μ , 103a. *Oe. pyrulum* var. *amplius*
168. Oospore subglobose, 22-26 $\times$ 21-24 μ , 104. *Oe. loricatum*
168. Oospore globose, 25-29 μ in diameter, 106. *Oe. pithophoræ*
169. Diameter of vegetative cell 8-16 μ , 170
169. Diameter of vegetative cell 16-20 μ , 107. *Oe. autumnale*
169. Diameter of vegetative cell 20-27 μ , 176
170. Oospore not filling oogonium, 171
170. Oospore filling oogonium, 172
171. Diameter of oogonium 30-38 μ , 109d. *Oe. crispum* var. *hawaiense*
171. Diameter of oogonium 40-43 μ , 108. *Oe. obesum*
172. Diameter of oogonium 23-28 μ , 104. *Oe. loricatum*
172. Diameter of oogonium 30-38 μ , 173
172. Diameter of oogonium 38-50 μ , 174
173. Vegetative cell 1-3 diameters long, 109e. *Oe. crispum* var. *uruguayense*
173. Vegetative cell 3-5 diameters long, 109c. *Oe. crispum* var. *gracilescens*
174. Oogonium obovoid-globose, 175
174. Oogonium subglobose, 109a. *Oe. crispum* f. *granulosum*
175. Diameter of oogonium 38-45 μ , 109. *Oe. crispum*
175. Diameter of oogonium 40-50 μ , 109b. *Oe. crispum* f. *inflatum*
176. Diameter of oogonium 44-49 μ , 110a. *Oe. rupestre* f. *pseudautumnale*
176. Diameter of oogonium 48-58 μ , 110. *Oe. rupestre*
177. Diameter of oogonium 9-14 μ , 201. *Oe. selandiae**
177. Diameter of oogonium 13-19 μ , 178
177. Diameter of oogonium 20-28 μ , 179
177. Diameter of oogonium 35-42 μ , 111. *Oe. ahlstrandii*
177. Diameter of oogonium 50-70 μ , 112. *Oe. pachydermum*
178. Length of oogonium 25-28 μ , 91. *Oe. trioicum*
178. Length of oogonium 34-40 μ , 114. *Oe. gracillimum*
179. Oospore nearly or quite filling oogonium, 180
179. Oospore not filling oogonium longitudinally, 181
180. Vegetative cell 7-9 $\times$ 11-36 μ , 115. *Oe. oblongellum*
180. Vegetative cell 8-15 $\times$ 16-60 μ , 113. *Oe. kirchneri*
181. Oospore oblong-ellipsoid, 114a. *Oe. gracillimum* f. *majus*
181. Oospore globose, 116b. *Oe. oblongum* f. *sphaericum*
181. Oospore ellipsoid, 182
182. Diameter of oogonium 20-26 μ , 116. *Oe. oblongum*
182. Diameter of oogonium 26-28 μ , 116a. *Oe. oblongum* f. *majus*
183. Diameter of oogonium 9-14 μ , 201. *Oe. selandiae**
183. Diameter of oogonium 28-33 μ , 202. *Oe. sancti thomæ**
183. Diameter of oogonium 35-44 μ , 204. *Oe. rhodosporum**
183. Diameter of oogonium 40-45 μ , 205. *Oe. vesicatum**
183. Diameter of oogonium 40-50 μ , 206. *Oe. montagnei**

184. Oogonium with mammiform projections.....118. *Oe. mammiferum*
 184. Oogonium with conically obtuse projections.....185
 184. Oogonium with projections intermediate between the two above,
 117a. *Oe. itzigsohnii* f. *heteromorphum*
 185. Diameter of vegetative cell 5-7 μ117b. *Oe. itzigsohnii* var. *minus*
 185. Diameter of vegetative cell 8-10 μ117. *Oe. itzigsohnii*

DIOECIOUS, NANNANDROUS, WITH CYLINDRICAL VEGETATIVE CELLS.

186. Oogonium opening by a pore.....187
 186. Oogonium opening by a lid.....228
 187. Pore median, or a little above median.....188
 187. Pore inframedian.....123a. *Oe. hystrix* var. *canadense*
 187. Pore supramedian.....201
 187. Pore superior.....202
 187. Pore inferior.....226
 188. Wall of oospore echinate.....189
 188. Wall of oospore punctate.....120. *Oe. nebraskense*
 188. Wall of oospore spirally ribbed.....194
 188. Wall of oospore smooth.....197
 189. Spines more than 5 μ in length.....124a. *Oe. echinospermum* var. *horridum*
 189. Spines less than 5 μ in length.....190
 190. Suffultory cell enlarged.....122. *Oe. hystricinum*
 190. Suffultory cell not or very slightly enlarged.....191
 191. Diameter of vegetative cell 7-13 μ121. *Oe. aster*
 191. Diameter of vegetative cell 12-16 μ125. *Oe. pungens*
 191. Diameter of vegetative cell 17-30 μ192
 192. Diameter of oogonium 36-38 μ123b. *Oe. hystrix* var. *subglobosum*
 192. Diameter of oogonium 38-50 μ193
 193. Oospore globose.....124. *Oe. echinospermum*
 193. Oospore ellipsoid.....123. *Oe. hystrix*
 194. Suffultory cell enlarged.....195
 194. Suffultory cell not enlarged.....196
 195. Oogonium 40-44 μ in diameter.....126. *Oe. exospirale*
 195. Oogonium 51-60 μ in diameter.....127. *Oe. illinoisense*
 196. Diameter of oospore 37-45 μ128a. *Oe. spirale* var. *acutum*
 196. Diameter of oospore 46-56 μ128. *Oe. spirale*
 197. Oospore globose to subglobose.....198
 197. Oospore sexangular-ellipsoid.....200
 197. Oospore ovoid.....132b. *Oe. braunii* var. *zehneri*
 198. Vegetative cell 8-9 μ in diameter.....129. *Oe. depressum*
 198. Vegetative cell 9-12 μ in diameter.....130. *Oe. semiapertum*
 198. Vegetative cell 12-23 μ in diameter.....199
 199. Diameter of oogonium 23-30 μ132a. *Oe. braunii* var. *hafniense*
 199. Diameter of oogonium 30-37 μ132. *Oe. braunii*
 199. Diameter of oogonium 43-48 μ131. *Oe. gallicum*
 199. Diameter of oogonium 49-52 μ133. *Oe. flavescens*
 200. Diameter of oogonium 19-30 μ135a. *Oe. sexangulare* var. *angulosum*
 200. Diameter of oogonium 29-33 μ135. *Oe. sexangulare*
 200. Diameter of oogonium 36-42 μ135b. *Oe. sexangulare* var. *majus*
 201. Oospore wall punctate.....120. *Oe. nebraskense*
 201. Oospore wall spirally ribbed.....128b. *Oe. spirale* var. *latviense*
 201. Oospore wall smooth.....143a. *Oe. multisporum* var. *magnum*
 201. Oospore wall echinate.....134. *Oe. manschuricum*
 202. Oospore wall smooth.....203
 202. Oospore wall dentate.....136. *Oe. stellatum*
 202. Oospore wall spirally ribbed.....128b. *Oe. spirale* var. *latviense*
 202. Oospore wall echinate.....215
 202. Oospore wall longitudinally ribbed.....218
 202. Oospore wall pitted.....225

*See footnote, page 53.

203.	Suffultory cell enlarged.....	204
203.	Suffultory cell not enlarged.....	207
	204. Diameter of oogonium 35-50 μ	205
	204. Diameter of oogonium 53-60 μ 138a. <i>Oe. borisianum</i> var. <i>westii</i>	
	204. Diameter of oogonium 65-80 μ	206
205.	Vegetative cell 12-15 $\times$ 60-120 μ	137. <i>Oe. silvaticum</i>
205.	Vegetative cell 15-23 $\times$ 45-150 μ	138. <i>Oe. borisianum</i>
	206. Suffultory cell 26-40 μ in diameter.....	139. <i>Oe. alternans</i>
	206. Suffultory cell about 44 μ in diameter.....	140. <i>Oe. kitulæ</i>
207.	Diameter of vegetative cell 10-20 μ	208
207.	Diameter of vegetative cell 21-40 μ	210
	208. Oogonium 24-35 μ in diameter.....	143. <i>Oe. multisporum</i>
	208. Oogonium 35-45 μ in diameter.....	209
209.	Oospore globose.....	144. <i>Oe. irregulare</i>
209.	Oospore depressed-globose.....	144a. <i>Oe. irregulare</i> var. <i>condensatum</i>
	210. Diameter of oogonium 39-60 μ	211
	210. Diameter of oogonium 65-80 μ	139. <i>Oe. alternans</i>
211.	Vegetative cell 1-3 diameters long.....	212
211.	Vegetative cell 3-6 diameters long.....	214
	212. Oospore not filling oogonium.....	141. <i>Oe. victoriense</i>
	212. Oospore nearly or quite filling oogonium.....	213
213.	Oogonium 39-54 $\times$ 42-65 μ	142a. <i>Oe. crassiusculum</i> var. <i>archavaletæ</i>
213.	Oogonium 55-60 $\times$ 60-75 μ	142c. <i>Oe. crassiusculum</i> var. <i>cataractum</i>
	214. Gynandrosporous.....	142. <i>Oe. crassiusculum</i>
	214. Idioandrosporous.....	142b. <i>Oe. crassiusculum</i> var. <i>idioandrosporum</i>
215.	Spines arranged spirally.....	216
215.	Spines not arranged spirally.....	217
	216. Diameter of antheridium 6-9 μ	136. <i>Oe. stellatum</i>
	216. Diameter of antheridium 9-10 μ	145a. <i>Oe. donnellii</i> var. <i>wittrockiana</i>
	216. Diameter of antheridium 14-15 μ	145. <i>Oe. donnellii</i>
217.	Diameter of oogonium 29-33 μ	146. <i>Oe. armigerum</i>
217.	Diameter of oogonium 35-36 μ	147. <i>Oe. echinatum</i>
217.	Diameter of oogonium 46-57 μ	148. <i>Oe. lindmanianum</i>
	218. Ribs 16-25 in number.....	219
	218. Ribs 25-40 in number.....	223
219.	Oogonium 45-50 μ in diameter.....	149a. <i>Oe. cyathigerum</i> f. <i>americanum</i>
219.	Oogonium 48-65 μ in diameter.....	220
219.	Oogonium 65-81 μ in diameter.....	149c. <i>Oe. cyathigerum</i> f. <i>perfectum</i>
	220. Oospore usually completely filling oogonium.....	221
	220. Oospore incompletely filling oogonium,.....	149d. <i>Oe. cyathigerum</i> var. <i>ellipticum</i>
221.	Dwarf male 47-58 μ in length.....	222
221.	Dwarf male 60-75 μ in length.....	149b. <i>Oe. cyathigerum</i> f. <i>ornatum</i>
	222. Diameter of oogonium 48-57 μ	149e. <i>Oe. cyathigerum</i> var. <i>hormosporum</i>
	222. Diameter of oogonium 57-66 μ	149. <i>Oe. cyathigerum</i>
223.	Ribs crenulate.....	150a. <i>Oe. wolleanum</i> var. <i>concinnum</i>
223.	Ribs entire.....	224
	224. Ribs 25-35 in number.....	150. <i>Oe. wolleanum</i>
	224. Ribs 35-40 in number.....	150b. <i>Oe. wolleanum</i> f. <i>insigne</i>
225.	Pits more or less in longitudinal rows.....	151. <i>Oe. concatenatum</i>
225.	Pits not in longitudinal rows.....	151a. <i>Oe. concatenatum</i> f. <i>hutchinsiae</i>
225.	Pits in vertical and transverse rows.....	151b. <i>Oe. concatenatum</i> var. <i>rectangulare</i>
	226. Oospore ribbed.....	152. <i>Oe. huntii</i>
	226. Oospore echinate.....	153. <i>Oe. hispidum</i>
	226. Oospore dentate.....	227
227.	Oogonium 47-60 $\times$ 48-63 μ	154. <i>Oe. cleveanum</i>
227.	Oogonium 39-46 μ $\times$ 40-47 μ	154a. <i>Oe. cleveanum</i> f. <i>exoticum</i>
	228. Division of oogonium median.....	229
	228. Division of oogonium suprmedian.....	233
	228. Division of oogonium inframedian.....	236
	228. Division of oogonium superior.....	240
	228. Division of oogonium supreme†.....	261

†Some forms of *Oe. ciliatum* may be sought here.

229.	Oogonium longitudinally plicate.....	230
229.	Oogonium not longitudinally plicate.....	231
230.	Diameter of oogonium 16-18 μ	233a. <i>Oe. crenulatum</i> var. <i>gracilius</i> *
230.	Diameter of oogonium 25-27 μ	233. <i>Oe. crenulatum</i> *
231.	Diameter of oogonium 13-16 μ	168. <i>Oe. longicolle</i>
231.	Diameter of oogonium 16-20 μ	168a. <i>Oe. longicolle</i> var. <i>senegalense</i>
231.	Diameter of oogonium 21-26 μ	169. <i>Oe. rothii</i>
231.	Diameter of oogonium 25-31 μ	232
231.	Diameter of oogonium 30-38 μ	171. <i>Oe. decipiens</i>
231.	Diameter of oogonium 39-46 μ	172. <i>Oe. macrospermum</i>
231.	Diameter of oogonium 89-100 μ	162. <i>Oe. perspicuum</i>
232.	Gynandrosporous.....	171c. <i>Oe. decipiens</i> var. <i>bernardense</i>
232.	Idioandrosporous.....	171b. <i>Oe. decipiens</i> var. <i>africanum</i>
233.	Oogonium with verticillate folds.....	234
233.	Oogonium without verticillate folds.....	235
234.	Diameter of oogonium 23-28 μ	229. <i>Oe. plicatulum</i> *
234.	Diameter of oogonium 50-55 μ	173. <i>Oe. costatum</i>
234.	Diameter of oogonium 58-65 μ	174. <i>Oe. boreale</i>
235.	Vegetative cell 4-7 μ in diameter.....	170. <i>Oe. danicum</i>
235.	Vegetative cell 13-24 μ in diameter.....	175. <i>Oe. mirandrium</i>
236.	Oogonium plicate.....	237
236.	Oogonium not plicate.....	176. <i>Oe. contortum</i>
237.	Suffultory cell enlarged.....	238
237.	Suffultory cell not enlarged.....	239
238.	Diameter of oogonium 23-26 μ	232. <i>Oe. pulchrum</i> *
238.	Diameter of oogonium 28-32 μ	167b. <i>Oe. platygynum</i> var. <i>continuum</i>
239.	Oogonium 16-23 μ in diameter.....	230. <i>Oe. lagerstedtii</i> *
239.	Oogonium 23-29 μ in diameter.....	231. <i>Oe. uleanum</i> *
240.	Oospore wall smooth.....	241
240.	Oospore wall pitted.....	177. <i>Oe. schmidlei</i>
240.	Oospore wall echinate.....	212. <i>Oe. velatum</i> *
240.	Oospore wall scrobiculate.....	258
241.	Vegetative cell 3-8 μ in diameter.....	242
241.	Vegetative cell 8-32 μ in diameter.....	245
242.	Diameter of oogonium 14-20 μ	243
242.	Diameter of oogonium 19-23 μ	179b. <i>Oe. rugulosum</i> f. <i>rotundatum</i>
243.	Oospore 15-20 μ in length.....	244
243.	Oospore 19-25 μ in length.....	179. <i>Oe. rugulosum</i>
244.	Oogonium ovoid to ellipsoid.....	178. <i>Oe. longatum</i>
244.	Oogonium obovoid to ovoid-ellipsoid,	179a. <i>Oe. rugulosum</i> f. <i>minutum</i>
245.	Oospore globose or subglobose.....	246
245.	Oospore ellipsoid or subellipsoid.....	253
246.	Suffultory cell enlarged.....	247
246.	Suffultory cell not enlarged.....	248
247.	Diameter of vegetative cell 9-15 μ	180. <i>Oe. hians</i>
247.	Diameter of vegetative cell 13-20 μ	175. <i>Oe. mirandrium</i>
248.	Vegetative cell $\frac{3}{4}$ to $2\frac{1}{2}$ diameters long.....	249
248.	Vegetative cell 2 to 3 diameters long,	184e. <i>Oe. macrandrium</i> var. <i>kohenackerii</i>
248.	Vegetative cell $2\frac{1}{2}$ to 6 diameters long.....	250
248.	Vegetative cell 6 to 14 diameters long.....	181. <i>Oe. hoersholmiense</i>
249.	Diameter of vegetative cell 12-14 μ	182. <i>Oe. laetevirens</i>
249.	Diameter of vegetative cell 13-21 μ	184c. <i>Oe. macrandrium</i> var. <i>lundense</i>
249.	Diameter of vegetative cell 21-29 μ	183. <i>Oe. pluviale</i>
250.	Length of oogonium 28-44 μ	251
250.	Length of oogonium 44-55 μ	252
251.	Oogonium usually single or in two's.....	184d. <i>Oe. macrandrium</i> var. <i>propinquum</i>
251.	Oogonium usually in groups of 2 to 6.....	184b. <i>Oe. macrandrium</i> f. <i>aemulans</i>
252.	Oospore 31-37 $\times$ 33-39 μ	184. <i>Oe. macrandrium</i>
252.	Oospore 35-40 $\times$ 36-45 μ	184a. <i>Oe. macrandrium</i> f. <i>acuminatum</i>

*See footnote, page 53.

253.	Diameter of vegetative cell 7-12 μ	181.	<i>Oe. hoersholmiense</i>	
253.	Diameter of vegetative cell 14-32 μ			254
254.	Terminal cell setiform.....	185.	<i>Oe. ciliatum</i>	
254.	Terminal cell not setiform.....			255
255.	Diameter of oospore 27-32 μ	207.	<i>Oe. flexuosum</i> *	
255.	Diameter of oospore 32-42 μ			256
255.	Diameter of oospore 41-48 μ	186.	<i>Oe. spectabile</i>	
256.	Vegetative cell $\frac{3}{4}$ to 3 diameters long.....			257
256.	Vegetative cell 3 to 5 diameters long.....	187.	<i>Oe. implexum</i>	
257.	Oogonium globose-obovoid.....	208.	<i>Oe. fonticola</i> *	
257.	Oogonium subglobose.....	183.	<i>Oe. pluviale</i>	
258.	Diameter of oogonium 30-39 μ			259
258.	Diameter of oogonium 40-50 μ			260
258.	Diameter of oogonium 56-68 μ	188b.	<i>Oe. monile</i> var. <i>eminens</i>	
259.	Suffultory cell tumid.....	188.	<i>Oe. monile</i>	
259.	Suffultory cell not tumid.....	184f.	<i>Oe. macrandrium</i> var. <i>scrobiculatum</i>	
260.	Suffultory cell 27-28 μ in diameter.....	188c.	<i>Oe. monile</i> f. <i>victoriense</i>	
260.	Suffultory cell 35-38 μ in diameter.....	188a.	<i>Oe. monile</i> f. <i>borgei</i>	
261.	Oospore wall smooth.....			262
261.	Oospore wall longitudinally ribbed.....			264
262.	Diameter of oogonium 30-36 μ	181.	<i>Oe. hoersholmiense</i>	
262.	Diameter of oogonium 42-55 μ			263
262.	Diameter of oogonium 55-58 μ	189a.	<i>Oe. obtruncatum</i> var. <i>completum</i>	
263.	Length of oogonium 56-68 μ	189.	<i>Oe. obtruncatum</i>	
263.	Length of oogonium 68-75 μ	189b.	<i>Oe. obtruncatum</i> var. <i>ellipsoideum</i>	
264.	Ribs 11-17 in number.....	195b.	<i>Oe. acrosporum</i> var. <i>bathmidosporum</i>	
264.	Ribs 23-30 in number.....			265
264.	Ribs 40-45 in number.....	193.	<i>Oe. tentoriale</i>	
265.	Vegetative cell 7-8 μ in diameter.....	195c.	<i>Oe. acrosporum</i> var. <i>floridense</i>	
265.	Vegetative cell 12-21 μ in diameter.....			266
266.	Plant usually less than 10-celled.....	195a.	<i>Oe. acrosporum</i> f. <i>boreale</i>	
266.	Plant usually more than 15-celled.....			267
267.	Diameter of oogonium 35-48 μ	195.	<i>Oe. acrosporum</i>	
267.	Diameter of oogonium 44-56 μ	195d.	<i>Oe. acrosporum</i> var. <i>majusculum</i>	
268.	Oospore wall smooth.....			269
268.	Oospore wall punctate.....	41c.	<i>Oe. fabulosum</i> var. <i>punctatum</i>	
269.	Vegetative cell 56-70 μ in diameter.....	41b.	<i>Oe. fabulosum</i> var. <i>columbianum</i>	
269.	Vegetative cell 70-93 μ in diameter.....			270
270.	Oospore globose to ellipsoid.....	41.	<i>Oe. fabulosum</i>	
270.	Oospore subglobose to subcylindrical,			
	41a. <i>Oe. fabulosum</i> var. <i>maximum</i>			

*See footnote, page 53.

1. *Oe. laeve* Wittrock.

(Pl. XI, fig. 100.)

1874, p. 8; De Toni 1889, p. 35; Hirn 1900, p. 75.

Monoecious; oogonium single,* depressed-globose, pore median; oospore depressed-globose, filling oogonium, spore wall smooth; antheridium in groups of 1-2,* subepigynous, sperm 1; vegetative cell $10-14 \times 20-70\mu$; oogonium $32-38 \times 24-30\mu$; oospore $30-35 \times 23-26\mu$; antheridium $9-10 \times 9-13\mu$.

United States: Illinois, Michigan; France.

Originally distributed from France, the species was found subsequently in Illinois and Michigan (Transeau), and the above description is made from the American material. The large oogonium in relation to the diameter of the vegetative cell and the depressed-globose oospores are chief characteristics. It is clearly separated from *Oe. curvum*, in addition, by the absence of any marked tendency toward irregularly curved filaments.

2. *Oe. curvum* Pringsheim.

(Pl. XI, figs. 103 and 104.)

1858, p. 69, Pl. 5, fig. 3; Hirn 1900, p. 74, Pl. I, fig. 3; Tiffany 1926, p. 99, Pl. III, figs. 25-27.

Monoecious; oogonium 1-6, depressed globose, pore median; oospore depressed-globose, filling or not filling oogonium, spore wall smooth; antheridium 1-7, sperm 1; basal cell elongate; filament usually irregularly curved; vegetative cells $5-10 \times 10-40\mu$; oogonium $21-25 \times 18-24\mu$; oospore $19-23 \times 14-19\mu$; antheridium $6-9 \times 6-9\mu$.

United States: Iowa, Alabama; Germany, Spain, Sweden.

This species is to be compared with *Oe. cryptoporum* below. The irregularly curved filaments, particularly in the region of the antheridia, are usually characteristic. The pore is sometimes a little above the median position.

3. *Oe. cryptoporum* Wittrock.

(Pl. XI, fig. 101.)

1870, p. 119; Hirn 1900, p. 72, Pl. I, fig. 1; P. B. A. No. 1525.

Monoecious; oogonium 1, subdepressed-obovoid-globose or subdepressed-globose, pore median; oospore subdepressed-globose, quite filling oogonium, spore wall smooth; antheridium 1-7, scattered or subhypogynous or subepigynous; sperm 1; vegetative cell $7-10 \times 28-60\mu$; oogonium $23-28 \times 26-31\mu$; oospore $22-27 \times 19-21\mu$; antheridium $6-8 \times 7-11\mu$.

United States: Utah; Austria, Denmark, Norway, Sweden.

*Hereafter a numeral (or numerals) only will indicate the number of oogonia and antheridia; thus 1=a single oogonium or antheridium and 2-7 means that oogonia (or antheridia) may occur in groups of from two to seven seriate individuals. The same applies to androsporangia.

3a. Var. **vulgare** Wittrock.

(Pl. XI, fig. 102.)

1874, p. 7; Wolle 1887, p. 70, Pl. 74, figs. 1 and 2; *Oe. cryptoporum* Wittr. in Hirn 1896, p. 2; Hirn 1900, p. 73, Pl. I, fig. 2; (?) *Oe. spetsbergense* Wittr. 1874, p. 37 and in Hirn 1900, p. 316, Pl. L, fig. 322.

Smaller; oogonium 1-5; vegetative cell $5-8 \times 15-48\mu$; oogonium $18-25 \times 18-26\mu$; oospore $16-23 \times 15-19\mu$; antheridium $5-7 \times 9-12\mu$.

United States: New Jersey, Michigan, Illinois, Massachusetts, Pennsylvania, Utah; Australia; Ko Chang; Ceylon; Austria, England, France, Germany, Italy, Spain, Norway, Sweden.

The variety is much more widely distributed and has smaller vegetative cells and shorter oogonia than the species. The small sized filaments with rather depressed-globose oogonia are characteristic.

4. **Oe. rufescens** Wittrock.

(Pl. XI, fig. 105.)

1870, p. 134; 1874, p. 32; Hirn 1900, p. 76, Pl. I, fig. 4; Collins 1909, p. 229; P. B. A. No. 521.

Dioecious, macrandrous; oogonium 1-3, obovoid- or depressed-obovoid-globose, pore median, rimiform; oospore globose or depressed-globose, filling oogonium or nearly so, spore wall smooth; antheridium to 12-seriate; sperm 1; female vegetative cell $8-10 \times 34-70\mu$, male $7-9 \times 30-54\mu$; oogonium $22-24 \times 22-30\mu$; oospore $21-23 \times 17-22\mu$; antheridium $6-8 \times 8-12\mu$.

United States: Connecticut, Rhode Island, Michigan, Illinois, Alabama, Ohio; Austria, Denmark, England, Germany, Sweden, Latvia; South Africa.

4a. Var. **exiguum** (Elfving) Tiffany nov. comb.

(Pl. XI, fig. 106.)

Oe. rufescens f. *exiguum* (Elfv.) Hirn 1900, p. 76, Pl. 1, fig. 5; *Oe. exiguum* Elfv. in Hirn 1895, p. 19.

Vegetative cells a little smaller; oospore subdepressed-globose; antheridium 3; vegetative cell $5-9 \times 22-88\mu$; oogonium $22-24 \times 20-28\mu$; oospore $20-22 \times 17-23\mu$; antheridium $5 \times 10-12\mu$.

United States: Illinois, Alabama, Mississippi; Finland, France; Southern Tibet.

4b. Var. **lundellii** (Wittrock) Tiffany.

(Pl. XI, fig. 107.)

1929, p. 75; *Oe. Lundellii* Wittr. 1874, p. 32; *Oe. Rothii* Breb. in Rabenh. Alg. Eur. No. 1374, 1862; *Oe. rufescens* Wittr. subsp. *Lundellii* (Wittr.) Hirn 1900, p. 77, Pl. I, fig. 6; *Oe. rufescens* Wittr. subsp. *Lundellii* (Wittr.) Hirn f. *elongatum* Hirn 1900, p. 78, Pl. I, fig. 8; P. B. A. No. 1428.

Vegetative cells wider in proportion to oogonium than in type; oogonium depressed-globose; vegetative cell $8-12 \times 30-75\mu$; oogonium $22-25 \times 21-27\mu$; oospore $19-22 \times 15-22\mu$; antheridium $7-9 \times 6-12\mu$.

United States: Maine, California; Austria, England, Finland, France, Germany, Hungary, Sweden.

The species and the varieties resemble *Oe. cryptoporum* (No. 3) except in their dioecious habit and are separated from each other chiefly on the basis of relative dimensions. The varieties are so widely distributed and so uniformly separable from the species that they must be regarded as distinct even though similar.

5. *Oe. calcareum* Cleve.

(Pl. XI, fig. 108.)

In Wittrock 1870, p. 135; (?) *Vesiculifera compressa* Hassall 1845, p. 204, Pl. 53, fig. 4; Hirn 1900, p. 78, Pl. I, fig. 9.

Dioecious, macrandrous; oogonium 1 or rarely 2, depressed-globose, with median pore; oospore depressed-globose, filling oogonium, wall smooth; antheridium to 8-seriate; sperm 1; vegetative cell $11-14 \times 22-56\mu$; oogonium $27-30 \times 21-23\mu$; oospore $26-28 \times 20-21\mu$; antheridium $10-11 \times 9-12\mu$.

England, Denmark, Sweden, Latvia; South Africa.

The filaments of this species are often incrustated with lime. The depressed-globose oospore completely filling the oogonium and the robust appearance of the short-celled vegetative filaments are characteristic.

6. *Oe. sociale* Wittrock.

(Pl. XI, fig. 109.)

In Wittr. & Nordst. Alg. Exs. No. 401, 1882; *Oe. ochroleucum* Kuetz. in Rabenh. Alg. Eur. No. 1280, 1862; Hirn 1900, p. 79, Pl. II, fig. 12.

Dioecious, macrandrous; oogonium 1, subglobose, pore median; oospore globose (or subglobose), quite filling oogonium, cell wall smooth; antheridium 1-5, sperms 2, division horizontal; vegetative cell $9-16 \times 30-130\mu$; oogonium $30-38 \times 33-42\mu$; oospore (26-) $28-35 \times 28-35\mu$; antheridium $12-14 \times 8-11\mu$.

United States: Illinois, Alabama, Mississippi; British Columbia; Austria, Germany, Sweden, Latvia; India, Tibet, Burma.

The oospore is usually quite globose. It is easily separable from its dioecious relatives by its larger size and by the presence of two sperms in the antheridium.

7. *Oe. cymatosporum* Wittrock and Nordstedt.

(Pl. XII, figs. 112 and 113.)

In Wittr. 1870, p. 121; *Oe. Magnusii* Wittr. in Hirn 1895, p. 11; Hirn 1900, p. 80, Pl. II, fig. 13.

Monoecious; oogonium 1 or rarely 2, subdepressed-globose, pore median, rimiform; oospore depressed-globose, filling or not filling oogonium, spore wall of three layers: outer smooth, middle scrobiculate,

inner smooth; antheridium 1-4, subepigynous, subhypogynous, hypogynous, or scattered; sperm 1; vegetative cell $8-10 \times 32-70\mu$; oogonium $30-40 \times 27-40\mu$; oospore $27-35 \times 22-33\mu$; antheridium $7-10 \times 9-15\mu$.

United States: Massachusetts, Mississippi; Austria, England, Finland, France, Hungary, Sweden.

This species resembles *Oe. magnusii* below from which it is distinguished by its monoecious habit, its larger oogonia, and its oospores not usually filling the oogonia.

8. *Oe. magnusii* Wittrock.

(Pl. XII, fig. 115.)

1874, p. 38; Hirn 1900, p. 81, Pl. II, fig. 14; Wittr. and Nordst. Alg. Ex. No. 109.

Diocious, macrandrous (perhaps monoecious also); oogonium 1-3, depressed-globose, pore median, rimiform; oospore depressed-globose, quite filling oogonium, spore wall of three layers: outer layer smooth, middle layer scrobiculate, inner layer smooth; antheridium to 8-seriate, sperm 1; vegetative cell $7-10 \times 12-40\mu$; oogonium $24-27 \times 21-26\mu$; oospore $22-25 \times 18-23\mu$; antheridium $8-10 \times 5-11\mu$.

United States: Illinois, Michigan, Massachusetts; British Columbia; Finland, Germany, Sweden.

Easily separable from *Oe. cymatosporum* (No. 7). According to Wittrock it is sometimes monoecious.

9. *Oe. suecicum* Wittrock.

(Pl. XII, fig. 116.)

1872, p. 5; 1874, p. 30; West 1891, p. 109, Pl. 18, fig. 2; (?) *Oe. trichosporum* Itzig. in Rabenh. 1868, p. 426; Hirn 1900, p. 82, Pl. II, fig. 15; Collins 1909, p. 230; Heering 1914, p. 193, fig. 281; Tiffany 1926, p. 89, Pl. V, figs. 57 and 58.

Diocious, macrandrous; oogonium 1, subglobose, pore median; oospore globose, nearly filling oogonium, outer spore wall echinate, inner smooth; antheridium 2-6, sperm 1; basal cell elongate; terminal cell apically obtuse; vegetative cell $9-14 \times 30-90\mu$; oogonium $32-38 \times 34-41\mu$; oospore $30-37 \times 30-37\mu$; antheridium $10-12 \times 13-17\mu$.

United States: Iowa, Illinois, Massachusetts; Australia; Austria, Denmark, Finland, France, Germany, Latvia, Iceland, Norway, Sweden; British Columbia; South Africa.

9a. Form *australe* G. S. West.

(Pl. XII, fig. 117.)

1909, p. 45, fig. 6 E.

Vegetative cell somewhat larger; spines larger and more sparse; vegetative cell $14-16 \times 56-80\mu$; oogonium $40-41 \times 40-41\mu$; oospore (without spines) $31-33 \times 31-33\mu$.

Africa: Yan Yean Reservoir, Victoria.

The species and the variety constitute the only dioecious macrandrous members of the genus having echinate oospores.

10. *Oe. varians* Wittrock and Lundell.

(Pl. XII, fig. 120.)

In Wittr. 1874, p. 11; *Oe. polymorphum* Wölle 1887, p. 73, Pl. LXXIV, figs. 16-19; Hirn 1900, p. 89, Pl. IV, fig. 23; Collins 1909, p. 232; Heering 1914, p. 206, fig. 298; Tiffany 1926, p. 99, Pl. VII, figs. 75 and 76.

Monoecious, (or sometimes dioecious); oogonium 1, or rarely more, depressed- or subdepressed-pyriform-globose, pore nearly superior; oospore globose, not filling oogonium, spore wall smooth; antheridium to 9-seriate, scattered; sperms 2, division horizontal; basal cell elongate; terminal cell apically obtuse; vegetative cell $12-16 \times 35-144 \mu$; oogonium $34-50 \times 34-55 \mu$; oospore $31-41 \times 30-41 \mu$; antheridium $11-15 \times 5-7 \mu$.

United States: Iowa, New Jersey, Illinois; Patagonia; Austria, Finland, Sweden; South Africa.

The typical plants of this species have a nearly superior pore, and the oospore usually does not fill the irregularly shaped oogonium. Transeau has observed both dioecious and monoecious forms from Illinois. (See also Hirn *loc. cit.*).

11. *Oe. urbicum* Wittrock.

(Pl. XII, figs. 118 and 119.)

1874, p. 13; *Oe. tumidulum* Pringsh. 1855, p. 158, Pl. 1, figs. 26 and 27; Hirn 1900, p. 91, Pl. V, fig. 26.

Monoecious; oogonium 1, ellipsoid-globose, pore a little above median or nearly superior; oospore globose, not filling oogonium, spore wall smooth; antheridium few celled; sperms 2, division horizontal; vegetative cell $15-19 \times 40-110 \mu$; oogonium $46-55 \times 58-63 \mu$; oospore $33-46 \times 33-46 \mu$; antheridium $14-16 \times 5-7 \mu$.

Germany.

In general appearance this species is similar to *Oe. varians* above and *Oe. tyrolicum* (No. 14). It is distinguished from the former by its size and tendency toward ellipsoid oogonia; from both by the variable position of the pore.

12. *Oe. obsoletum* Wittrock.

(Pl. XII, fig. 121.)

1874, p. 9; *Oe. vernale* Wittr. 1872, p. 1; Hirn 1900, p. 83, Pl. II, fig. 16.

Monoecious; oogonium 1, subglobose or subdepressed-globose, pore a little above supramedian; oospore subdepressed-globose, not filling oogonium, spore wall smooth; antheridium 1-3, subepigynous; sperm 1; vegetative cell $9-15 \times 30-75 \mu$; oogonium $34-39 \times 34-43 \mu$; oospore $30-34 \times 28-32 \mu$; antheridium $8-9 \times 12-16 \mu$.

United States: Pennsylvania; England, Sweden.

Similar to *Oe. plusiosporum* below and distinguished from it by the usually depressed-globose appearance of the oospore. The oospore of *Oe. plusiosporum* is more regularly globose and its oogonium is more nearly ellipsoid.

13. *Oe. plusiosporum* Wittrock.

(Pl. XII, fig. 122.)

1874, p. 11; Wolle 1887, p. 72, Pl. 74, figs. 20 and 21; Hirn 1900, p. 84, Pl. II, fig. 17.

Monoecious; oogonium 1, subglobose or subellipsoid-globose, with pore a little above median; oospore globose, rarely subglobose, not quite filling oogonium, spore wall smooth; antheridium to 6-seriate; sperm (?) 1; terminal cell obtuse; vegetative cell $12-19 \times 24-77\mu$; oogonium (28-) $34-45 \times 35-50\mu$; oospore (25-) $30-39 \times 30-39\mu$; antheridium $12-14 \times 8-12\mu$.

United States: Pennsylvania, New Jersey, California; Columbia; British Columbia; Sweden, Finland, Rumania.

See *Oe. obsoletum* above.

14. *Oe. tyrolicum* Wittrock.

(Pl. XII, fig. 123.)

1874, p. 12; Hirn 1900, p. 91, Pl. IV, fig. 25; Heering 1914, p. 206, fig. 299; Collins 1918, p. 65; Tiffany 1926, p. 99, Pl. VII, fig. 77.

Monoecious; oogonium 1 (rarely 2), ellipsoid-globose or occasionally obovoid, pore nearly superior; oospore usually globose, not filling oogonium, spore wall smooth; antheridium 1-4; sperms 2, division horizontal; basal cell elongate; terminal cell apically obtuse; vegetative cell $15-24 \times 45-120\mu$; oogonium $45-53 \times 57-70\mu$; oospore $40-48 \times 40-48\mu$; antheridium $12-21 \times 9-11\mu$.

United States: Massachusetts, Iowa; Austria.

Distinguished by its more nearly ellipsoid oogonia and globose oospores. Separated from *Oe. varians* (No. 10) by its larger dimensions and from *Oe. cardiacum* var. *carbonicum* (No. 15c) by its monoecious habit.

15. *Oe. cardiacum* (Hassall) Wittrock.

(Pl. XIII, figs. 124 and 125.)

Vesiculifera cardiaca Hass. 1845, p. 203, Pl. 51, fig. 4; *Oe. cardiacum* Wittr. 1870, p. 135; *Oe. lautumnarum* Wittr. in Hirn 1896, p. 3; (?) *Pringsheimia inaequalis* Wood 1872, p. 195, Pl. 18, fig. 1; Hirn 1900, p. 85, Pl. III, fig. 19; Collins 1909, p. 231; Heering 1914, p. 194, fig. 273; Tiffany 1926, p. 89.

Dioecious, macrandrous; oogonium usually 1, subglobose to subcordiform-globose, with a supramedian pore; oospore globose, not filling the oogonium, spore wall smooth; antheridium 1-10; sperms 2, division horizontal; basal cell elongate; terminal cell apically obtuse; female vegetative cell $18-30 \times 60-200\mu$, male $15-25 \times 45-170\mu$; oogonium $48-70 \times 58-78\mu$; oospore $42-60 \times 42-60\mu$; antheridium $15-21 \times 10-14\mu$.

United States: Iowa, Illinois, Pennsylvania, Oregon, Ohio, Connecticut; Africa; Paraguay; Ontario; Australia; England, Denmark, Finland, France, Germany, Switzerland, Sweden.

15a. Form **interjectum** Hirn.

(Pl. XIII, fig. 127.)

1900, p. 86, Pl. III, fig. 21.

Pore of oogonium more nearly superior than in the type; vegetative cell $19-29 \times 60-190\mu$; oogonium $45-60 \times 63-81\mu$; oospore $44-58 \times 43-58\mu$.

England, Sweden.

15b. Form **pulchellum** (Hassall) Hirn.

(Pl. XIII, fig. 126.)

1900, p. 86, Pl. III, fig. 20; *Vesiculifera pulchella* Hass. 1845, p. 199, Pl. 50, fig. 3.

Vegetative cell $25-37 \times 40-125\mu$; oogonium $57-73 \times 68-78\mu$; oospore $56-66 \times 56-66\mu$.

England, France.

15c. Var. **carbonicum** Wittrock.

(Pl. XIII, fig. 128.)

In Wittr. and Nordst. Exs. 1883; *Oe. carbonicum* Wittr. 1874, p. 33; Hirn 1900, p. 87, Pl. IV, fig. 22; Tiffany 1926, p. 90, Pl. I, fig. 5; P. B. A. No. 1781.

Oogonium 1-2, obovoid to globose-ovoid, pore nearly superior; oospore obovoid-ellipsoid to globose-ellipsoid; female vegetative cell $14-30 \times 40-200\mu$, male $14-25 \times 40-150\mu$; oogonium $42-56 \times 50-80\mu$; oospore $40-52 \times 46-65\mu$; antheridium $13-20 \times 12-15\mu$.

United States: Iowa, Illinois, Ohio; Austria, England, Denmark, Germany, Sweden.

15d. Var. **minus** Lemmermann.

(Pl. XIII, fig. 129.)

1909, p. 191, as *v. minor*; Heering 1914, p. 194; Tiffany 1926, p. 90, Pl. I, fig. 8.

Oogonium and oospore smaller; female vegetative cell $16-28 \times 25-75\mu$, male $14-23 \times 25-75\mu$; oogonium $30-45 \times 35-50\mu$; oospore $28-42 \times 28-40\mu$; antheridium $12-15 \times 5-6\mu$.

United States: Iowa, Kentucky; Germany.

The "cardiacum" group is rather variable, and the varieties are not always clearly separated. The differentiating characters used in the key are those including shape and size of oospore and relative lengths of vegetative cells. Upon these bases the divisions above seem tenable. See also *Oe. glabrum* (No. 17).

16. **Oe. franklinianum** Wittrock.

(Pl. XIII, fig. 131.)

In Wittr. and Nordst. Alg. Exs. No. 309, 1880; Wille 1887, p. 89, figs. 7-9; Hirn 1900, p. 88, Pl. II, fig. 18; Collins 1909, p. 231; Heering 1914, p. 194; Tiffany 1926, p. 89, Pl. IV, figs. 43-44.

Diocious, macrandrous; oogonium 1, subglobose, with a supra-median pore; oospore globose, almost filling oogonium, spore wall

smooth; antheridium 1-4; sperms 2, division horizontal; female vegetative cell $9-12 \times 30-95\mu$, male $8-10 \times 25-90\mu$; oogonium $26-31 \times 29-41\mu$; oospore $24-30 \times 24-30\mu$; antheridium $8-9 \times 5-7\mu$.

United States: Iowa, Illinois, New Jersey, Pennsylvania; Brazil; Australia.

This species also resembles *Oe. cardiacum* (No. 15) but is distinguished by its smaller dimensions throughout.

17. *Oe. glabrum* Hallas.

(Pl. XIII, fig. 130.)

1905, p. 408, fig. 18; Hirn 1906, p. 14, Pl. I, fig. 2.

Dioecious, macrandrous; oogonium 1 (rarely 2), obovoid, pore superior; oospore globose or subglobose, not filling oogonium, spore wall smooth; antheridium 5-8; sperms 2, division vertical; terminal cell (sometimes the oogonium) broadly apiculate; female vegetative cell $16-26 \times 120-240\mu$, male $21-25 \times 60-150\mu$; oogonium $56-58 \times 63-86\mu$; oospore $42-49 \times 44-63\mu$; antheridium $20-22 \times 7-9\mu$.

Denmark.

This species in general appearance is near the "cardiacum" group. Its vertical division of the two sperms, the small size of its oospore in comparison to the oogonium, and the uniformly superior pore are characters that identify it and make it tenable as a species.

18. *Oe. lautumniarum* Wittrock.

(Pl. XIV, figs. 132 and 133.)

In Wittr. and Nordst. Alg. Exs. No. 7, 1877; Hirn 1900, p. 92, Pl. V, fig. 27.

Dioecious, macrandrous; oogonium 1, rarely 2, subobovoid-globose, pore a little above median (rarely nearly superior); oospore subglobose, or sometimes subangular, filling oogonium, wall smooth, rarely thickened; antheridium 1-4; female vegetative cell $16-22 \times 40-110\mu$, male $15-20 \times 45-100\mu$; oogonium $40-49 \times 45-51\mu$; oospore $36-46 \times 35-47\mu$; antheridium $14-17 \times 7-10\mu$.

British Columbia; Finland, Germany, Sweden, Latvia.

This species is separable from the "cardiacum" group largely by the varying position of the oogonial pore (from nearly median to nearly superior) and by a combination of cell sizes and shapes not found in any one of this group.

19. *Oe. intermedium* Wittrock.

(Pl. XIV, fig. 134.)

In Wittr. and Nordst. Alg. Exs. No. 708, 1886; *Oe. fragile* Wittr. in *ibid.* No. 4, 1877; *Oe. vaucherii* (Le Cl.) Al. Br. in *ibid.* No. 14, 1877, and in Rabenh. Alg. Eur. No. 2499, 1877; Hirn 1900, p. 94, Pl. V, fig. 31.

Monoecious; oogonium 1, obovoid to obovoid-globose, pore superior; oospore globose or obovoid-globose, filling oogonium or nearly so, spore

wall smooth and thick; antheridium 1-4, subepigynous or hypogynous or rarely scattered, sperms 2, division horizontal; vegetative cell $15-18 \times 45-80\mu$; oogonium $31-37 \times 34-45\mu$; oospore $30-36 \times 33-41\mu$; antheridium $14-16 \times 5-10\mu$.

United States: Illinois, Ohio, Mississippi; Austria, Finland, Sweden, Latvia, Germany; India; Siberia.

19a. Var. **fennicum** Tiffany.

(Pl. XIV, fig. 135.)

1929, p. 74; *Oe. intermedium* Wittr. forma *valida* Hirn 1900, p. 95, Pl. V, fig. 32; *Oe. intermedium* Wittr. forma West 1909a, p. 239.

Oogonium and oospore considerably larger; vegetative cell $17-19 \times 50-120\mu$; oogonium $38-46 \times 42-60\mu$; oospore $35-40 \times 35-40\mu$; antheridium $13-18 \times 6-8\mu$.

United States: Michigan; Finland; Egypt.

Oe. intermedium resembles *Oe. fragile* (No. 24) in general appearance, particularly those specimens that reach the upper limits of dimensions. It is rather intermediate between *Oe. fragile* and *Oe. globosum* (No. 21). Its size and location of antheridia are distinctive characters in this monoecious group with regularly superior pore. The variety *fennicum* is larger and its oogonium is not filled by the oospore.

20. **Oe. hirnii** Gutwinski.

(Pl. XIV, figs. 136 and 137.)

1896, p. 2, Pl. 5, fig. 1; Hirn 1900, p. 93, Pl. V, fig. 29; West and West 1902, p. 12, Pl. I, fig. 1-3.

Monoecious; oogonium 1, subglobose or subobovoid, with superior pore; oospore globose, not filling oogonium, spore wall smooth; antheridium 1-2, subepigynous; sperms 2, division horizontal; vegetative cell (sometimes slightly capitellate) $8-13 \times 28-80\mu$; oogonium $32-37 \times 32-39\mu$; oospore $28-31 \times 28-31\mu$; antheridium $8-11 \times 4-9\mu$.

Austria, Ireland.

20a. Var. **africanum** G. S. West.

(Pl. XIV, fig. 138.)

1907, p. 98; Tiffany 1926, p. 100, Pl. III, fig. 24.

Oogonium globose to subglobose; antheridium 1-3; vegetative cell, broadly capitellate, $13-15 \times 33-60\mu$; oogonium $39-40 \times 38-40\mu$; oospore $33-36 \times 33-36\mu$; antheridium $13-14 \times 5-6\mu$.

United States: Iowa; Africa.

The species is near *Oe. globosum* below in description, differing in its smaller size and its commonly subobovoid oogonia. The variety *africanum* is very distinct and is the only poriferous form in the monoecious group of the genus having broadly capitellate cells.

21. *Oe. globosum* Nordstedt.

(Pl. XIV, fig. 139.)

1878, p. 20, Pl. 2, fig. 16; Hirn 1900, p. 94, Pl. V, fig. 30.

Monoecious; oogonium 1, globose or subglobose, pore superior; oospore globose, quite filling oogonium, spore wall smooth; antheridium 1-7, subepigynous or subhypogynous or scattered; sperms 2, division horizontal; basal cell elongate; terminal cell setiform; vegetative cell $10-14 \times 40-95\mu$; oogonium $32-40 \times 32-46\mu$; oospore $30-37 \times 28-37\mu$; antheridium $9-12 \times 4-8\mu$.

United States: Illinois, Massachusetts; Australia; Greece, Serbia.

The relation of this species to *Oe. hirnii* (No. 20) and to *Oe. intermedium* (No. 19) is discussed under these latter forms.

22. *Oe. zigzag* Cleve.

(Pl. XV, figs. 143 and 144.)

In Wittr. 1870, p. 120; Hirn 1900, p. 101, Pl. VII, fig. 42.

Monoecious; oogonium 1, globose or obovoid-globose, pore superior; oospore of same form as oogonium and quite filling it, spore wall smooth, thick; antheridium 1, subepigynous or subhypogynous, alternating with oogonium and vegetative cell; sperms 2, division horizontal; basal cell elongate; terminal cell apically obtuse; vegetative cell $15-20 \times 30-80\mu$; oogonium $45-63 \times 52-63$ (-67) μ ; oospore $43-60 \times (44-) 48-58$ (-63) μ ; antheridium $15-19 \times 8-15\mu$.

Sweden.

The short, few-celled filaments of *Oe. zigzag* are quite unmistakable. It bears some resemblance to the dioecious *Oe. alternans* (No. 139) and is perhaps closely related to *Oe. curtum* (No. 23). From the latter it is distinguished by its shorter filaments, by its single oogonium, and by its single antheridium alternating with a vegetative cell.

23. *Oe. curtum* Wittrock and Lundell.

(Pl. XV, fig. 149.)

In Wittr. 1870, p. 121; *Oe. curvum* Pringsh. in Roumeguere Alg. Exs. No. 793; Hirn 1900, p. 102, Pl. VI, fig. 41; Skuja 1927, p. 98, Pl. II, fig. 11.

Monoecious; oogonium 1-4, obovoid-globose or subglobose, pore superior; oospore of same form as oogonium, which it nearly fills, spore wall smooth, often thick; suffultory cell occasionally larger than vegetative cell; antheridium 1-4, subepigynous, sometimes terminal; sperms 2, division horizontal; basal cell elongate; vegetative cell $12-22 \times 25-110\mu$; oogonium $38-55 \times 37-54\mu$; oospore $36-52 \times 35-51\mu$; antheridium $10-17 \times 8-13\mu$.

Finland, Germany, Sweden, Latvia.

Similar to *Oe. zigzag* (No. 22). Skuja (1927) reports *Oe. curtum* with 10-12 cells (and often 20) in a filament. *Oe.*

zigzag rarely has more than 6-8 cells in the filament. Skuja (1927) also records the membrane of the vegetative cell as being spirally punctate.

24. *Oe. fragile* Wittrock.

(Pl. XV, fig. 145.)

1870, p. 120; 1872a, p. 24; Wolle 1887, p. 71, Pl. 74, figs. 4-6; *Oe. Candollei* (Le Cl.) Breb. in Roumeguere Alg. Exs. No. 347; Hirn 1900, p. 96, Pl. V, figs. 33 and 34; Collins 1909, p. 232, fig. 81; Heering 1914, Pl. IV, fig. 36; P. B. A. No. 1477.

Monoecious; oogonium 1, globose or subobovoid-globose, with superior pore; oospore globose, filling oogonium, spore wall smooth; antheridium 1-3; sperms 2, division horizontal; basal cell elongate; vegetative cell $12-17 \times 50-120\mu$; oogonium $42-50 \times 44-55\mu$; oospore $39-46 \times 39-46\mu$; antheridium $12-15 \times 10-12\mu$.

United States: Iowa, Illinois, Mississippi, Massachusetts, Pennsylvania, Michigan; British Columbia; Patagonia; South Africa; Australia; France, Finland, Sweden, Latvia.

24a. Var. *abyssinicum* Hirn.

(Pl. XV, fig. 147.)

1900, p. 97, Pl. VI, fig. 35.

Oogonium smaller than the type; antheridium to 6-seriate; vegetative cell $12-17 \times 25-100\mu$; oogonium $38-45 \times 40-50\mu$; oospore $36-43 \times 36-44\mu$; antheridium $11-15 \times 6-14\mu$.

Africa.

24b. Var. *robustum* (West and West) Tiffany nov. comb.

(Pl. XV, fig. 146.)

Oe. zigzag Cleve var. *robustum* West and West 1903, p. 36; *ibid.* in West 1904, p. 60, fig. 12B; *ibid.* in Hirn 1906, p. 24, Pl. I, fig. 3.

Oogonium usually globose; antheridium 2; vegetative cell $19-23 \times 40-90\mu$; oogonium $53-58 \times 50-60\mu$; oospore $51-56 \times 47-53\mu$; antheridium $19 \times 5-7\mu$.

England.

Oe. fragile is characterized by its usually globose spores completely filling the oogonia, its size, and monoecious habit. Cf. *Oe. globosum* (No. 21) and *Oe. intermedium* (No. 19). The variety *abyssinicum* is more slender and has a larger number of seriate antheridia than the species. Variety *robustum* is considerably larger than either. The latter was originally described by W. and G. S. West as a variety of *Oe. zigzag* (No. 22). Its pluricellular habit and number of antheridia place it outside the few-celled *zigzag*. Its globose oospore completely filling the oogonium and its subepigynous and subhypogynous antheridia are characters that seem nearest to *Oe. fragile*.

25. *Oe. vaucherii* (Le Clerc) Al. Braun; Wittrock.

(Pl. XV, figs. 150 and 151.)

(?) *Prolifera vaucherii* Le Cl. 1817, p. 474, Pl. 23, fig. 4; (?) *Oe. vaucherii* Braun 1855, p. 40, Pl. 2, fig. 13; Wittr. 1870, p. 121; (?) *Oe. diandronites* Carter 1858, p. 31, Pl. 3, figs. 3, 9-11; (?) *Oe. monandronites* Carter 1858, p. 38; Hirn 1900, p. 97, Pl. VI, fig. 36; *Oe. vaucherii* (Le Cl.) Al. Br.; Wittr. f. *insulare* Hirn 1900, p. 99, Pl. VI, fig. 37; Collins 1909, p. 232; Heering 1914, p. 207, fig. 303; Tiffany 1926, p. 100, Pl. IV, fig. 35; P. B. A. No. 1786.

Monococious; oogonium 1, obovoid to subovoid-globose, pore superior; oospore globose to subglobose, not filling oogonium, spore wall smooth and sometimes thick; antheridium 1-4; sperms 2, division horizontal; basal cell elongate; vegetative cell $20-30 \times 32-118\mu$; oogonium $40-58 \times 45-65\mu$; oospore $35-54 \times 35-55\mu$; antheridium $17-30 \times 6-15\mu$.

United States: Massachusetts, Michigan, Iowa, Illinois, Ohio, Mississippi; India; Denmark, France, Germany, Switzerland, Italy, Sweden, Latvia.

Oe. vaucherii resembles *Oe. intermedium* (No. 19) and *Oe. fragile* (No. 24). Hirn's f. *insulare* is not distinct and is included with the species. The dimensions given above include the variations observed for the American material.

26. *Oe. richterianum* Lemmermann.

(Pl. XVI, fig. 156.)

1895, p. 26, figs. 1-3; *Oe. paludosum* (Hass.) Kuetz. in Hirn 1895, p. 14; Hirn 1900, p. 117, Pl. XII, fig. 63.

Monococious; oogonium 1-2, obovoid or subellipsoid, pore superior; oospore subobovoid or subellipsoid (rarely globose-ellipsoid), filling oogonium or not, spore wall smooth; antheridium 1-6, subhypogynous or subepigynous or scattered; sperms 2, division horizontal; basal cell elongate; vegetative cell $12-21 \times 36-126\mu$; oogonium $36-48 \times 48-74\mu$; oospore $35-43 \times 43-59\mu$; antheridium $12-15 \times 6-10\mu$.

United States: Massachusetts; British Columbia; Germany, Finland.

Distinguished by its combination of dimensions and by the horizontal division of the antheridium. Cf. *Oe. upsaliense* (No. 29).

27. *Oe. pseudoboscii* Hirn.

(Pl. XX, fig. 186.)

1900, p. 291, Pl. XIII, fig. 67; *Oe. neglectum* Hirn 1895, p. 21, Pl. 1, fig. 1; Silf-venius 1903, p. 15, fig.; Iwanoff 1901, p. 48; Hirn 1906, p. 48, Pl. II, fig. 6.

Monococious; oogonium 1, subobovoid, pore superior; oospore ellipsoid-ovoid or ellipsoid, inflating lower part of oogonium, otherwise not filling it, spore wall smooth; antheridium 1-2, subepigynous; sperms 2, division horizontal; vegetative cell $8-14 \times 64-275\mu$; oogonium $41-50 \times 75-105\mu$; oospore $38-45 \times 48-60\mu$; antheridium $11-12 \times 10-11\mu$.

United States: Illinois, Alabama, Massachusetts; Finland, Russia.

Iwanoff and Silfvenius (*loc. cit.*) both found material of this species which proved it to be monoecious and not dioecious (Hirn 1900, 1906). It is characterized by the relatively short oospore within a much longer oogonium. The diameter of the oospore is such that the lower part of the oogonium is really inflated. It bears some resemblance to *Oe. paludosum* (No. 53) and *Oe. boscii* (No. 55). From both it is distinguished by its smooth oospores. It is unlike *Oe. boscii* in its monoecious habit.

28. *Oe. sodiroanum* Lagerheim.

(Pl. XVI, figs. 153 and 154.)

1890, p. 81; Hirn 1900, p. 118.

Monoecious; oogonium 1, ellipsoid-obovoid, pore superior; oospore ellipsoid, not filling oogonium at upper end, spore wall smooth; antheridium subhypogynous or subepigynous or scattered, 1-9; sperms 2, division vertical; vegetative cell $20-24 \times 44-84\mu$; oogonium $40-45 \times 70-90\mu$; oospore $38-42 \times 56-64\mu$; antheridium $20-22 \times 6-8\mu$.

United States: Michigan; Ecuador.

This species originally described incompletely from Ecuador by Lagerheim was collected nearly forty years later in Michigan by Miss Alma B. Ackley. The above description is based on the Michigan material. There can be no question of the validity of the species. It resembles in general appearance *Oe. oviforme* (No. 30), *Oe. upsaliense* (No. 29), and *Oe. richterianum* (No. 26). It is readily distinguished from these (Cf. No. 29) in dimensions and habit. The antheridia occur in series and when present at the end of the filament give the plant a curved appearance, not unlike that of the androsporangial filaments of *Oe. spectabile* (No. 186).

29. *Oe. upsaliense* Wittrock.

(Pl. XVI, fig. 157.)

1870, p. 125; 1872a, p. 22; 1874, p. 14, Pl. I, fig. 4; *Oe. tumidulum* (Roth) in Areschoug Alg. Exs. No. 236, 1864; Hirn 1900, p. 115, Pl. XII, fig. 60; Wittr. and Nordst. Alg. Exsicc. No. 18; P. B. A. No. 1576.

Monoecious; oogonium 1, obovoid or suboblong-ellipsoid, pore superior; oospore same form as oogonium (which it fills), spore wall smooth; antheridium 1-3, subhypogynous; antheridium, oogonium, and vegetative cell alternating; sperms 2, division vertical; vegetative cell varying much in same filament; suffultory cell up to 27μ in diameter; basal cell elongate; terminal cell obtuse; vegetative cell $13-20 \times 55-160\mu$; oogonium $45-50 \times 66-100\mu$; oospore $42-47 \times 60-75\mu$; antheridium $15-18 \times 7-10\mu$.

United States: Michigan, New Hampshire, Ohio, Illinois; Greenland; France, Germany, Sweden, Latvia.

29a. Var. *fennicum* Hirn.

(Pl. XVI, fig. 158.)

1895, p. 14; 1900, p. 116, Pl. XII, fig. 61.

Oogonium and oospore larger, ellipsoid; vegetative cell $13-20 \times 55-175\mu$; suffultory cell $25-33 \times 40-66\mu$; oogonium $63-68 \times 75-85\mu$; oospore $61-64 \times 70-78\mu$; antheridium $15-20 \times 8-13\mu$

Finland.

This smooth-spored species is one of a few in the genus combining the monoecious habit with the tendency toward ellipsoid oogonia. It is distinguished from *Oe. oviforme* (No. 30) by its slightly smaller size, its varying diameter of vegetative cell, and by the rather characteristic alternation of antheridium, oogonium, and vegetative cell in the filament. *Oe. sodiroanum* (No. 28) is smaller and has shorter vegetative cells. *Oe. upsaliense* bears some resemblance to *Oe. richterianum* (No. 26) but in the latter the sperms result from a horizontal division of the antheridium, not a vertical division. The variety *fennicum* has much larger oogonia and antheridia. The smaller suffultory cell is usually characteristic of both the species and the variety.

30. *Oe. oviforme* (Lewin) Hirn.

(Pl. XVI, fig. 152.)

1900, p. 116, Pl. XII, fig. 62; *Oe. urbicum* Wittr. var. *oviforme* Lewin 1888, p. 17, Pl. 2, fig. 43; Heering 1914, p. 210, fig. 308; Tiffany 1926, p. 100, Pl. VIII, figs. 78 and 79.

Monoecious; oogonium 1, obovoid to ellipsoid-obovoid, pore superior; oospore obovoid to ovoid (rarely almost globose), filling or not filling the oogonium, spore wall smooth, thick; antheridium 1-4, epigynous, hypogynous, or scattered; sperms 2, division vertical; basal cell elongated; vegetative cell $15-23 \times 40-135\mu$; oogonium $48-55 \times (56-65-80\mu)$; oospore $46-53 \times 50-63\mu$; antheridium $16-19 \times 7-12\mu$.

United States: Iowa; Spain.

The material distributed from Spain shows a more uniformly obovoid oogonium than is common in the Iowa samples: the latter approach the ellipsoid-obovoid variation. Cf. *Oe. upsaliense* (No. 29).

31. *Oe. geniculatum* Hirn.

(Pl. XVI, fig. 155.)

1900, p. 106, Pl. VIII, fig. 48; P. B. A. No. 411.

Monoecious; oogonium 1, obovoid or obovoid-globose, pore superior; oospore globose or subdepressed-globose, not filling oogonium, spore wall smooth and thick; antheridium 1-5, subepigynous or subhypogynous or scattered, sometimes alternating with vegetative cells; sperms (?) 2, division (?) horizontal; vegetative cell $37-48 \times 60-135\mu$; oogonium $56-63 \times 56-68\mu$; oospore $48-59 \times 48-59\mu$; antheridium $37-44 \times 5-9\mu$.

United States: California.

This species is near *Oe. capilliforme* (No. 37) and *Oe. anomalum* (No. 40). It is larger than either and has somewhat larger oogonia in proportion to the size of the vegetative cells than is the case in *Oe. anomalum*. The peculiar curvature (from which the specific name is derived) of the filaments, due to the opening of the antheridia, is usually characteristic of the species.

32. *Oe. suboctangulare* West and West.

(Pl. XVII, fig. 160.)

1902, p. 129, Pl. 17, figs. 1 and 2; Hirn 1906, p. 23, Pl. I, fig. 4.

Monoecious; oogonium 1, longitudinally rectangular in optical section, broadly inflated, membrane $3.4-3.8\mu$ thick, yellow, pore superior; oospore subrectangular-ellipsoid, filling oogonium except at the corners; antheridium 2, scattered; vegetative cell $50-54 \times 175-240\mu$; oogonium $53-67 \times 82-92\mu$; oospore $50-60 \times 73-85\mu$; antheridium $41-48 \times 25-27\mu$.

Ceylon.

The rectangularly appearing oogonium slightly exceeding the vegetative cells in diameter is a distinctive characteristic.

33. *Oe. martinicense* Hirn.

(Pl. XVII, fig. 159.)

1900, p. 134, Pl. XVI, fig. 92; *Oe. crassum* (Hass.) Wittr. in Wolle 1887, p. 74, Pl. 76, figs. 2 and 3; Collins 1909, p. 240.

Monoecious; oogonium 1, obovoid or subobovoid, pore superior; oospore obovoid or obovoid-ellipsoid, quite filling oogonium, spore wall smooth, often thick; antheridium 1-5, hypogynous; sperms 2, division vertical; vegetative cell $33-37 \times 115-240\mu$; oogonium $(63-68-74 \times 96-125\mu)$; oospore $66-72 \times 81-96\mu$; antheridium $33-35 \times 5-7\mu$.

United States: Iowa, Kansas; Island of Martinique.

In appearance this species resembles *Oe. landsboroughi* (No. 48) or *Oe. crassum* (No. 49), both of which, however, are dioecious.

34. *Oe. kurzii* Zeller.

(Pl. XVII, fig. 163.)

1873, p. 189; 1873a, p. 189; Hirn 1900, p. 135, Pl. XVI, fig. 93.

Monoecious; oogonium 1, rarely in series, obovoid or subellipsoid, pore superior; oospore subglobose or ellipsoid, not filling oogonium, spore wall smooth; antheridium hypogynous or scattered, to 15-seriate; sperms 2, division vertical; terminal cell apically obtuse; vegetative cell $44-52 \times 90-250\mu$; oogonium $70-95 \times 111-130\mu$; oospore $67-86 \times 80-93\mu$; antheridium $44-52 \times 6-16\mu$.

United States: Pennsylvania, Arkansas, Mississippi; India.

This species is quite easily recognized by its monoecious habit, large size, and subglobose to ellipsoid oospores not filling the oogonia.

35. **Oe. capillare** (Linnaeus) Kuetzing.

(Pl. XVIII, figs. 164 and 165.)

Conferva capillate L. 1753, p. 1166; *Oe. capillare* Kuetz. 1843, p. 255, Pl. 12, fig. 11, 1-10; Kuetz. 1853, p. 13, Pl. 40, fig. 3; *Oe. capillare* Kuetz. var. *flavescens* in Rabenh. Alg. Europe 1861; *Oe. stagnale* Kuetz. in Rabenh. *ibid.* 1883; Hirn 1900, p. 112, Pl. XI, fig. 58; Heering 1914, p. 197, fig. 278; Tiffany 1926, p. 91, Pl. IV, figs. 39 and 40.

Diocious, macrandrous; oogonium 1, not or scarcely exceeding the vegetative cell in diameter, cylindrical to subcylindrical, pore superior; oospore globose to cylindric-globose to ovoid, not or completely filling oogonium, spore wall smooth; antheridium 1-4, often alternating with vegetative cell; sperms 2, division horizontal; basal cell elongate, terminal cell broadly apiculate to obtuse; female vegetative cell $35-56 \times 36-120\mu$, male $35-50 \times 35-90\mu$; oogonium $40-60 \times 45-75\mu$; oospore $30-52 \times 35-65\mu$; antheridium $30-48 \times 5-10\mu$.

United States: Iowa; Austria, Denmark, Finland, France, Germany, Spain, Italy, Russia, Sweden, Latvia; Africa (Victoria).

35a. Form **stagnale** (Kuetzing; Wittrock) Hirn.

(Pl. XVIII, fig. 166.)

1900, p. 113, Pl. X, fig. 57; *Oe. stagnale* Kuetz. 1849, p. 368, and 1853, p. 13, Pl. 41, fig. 2; Wittrock 1874, p. 31.

Oospore subcylindrical or cylindrical-globose, sometimes constricted at the middle, not filling oogonium; female vegetative cell $38-50 \times 40-100\mu$, male $35-45 \times 36-90\mu$; oogonium $40-60 \times 55-75\mu$; oospore $36-57 \times 40-60\mu$; antheridium $33-42 \times 5-9\mu$.

Germany, Latvia, Spain.

This species and its variety are rather readily distinguished from others by having oogonia which scarcely exceed the vegetative cells in diameter. They are to be compared with *Oe. suboctangulare* (No. 32) and *Oe. geniculatum* (No. 31), both of which are monoecious. The variety *stagnale* is separable from the type on dimensions and shape of oogonium.

36. **Oe. plagiostomum** Wittrock.

(Pl. XIV, fig. 140.)

Oe. sp. Wittr. 1872a, p. 24, Pl. 1, fig. 11; Wittr. 1874, p. 41; Hirn 1900, p. 100, Pl. VI, fig. 39; Collins 1918, p. 62; Heering 1914, p. 196, fig. 276; Tiffany 1926, p. 92.

Diocious, macrandrous; oogonium 1, obovoid-globose, with superior pore; oospore globose to subglobose, usually filling oogonium, spore wall smooth and thickened; antheridium 1-6, often alternating with vegetative cell; basal cell elongate; vegetative cell $22-27 \times 65-120\mu$; oogonium $42-49 \times 50-60\mu$; oospore $41-47 \times 42-49\mu$; antheridium $20-24 \times 8-10\mu$.

United States: Illinois, Massachusetts, Iowa, Ohio, Mississippi; Denmark, Sweden; India; South Africa.

36a. Var. **gracilius** Wittrock.

(Pl. XIV, figs. 141 and 142.)

1878, p. 142; Hirn 1900, p. 101, Pl. VI, fig. 40; Collins 1909, p. 233; Tiffany 1926, Pl. IV, figs. 37 and 38.

Smaller than the type, with vegetative cell usually shorter; vegetative cell $20-25 \times 40-100\mu$; oogonium $36-42 \times 46-57\mu$; oospore $34-39 \times 36-44\mu$; antheridium $19-22 \times 7-10\mu$.

United States: Iowa, New York, Illinois, Ohio, Mississippi; Germany; Mexico; South Africa.

The species and variety are separated mainly on the basis of size. The rather regularly globose oospores, with heavy walls, within obovoid-globose oogonia are characteristic.

37. **Oe. capilliforme** Kuetzing; Wittrock.

(Pl. XIX, figs. 172 and 173.)

Kuetz. 1853, p. 12, Pl. 37, fig. 3; Wittrock 1872a, p. 21; 1874, p. 31; *Oe. dioicum* Petrovsky 1861, p. 611, Pl. 13, fig. 6-10; Hirn 1900, p. 107, Pl. VIII, fig. 49; Collins 1909, p. 234; Heering 1914, p. 196, fig. 282; Tiffany 1926, p. 92, Pl. III, figs. 22 and 23.

Dioecious, macrandrous; oogonium 1, obovoid to subovoid, with superior pore; oospore variable, ovoid-globose, cylindric-globose, sub-globose, or globose, not filling oogonium, spore wall smooth; antheridium 2-8, often alternating with the vegetative cell; sperms 2, division horizontal; basal cell elongate, terminal cell apically obtuse or apiculate; female vegetative cell $28-38 \times 42-120\mu$, male $25-30 \times 40-100\mu$; oogonium $42-50 \times 51-62\mu$; oospore $37-45 \times 40-50\mu$; antheridium $20-25 \times 8-10\mu$.

United States: Missouri, Iowa, Ohio, Mississippi, Illinois; France, Russia, Sweden; Patagonia; S. Africa.

37a. Form **debaryanum** (Chmielevsky) Hirn.

(Pl. XIX, fig. 174.)

1900, p. 108, Pl. VIII, fig. 50; *Oe. De Baryanum* Chmiel. 1889, p. 1, Pl. 1, figs. 1 and 2; Tiffany 1926, p. 92, Pl. III, figs. 20 and 21.

Oospore globose or subglobose; female vegetative cell $25-37 \times 38-130\mu$, male $22-27 \times 35-100\mu$; oogonium $46-56 \times 52-66\mu$; oospore $38-50 \times 40-53\mu$; antheridium $18-24 \times 5-10\mu$.

United States: Iowa; Greenland; Russia.

37b. Form **lorentzii** (Magnus and Wille) Hirn.

(Pl. XIX, fig. 175.)

1900, p. 109, Pl. IX, fig. 51; *Oe. lorentzii* Magnus and Wille in Wille 1884, p. 51, Pl. 3, figs. 99-101.

Oogonium obovoid to obovoid-globose; oospore usually globose or subglobose, sometimes cylindric-globose; female vegetative cell $26-34 \times$

25–100 μ , male 24–30 $\times$ 25–90 μ ; oogonium 40–53 $\times$ 40–65 μ ; oospore 37–47 $\times$ 38–54 μ ; antheridium 22–30 $\times$ 5–10 μ .

Uruguay.

37c. Var. **australe** Wittrock.

(Pl. XIX, figs. 176 and 177.)

In Wittr. and Nordst. Exs. 1886; *Oe. princeps* (Hass.) Wittr. in Tilden Amer. Alg. 1894; Hirn 1900, p. 109, Pl. IX, fig. 52; *Oe. capilliforme* Kuetz.; Wittr. var. *australe* Wittr. f. *uberosporum* Hirn 1900, p. 110, Pl. IX, fig. 53; Tiffany 1926, p. 92.

Oogonium obovoid-globose to subglobose; female vegetative cell 24–36 $\times$ 26–100 μ , male 22–33 $\times$ 40–100 μ ; oogonium 37–53 $\times$ 43–60 μ ; oospore 35–50 $\times$ 36–55 μ ; antheridium 21–26 $\times$ 4–9 μ .

United States: Minnesota, Iowa, New York, South Dakota, Nebraska, California; Africa; Uruguay, Argentine, Brazil; Mexico; Vancouver.

37d. Var. **diversum** (Hirn) Tiffany.

(Pl. XIX, fig. 178.)

1929, p. 75; *Oe. capilliforme* Kuetz.; Wittr. var. *australe* Wittr. f. *diversum* Hirn 1900, p. 110, Pl. IX, fig. 54; *Oe. stagnale* Kuetz. in Tilden Amer. Alg. 1896.

Dimensions larger throughout; female vegetative cell 34–46 $\times$ 45–130 μ , male 33–40 $\times$ 50–120 μ ; oogonium 46–56 $\times$ 46–70 μ ; oospore 43–52 $\times$ 40–58 μ ; antheridium 30–37 $\times$ 6–11 μ .

United States: Iowa, Colorado, Montana, Minnesota.

The “capilliforme” group is readily recognized by the smaller male vegetative filaments and shape of oogonia. The group is variable within itself, however, and the above varieties are recognized on the basis of differences in shape of oogonia and in dimensions.

38. **Oe. princeps** (Hassall) Wittrock.

(Pl. XVIII, figs. 167–169.)

Vesiculifera princeps Hass. 1842, p. 388; *V. capillaris* Hass. 1845, p. 195, Pl. 50, figs. 1 and 2; *Oe. princeps* Wittr. 1874, p. 42; Hirn 1900, p. 289, Pl. X, fig. 56.

Dioecious, macrandrous; oogonium 1, little tumid, subobovoid, pore superior; oospore globose or subglobose, not filling oogonium, spore wall smooth; terminal cell apiculate; antheridium 1–7; vegetative cell 33–42 $\times$ 40–133 μ ; oogonium 54–63 $\times$ 67–80 μ ; oospore 48–58 $\times$ 50–60 μ ; antheridium 32–38 $\times$ 8–20 μ .

United States: Minnesota, Mississippi; England.

I have seen specimens of this alga from both Minnesota and Mississippi with immature antheridia which appear to belong with the oogonial filaments. I have therefore placed the species in the dioecious group. The cells of the antheridial filaments are somewhat longer than those of the male plants. It should be compared with the monoecious *Oe. geniculatum* (No. 31) and the dioecious *Oe. capilliforme* (No. 37). It is more robust than the latter.

39. *Oe. biforme* Nordstedt.

(Pl. XVII, figs. 161 and 162.)

In Wittrock and Nordstedt Exs. 1880; Hirn 1900, p. 147, Pl. XXII, fig. 112.

Dioecious, macrandrous; oogonium 1 (rarely 2), pore superior, cylindric-obovoid; oospore oblong-ellipsoid to subglobose or nearly globose, generally not filling oogonium, spore wall smooth; antheridium to 21-seriate; sperms 2, division vertical; female vegetative cell $22-32 \times 55-140\mu$; male $18-28 \times 45-140\mu$; oogonium $37-52 \times 48-90\mu$; oospore $36-48 \times 40-70\mu$; antheridium $18-28 \times 6-15\mu$.

Brazil, Paraguay, Ecuador.

Near *Oe. oboviforme* (No. 46) and *Oe. grande* (No. 45); distinguished chiefly by its cylindric-obovoid oogonium.

40. *Oe. anomalum* Hirn.

(Pl. XVIII, figs. 170 and 171.)

1900, p. 112, Pl. X, fig. 55; *Oe. stagnale* Kuetz. var. *variabilis* Lewin 1888, p. 17, Pl. 3, figs. 44-48; Heering 1914, p. 197, fig. 277; Tiffany 1926, p. 91, Pl. I, figs. 6 and 7.

Dioecious, macrandrous; oogonium 1, subovoid or cylindric-ovoid, pore superior; oospore globose or subglobose, not filling oogonium, spore wall smooth; antheridium 4-13, sperms 2, division vertical; terminal cell apically obtuse; female vegetative cell $40-50 \times 80-300\mu$, male $33-42 \times 80-240\mu$; oogonium $54-64 \times 68-75\mu$; oospore $48-60 \times 52-61\mu$; antheridium $30-40 \times 6-18\mu$.

United States: Iowa; Spain.

This species has some of the appearances of *Oe. capillare* (No. 35), particularly in the relatively small difference between the diameter of the vegetative cells and that of the oogonia.

41. *Oe. fabulosum* Hirn.

(Pl. XXV, figs. 215 and 216.)

1900, p. 114, Pl. XI, fig. 59.

Dioecious, macrandrous; oogonium 1, little tumid, cylindrical, pore superior; oospore globose or ellipsoid, not filling oogonium in length, spore wall smooth; antheridium seriate; sperms 2, division vertical; vegetative cell $74-85 \times 115-210\mu$; oogonium $81-96 \times 104-133\mu$; oospore $75-89 \times 78-104\mu$; antheridium $68-80 \times 7-22\mu$.

Brazil.

41a. Var. *maximum* (West) Hirn.

(Pl. XXV, figs. 217 and 218.)

1906, p. 13, Pl. II, fig. 5; *Oe. maximum* West 1901, p. 75, Pl. IV, figs. 39-41.

Oogonium wall thick; oospore subglobose to subcylindrical, filling oogonium or nearly so; basal cell elongate; terminal cell obtuse or broadly apiculate; vegetative cell $70-93 \times 105-335\mu$; oogonium $75-107 \times 75-136\mu$; oospore $70-100 \times 65-110\mu$; antheridium $65-86 \times 7-20\mu$.

Siam (Koh-chang).

41b. Var. **columbianum** G. S. West.

(Pl. XXV, fig. 219.)

1914, p. 1048, text fig. 3D.

Vegetative cell more slender; oogonium a little longer and more tumid; oospore more ellipsoid than the type; vegetative cell $56-70 \times 248-425\mu$; oogonium $98-100 \times 163-174\mu$; oospore $81-84 \times 124-128\mu$.

Columbia.

41c. Var. **punctatum** Lemmermanni.

1910, p. 313.

Oogonium not tumid; oospore globose, not filling oogonium in length, spore wall yellow-brown, densely punctate; vegetative cell $64-75 \times 210-245\mu$; oogonium $64-75 \times 101-113\mu$; oospore $63-71 \times 63-71\mu$.

Paraguay.

Oe. fabulosum with its three varieties represents the largest size attained by any species of the genus so far recorded. The variety *punctatum* is easily separable by its punctate oospore, although unfortunately Lemmermann left no figure; *columbianum* is the most slender; *maximum* is slightly larger than the species.

42. **Oe. rivulare** (Le Clerc) Al. Braun.

(Pl. XXIV, figs. 209 and 210.)

(?) *Prolifera rivularis* Le Cl. 1817, p. 472, Pl. 23, fig. 1; *Oe. rivulare* Al. Br. 1855, p. 23, Pl. 1, figs. 1-10; Hirn 1900, p. 119, Pl. XII, fig. 66.

Diocious, macrandrous; oogonium 1-7, obovoid, pore superior; oospore obovoid, ellipsoid, or subglobose, not filling oogonium longitudinally, spore wall smooth; antheridium to 13-seriate; sperms 2, division horizontal; basal cell elongate; female vegetative cell $35-45 \times 110-350\mu$, male $30-36 \times 120-280\mu$; oogonium $70-85 \times 130-160\mu$; oospore $55-70 \times 65-100\mu$; antheridium $21-28 \times 14-26\mu$.

United States: Iowa, Florida, Massachusetts; Denmark, England, Austria, France, Scotland, Germany, Sweden.

Very readily recognized by its large size and particularly by its oospore which is much smaller than the oogonium.

43. **Oe. pachyandrium** Wittrock.

(Pl. XX, figs. 179 and 180.)

In Wittr. and Nordst. Alg. Exs. No. 5, 1877; Hirn 1900, p. 142, Pl. XX, fig. 104.

Diocious, macrandrous; oogonium 1-3, subobovoid or obovoid-ellipsoid, pore superior; oospore ellipsoid, not filling oogonium (rarely inflating it), spore wall smooth; antheridium 1-4; sperms 2, division vertical; female vegetative cell $30-36 \times 80-220\mu$, male $32-45 \times 60-160\mu$; oogonium $53-57 \times 84-108\mu$; oospore $51-54 \times 69-85\mu$, antheridium $30-43 \times 11-20\mu$.

United States: Pennsylvania, New Jersey, Massachusetts; Sweden.

This plant is very near *Oe. grande* (No. 45), but is distinguished by having male filaments larger than the female.

44. *Oe. mexicanum* Wittrock.

(Pl. XXI, figs. 188 and 189.)

1878, p. 138; Hirn 1900, p. 147, Pl. XXII, fig. 111.

Dioecious, macrandrous; oogonium 1, cylindric-obovoid, pore superior; oospore cylindric-obovoid, quite or nearly filling oogonium; spore wall smooth; antheridium to 8-seriate; sperms 2, division vertical; female vegetative cell $34-41 \times 60-140\mu$, male $30-38 \times 60-114\mu$; oogonium $53-63 \times 76-110\mu$; oospore $51-60 \times 63-80\mu$; antheridium $28-35 \times 7-17\mu$.

United States: South Dakota, Massachusetts; Mexico.

This species is close to *Oe. grande* below, being usually distinguished by its larger and considerably shorter vegetative cells and by its more cylindric oogonia.

45. *Oe. grande* Kuetzing; Wittrock.

(Pl. XXII, figs. 195 and 196.)

In Wittr. and Nordst. Alg. Exs. No. 502, 1883; Kuetz. 1845, p. 200, 1853, p. 12, Pl. 37, fig. 1; Hirn 1900, p. 143, Pl. XXI, figs. 105-106; Collins 1909, p. 242; Heering 1914, p. 202, fig. 288; Tiffany 1926, p. 93, Pl. II, figs. 14-16; P. B. A. No. 811.

Dioecious, macrandrous; oogonium 1-5, subobovoid, pore superior; oospore of the same form as oogonium, which it completely fills or not, spore wall smooth; antheridium 1-10; sperms 2, division vertical; basal cell elongated; female vegetative cell $28-37 \times 70-210\mu$, male $28-33 \times 70-175\mu$; oogonium $49-60 \times 86-110\mu$; oospore $47-58 \times 60-94\mu$; antheridium $25-33 \times 11-18\mu$.

United States: Iowa, Maine, Massachusetts, Pennsylvania, California, Michigan, Illinois, Ohio, Minnesota, Mississippi, Connecticut; Ontario; Australia; Austria, Sweden, Switzerland.

45a. Form *gemelliparum* (Pringsheim) Hirn.

(Pl. XXII, fig. 200.)

1900, p. 145, Pl. XXI, fig. 107; *Oe. gemelliparum* Pringsh. 1858, p. 71, Pl. 5, fig. 10; *Oe. landsboroughi* (Hass.) Wittr. var. *gemelliparum* (Pringsh.) Wittr. 1874, p. 36.

Oogonium ellipsoid-ovoid, antheridium seriate; vegetative cell $20-27 \times 60-200\mu$; oogonium $55-57 \times 75-80\mu$; oospore $49-51 \times 65-69\mu$.

Germany.

45b. Var. *aequatoriale* Wittrock.

(Pl. XXII, fig. 201.)

In Wittr. and Nordst. Alg. Exs. No. 1016, 1893; Hirn 1900, p. 145, Pl. XXI, fig. 108; *Oe. grande* Kuetz.; Wittr. var. *aequatoriale* Wittr. f. *hortense* Wittr. in Hirn 1900, p. 145, Pl. XXI, fig. 109.

Vegetative cell $26-33 \times 70-165\mu$; oogonium $44-51 \times 75-100\mu$; oospore $42-49 \times 50-81\mu$.

Ecuador; Greenland.

45c. Var. **angustum** Hirn.

(Pl. XXII, figs. 198 and 199.)

1900, p. 146, Pl. XXI, fig. 110; P. B. A. No. 410.

Vegetative cell and oogonium elongated; antheridium to 36-seriate; oogonia sometimes 4; vegetative cell, female $19-30 \times 70-330\mu$, male $19-25 \times 80-225\mu$; oogonium $42-52 \times 62-110\mu$; oospore $40-50 \times 60-89\mu$; antheridium $18-22 \times 7-15\mu$.

United States: Massachusetts, Mississippi, Michigan, Illinois; Brazil.

45d. Var. **majus** Hansgirg.

(Pl. XXII, fig. 197.)

1886, p. 45; *Oe. grande* forma *robusta* Hirn 1900, p. 144; Collins 1918, p. 63; *Oe. grande* Kuetz.; Wittr. var. *robustum* (Hirn) Tiffany 1924, p. 186, Pl. III, fig. 7, and 1926, p. 94, Pl. II, fig. 19.

Larger than the type; female vegetative cell $36-46 \times 80-200\mu$, male $32-42 \times 80-200\mu$; oogonium $52-68 \times 75-90\mu$; oospore $50-64 \times 68-88\mu$; antheridium $30-36 \times 10-16\mu$.

United States: Illinois, Iowa, Michigan, Ohio, Massachusetts; Austria, Czecko Slovakia, Latvia.

Oe. grande is one of our commonest species. The sub-obovoid oogonia often in series, with superior pore, the accompanying smaller male filaments with vertical division of the antheridium are distinctive characters. The varieties are distinguished on the basis of size and differences in shape of oogonia.

The common occurrence of the distinct, robust variety of *Oe. grande* led to the varietal name *robustum* (Tiffany 1924). Hansgirg's description of the variety *majus* is so near that given above that the prior name must be used, in spite of its relegation by Hirn (1900) to be synonymous with the robust form of the type. Skuja (1927) records a robust form of *Oe. grande* with slightly smaller dimensions, which I have included under the variety *majus*. The dimensions of this variety are from the American material.

46. **Oe. oboviforme** Wittrock.

(Pl. XX, figs. 184 and 185.)

1878, p. 140; De Toni 1889, p. 74; Hirn 1900, p. 141, Pl. XX, fig. 103; Collins 1909, p. 241.

Dioecious, macrandrous; oogonium 1, obovoid, pore superior; oospore obovoid (rarely ellipsoid-obovoid), about filling oogonium, spore wall smooth; antheridium to 19-seriate, sperms 2, division vertical; basal cell elongate; female vegetative cell $21-33 \times 72-280\mu$; male $21-31 \times 72-270\mu$; oogonium $55-65 \times 80-107\mu$; oospore $54-61 \times 70-85\mu$; antheridium $21-28 \times 5-13\mu$.

Brazil; Mexico.

Distinguished from *Oe. mexicanum* (No. 44) by longer vegetative cells. Its obovoid and slightly larger oogonia usually separate it from *Oe. grande*, above.

47. *Oe. subrectum* Hirn.

(Pl. XX, figs. 181-183.)

1900, p. 141, Pl. XX, fig. 102.

Dioecious, macrandrous; oogonium 1-2, subellipsoid or obovoid-ellipsoid, pore superior; oospore ellipsoid or globose-ellipsoid, filling or nearly filling oogonium, spore wall smooth, often thick; antheridium to 9-seriate, sperms 2, division vertical; basal cell elongate; terminal cell apically obtuse; vegetative cell, female, $27-35 \times 160-340\mu$, male $26-33 \times 104-300\mu$; oogonium $70-83 \times 83-103\mu$; oospore $65-80 \times 75-93\mu$; antheridium $22-28 \times 8-14\mu$.

Brazil.

Characterized among the species with obovoid-ellipsoid oogonia by its size and the relatively long vegetative cells.

48. *Oe. landsboroughi* (Hassall) Wittrock.

(Pl. XXIV, figs. 211 and 212.)

Vesiculifera landsboroughi Hass. 1842, p. 289, and 1845, p. 197, Pl. 51, fig. 2; *Oe. landsboroughi* Wittr. 1874, p. 35; *Oe. gemelliparum* Pringsh. in Rabenh. Alg. Eur. No. 1118, 1861; *Oe. gemelliparum* v. *majus* Wittr. 1870, p. 137; *Oe. tumidulum* Kuetz. in *ibid* No. 1468, 1863; Hirn 1900, p. 135, Pls. XVI and XVII, figs. 94 and 95; Collins 1909, p. 241; Heering 1914, p. 200, fig. 285; Tiffany 1926, p. 93, Pl. II, figs. 12 and 13; P. B. A. No. 663.

Dioecious, macrandrous; oogonium 1-2, (rarely 3), obovoid to ovoid, pore superior; oospore ovoid to ellipsoid, filling or not filling the oogonium, spore wall smooth; antheridium up to 30-seriate; sperms two, division vertical; basal cell elongate, terminal cell obtuse; female vegetative cell $31-40 \times 90-240\mu$, male $30-37 \times 120-225\mu$; oogonium $63-78 \times 85-115\mu$; oospore (55-) $59-70 \times 73-102\mu$; antheridium $27-35 \times 9-20\mu$.

United States: Iowa, Ohio, Connecticut, Michigan, Massachusetts, New York, Mississippi; British Columbia; Paraguay, Uruguay; Mexico; Austria, England, Finland, France, Germany, Norway, Russia, Sweden.

48a. Var. *norvegicum* Wittrock.

(Pl. XXIV, figs. 213 and 214.)

In Wittr. and Nordst. Alg. Exs. No. 204, 1879; *Oe. landsboroughi* var. *robustum* Wittr. in Wittr., Nordst. and Lagerh. Alg. Exs. No. 1215, 1896; Hirn 1900, p. 137, Pl. XVII, figs. 97 and 98; Tiffany 1926, p. 93.

Vegetative cells shorter but of larger diameter; female vegetative cell $38-45 \times 60-160\mu$, male $36-45 \times 55-135\mu$; oogonium $64-73 \times 70-105\mu$; oospore $60-70 \times 67-90\mu$; antheridium $35-45 \times 8-15\mu$.

United States: Iowa; Norway; Brazil; Mexico.

It is rather difficult to indicate the exact differences between the *landsboroughi* group and the *crassum* group in the genus,

although one can scarcely confuse the two after some familiarity with them. The shape of the oogonia is quite similar in each, but the oospore is perhaps the most distinctive characteristic. The oospore of *landsboroughi* varies from ovoid to ellipsoid while the oospore of *crassum* varies from ellipsoid to nearly globose. Globose oospores are never present in *Oe. landsboroughi*. The variety *norvegicum* has stouter and shorter vegetative cells. Compare also with *Oe. subrectum* (No. 47) and *Oe. grande* (No. 45).

49. *Oe. crassum* (Hassall) Wittrock.

(Pl. XXIII, figs. 202 and 203.)

(?) *Vesiculifera crassa* Hass. 1842, p. 389; *Oe. crassum* Wittr. 1872a, p. 20, Pl. 1, figs. 4-6; Hirn 1900, p. 139, Pl. XVIII, fig. 99; Heering 1914, p. 200, fig. 286; Tiffany 1926, p. 93, Pl. II, figs. 17 and 18.

Dioecious, macrandrous, oogonium 1-2, ovoid to obovoid-ellipsoid, pore superior; oospore ellipsoid to globose, filling or not filling oogonium, spore walls smooth; antheridium 2-25; sperms 2, division vertical; female vegetative cell $36-50 \times 72-340\mu$, male $30-36 \times 72-260\mu$; oogonium $60-78 \times 85-120\mu$; oospore $58-76 \times 75-96\mu$; antheridium $28-32 \times 10-20\mu$.

United States: Iowa, Florida, Ohio, Michigan, Mississippi, Alabama, Indiana; Austria, England, Germany, Sweden; South Africa.

49a. Form *amplum* (Magnus and Wille) Hirn.

(Pl. XXIII, figs. 204 and 205.)

1900, p. 139, Pls. XVIII and XIX, fig. 100; *Oe. amplum* Magnus and Wille in Wille 1884, p. 40, Pl. 2, figs. 65 and 66; *Oe. crassum* (Hass.) Wittr. in Wille 1884, p. 39, Pl. 2, fig. 64; (?) *Oe. rivulare* (Le Cl.) Al. Br. var. *major* Wolle 1887, p. 92, Pl. 76, fig. 7 and 8.

Of larger dimensions; vegetative cell, female $46-54 \times 70-160\mu$, male $42-50 \times 65-200\mu$; oogonium $75-90 \times 83-115\mu$; oospore $72-85 \times 77-100\mu$; antheridium $40-50 \times 8-20\mu$.

United States: Ohio, Illinois, Mississippi, (?) Florida, Michigan; Uruguay, Brazil.

49b. Var. *longum* Transeau.

(Pl. XXIV, fig. 208.)

1918, p. 237.

Elongated oogonium and oospore; vegetative cell, female $40-52 \times 100-240\mu$, male $36-44 \times 100-180\mu$; oogonium $68-84 \times 120-180\mu$; oospore $66-80 \times 100-125\mu$; antheridium $34-40 \times 8-14\mu$.

United States: New York (Short Point Bay, Oneida Lake).

49c. Var. *subtumidum* Hirn.

(Pl. XXIII, figs. 206 and 207.)

1900, p. 140, Pl. XIX, fig. 101.

Longer vegetative cells; oogonium and oospore larger than in type; vegetative cell, female $42-50 \times 120-250\mu$, male $40-48 \times 120-240\mu$; oogonium $75-85 \times 96-137\mu$; oospore $67-79 \times 84-100\mu$; antheridium $34-45 \times 7-15\mu$.

Brazil; South Africa.

For a discussion of the chief characteristics of the *crassum* group, see notes under *Oe. landsboroughi* (No. 48). The variety *amplum* is larger than the species proper and has shorter vegetative cells than the variety *subtumidum*. Fritsch's form of *crassum* from South Africa is nearest the variety *subtumidum*. The variety *longum* is readily characterized by its elongated oogonia and oospores.

50. *Oe. exocostatum* Tiffany.

(Pl. XXVII, figs. 235-237.)

1921, p. 272, Pl. I, figs. A-F, 1926, p. 94, Pl. V, figs. 51-54.

Dioecious, macrandrous, oogonium 1-2, ellipsoid to ellipsoid-globose, occasionally terminal, pore superior; oospore of the same form as the oogonium, which it very nearly completely fills, spore wall of two layers: the outer marked by 13-15 longitudinal ribs, inner smooth; suffultory cell swollen; male filament a little more slender than the female, antheridium 3-7; sperms 2, division horizontal; basal cell elongate; female vegetative cell (13-) $18-25 \times 72-140\mu$, male (13-) $16-20 \times 48-100\mu$; suffultory cell $22-30 \times 60-90\mu$; oogonium $40-52 \times 60-96\mu$; oospore $38-41 \times 56-68\mu$; antheridium $12-16 \times 7-12\mu$.

United States: Iowa, Ohio.

This species is readily distinguished by the presence of longitudinal ribs on the outer layer of the oospore wall. Otherwise it is similar in appearance to *Oe. crenulatocostatum* (No. 57).

51. *Oe. kjellmanii* Wittrock; Hirn.

(Pl. XXVI, fig. 223.)

1900, p. 127, Pl. XIV, fig. 82; Wittr. 1880, in Wittr. and Nordst. Alg. Exs. No. 306.

Dioecious, macrandrous; oogonium 1, obovoid or rarely subellipsoid, pore superior; oospore ellipsoid or subellipsoid, not completely filling oogonium, spore wall of 3 layers: outer layer internally ribbed, middle layer with 35-45 slightly crenulate longitudinal ribs, anastomosate, inner layer smooth; antheridium to 30-seriate; sperms 2, division vertical; female vegetative cell $15-22 \times 45-120\mu$; male $14-18 \times 56-120\mu$; oogonium $41-50 \times 60-75\mu$; oospore $39-47 \times 48-57\mu$; antheridium $12-15 \times 4-12\mu$.

Borneo.

Characterized by the large number of longitudinal ribs on the middle layer of the oospore.

52. *Oe. paulense* Nordstedt and Hirn.

(Pl. XXVI, fig. 224.)

In Hirn 1900, p. 292, Pl. XIV, fig. 80; Schmidle 1901, p. 344, Pl. XII, fig. 1; Hirn 1906, p. 45, Pl. II, fig. 5.

Monoecious; oogonium 1, subellipsoid or ellipsoid-ovoid, pore superior; oospore ellipsoid, not or quite filling oogonium, spore wall of three layers: outer layer smooth, middle layer with 18–22 longitudinal ribs, entire and sometimes anastomosate, inner layer smooth; antheridium 1–2, subepigynous; sperms 2, division horizontal; vegetative cell $10-15 \times 50-120\mu$; oogonium $33-38 \times 43-68\mu$; oospore $30-36 \times 38-45\mu$; antheridium $10-14 \times 6-8\mu$.

Brazil; Australia.

Schmidle's record of this species from South America definitely placed it in the monoecious group. Its small size and fewer ribs on the oospore are characteristic.

53. *Oe. paludosum* (Hassall) Wittrock.

(Pl. XXVI, fig. 231.)

Oe. paludosum Wittr. 1870, p. 124; (?) *Vesiculifera paludosa* Hass. 1845, p. 199, Pl. 52, fig. 3; Hirn 1900, p. 120, Pl. XIII, fig. 69; Collins 1909, p. 237.

Monoecious; oogonium 1, ellipsoid, pore superior; oospore ellipsoid, filling oogonium, spore wall in three layers: outer and middle layers with 27–35 longitudinal ribs, continuous, rarely anastomosate; antheridium 1–8, scattered, often in the upper part of the filament; sperms 2, division vertical; vegetative cell $15-20 \times 50-140\mu$; oogonium $39-48 \times 66-84\mu$; oospore $36-45 \times 54-63\mu$; antheridium $14-16 \times 6-13\mu$.

United States: Illinois, Pennsylvania; British Columbia, Norway, England, Sweden.

53a. Var. *americanum* Nordstedt.

(Pl. XXVII, fig. 233.)

In Hirn 1900, p. 121, Pl. XIII, fig. 72; Collins 1909, p. 237.

Oogonium larger; vegetative cell $14-23 \times 45-156\mu$; oogonium $54-63 \times 75-90\mu$; oospore $49-57 \times 69-75\mu$; antheridium $12-18 \times 7-12\mu$.

United States: South Carolina.

53b. Var. *parvisporum* Hirn.

(Pl. XXVI, fig. 232.)

1900, p. 120, Pl. XIII, fig. 70; (?) *Oe. pringsheimiana* Archer 1868, p. 295; (?) *Oe. archerianum* Cooke 1884, p. 157; Borge 1913, p. 63, fig. 1.

Oogonium subellipsoid or suboblong-ellipsoid; oospore not filling oogonium; antheridium subepigynous or subhypogynous or scattered; vegetative cell $15-20 \times 45-120\mu$; oogonium $38-44 \times 70-86\mu$; oospore $35-41 \times 54-60\mu$; antheridium $15-17 \times 7-12\mu$.

United States: Illinois, Michigan, Ohio; Spain, Sweden.

Oe. paludosum is to be compared with *Oe. boscii* (No. 55) from which it differs in being monoecious. The two varieties are distinguished chiefly by relative sizes and by differently shaped and incompletely filled oogonia.

54. *Oe. leiopleurum* Nordstedt and Hirn.

(Pl. XXVI, figs. 221 and 222.)

In Hirn 1900, p. 126, Pl. XIV, fig. 79; *Oe. kjellmanii* Wittr. in Nordst. 1888a, p. 196.

Dioecious, macrandrous; oogonium 1-4, obovoid (or ellipsoid-obovoid) or subellipsoid, pore superior; oospore obovoid-ellipsoid or subellipsoid, usually filling oogonium, spore wall in three layers: outer and middle layers with 26-30 longitudinal ribs, entire, sometimes anastomosate, inner layer smooth; antheridium to 24-seriate, sperms 2, division horizontal; basal cell elongate; terminal cell apically obtuse; female vegetative cell 15-25×45-170 μ , male 15-17×45-85 μ ; oogonium 40-51×65-83 μ ; oospore 39-49×50-68 μ ; antheridium 15-17×7-15 μ .

Africa.

To be compared with *Oe. boscii* below. Its large number of seriate antheridia and usually completely filled oogonia are characteristic.

55. *Oe. boscii* (Le Clerc) Wittrock.

(Pl. XXVI, figs. 225 and 226.)

1874, p. 34; (?) *Prolifera Boscii* Le Clerc 1817, p. 474, Pl. 23, fig. 5; *Conferva vesicata* Ag. in Desmaz. Exs. 1845; Hirn 1900, p. 122, Pl. XIII, fig. 73; Wittr., Nordst. and Lagerh. Alg. Ex. No. 1213.

Dioecious, macrandrous; oogonium 1 (rarely 2), oblong-ellipsoid, pore superior; oospore ellipsoid, not nearly filling oogonium longitudinally, spore wall of three layers: outer and middle layers with 27-35 continuous, rarely anastomosate, longitudinal ribs; antheridium 1-8, scattered, often in upper part of filament; sperms 2, division vertical; vegetative cell, female 14-23×45-135 μ , male 13-18×52-108 μ ; oogonium 39-51×75-110 μ ; oospore 36-43×56-70 μ ; antheridium 13-14×6-16 μ .

United States: Connecticut, Massachusetts, Iowa, California, Illinois, Ohio, Mississippi; Austria, England, Finland, France, Germany, Switzerland, Norway, Latvia, Sweden; Brazil; Greenland; British Columbia.

55a. Form *dispar* Hirn.

(Pl. XXVI, fig. 230.)

1900, p. 124, Pl. XIV, fig. 76; P. B. A. No. 1226.

Oospore smaller, ellipsoid-globose or subglobose; vegetative cell 14-21×60-170 μ ; oogonium 38-45×70-92 μ ; oospore 34-43×44-55 μ .

United States: Massachusetts, Pennsylvania, California.

55b. *Var. notabile* Lemmermann.

(Pl. XXVI, fig. 229.)

1898, p. 511, Pl. 5, figs. 21-23; Hirn 1900, p. 125, Pl. XIV, fig. 78.

Oogonium suboblong-obovoid; oospore obovoid or ellipsoid-obovoid; vegetative cell $15-20 \times 60-150\mu$; oogonium $45-50 \times 90-105\mu$; oospore $44-49 \times 75-85\mu$.

Germany.

55c. *Var. occidentale* Hirn.

(Pl. XXVI, figs. 227 and 228.)

1900, p. 125, Pl. XIV, fig. 77, P. B. A. No. 1671.

Smaller; vegetative cell $8-15 \times 50-165\mu$; oogonium $33-38 \times 75-100\mu$; oospore $32-37 \times 45-50\mu$; antheridium $12-13 \times 10-16\mu$.

United States: Maine.

To be compared with *Oe. paludosum* (No. 53) and *Oe. leiopleurum* (No. 54).

56. *Oe. margaritiferum* Nordstedt and Hirn.

(Pl. XXVII, fig. 234.)

In Hirn 1900, p. 128, Pl. XV, fig. 83.

Dioecious, macrandrous; oogonium 1, suboblong-ellipsoid or sub-obovoid-ellipsoid or subellipsoid, pore superior; oospore ellipsoid or globose-ellipsoid, usually not filling oogonium longitudinally (rarely quite filling it), spore wall of three layers: outer and middle layer with 30-35 granulate, anastomosate, longitudinal ribs, inner layer smooth; antheridium to 10-seriate; sperms 2, division vertical; female vegetative cell $23-28 \times 82-190\mu$, male $17-23 \times 70-160\mu$; oogonium $50-63 \times 82-100\mu$; oospore $48-61 \times 55-75\mu$; antheridium $18-20 \times 7-13\mu$.

Brazil.

The granulate ribs of the oospore are distinctive. Compare with *Oe. kjellmanii* (No. 51).

57. *Oe. crenulato-costatum* Wittrock.

(Pl. XXVIII, figs. 240 and 241.)

1878, p. 139; (?) *Oe. apiculatum* Wille 1877, p. 188; Hirn 1900, p. 129, Pl. XV, fig. 84; Collins 1909, p. 239; Heering 1914, p. 198, fig. 280; Tiffany 1926, p. 94, Pl. V, figs. 48-50.

Dioecious, macrandrous; oogonium 1-6, obovoid to subellipsoid, often terminal, pore superior; oospore of the same form as the oogonium, which it nearly or quite fills, outer spore wall smooth, median wall with 14-20 longitudinal ribs, crenulate and sometimes anastomosing, inner wall smooth; antheridium 2-6, often alternating with vegetative cell; sperms 2, division horizontal; terminal cell obtuse to broadly apiculate; female vegetative cell $10-18 \times 25-125\mu$, male $9-13 \times 32-80\mu$; oogonium $30-36 \times 40-65\mu$; oospores $28-34 \times 37-55\mu$; antheridium $9-12 \times 9-14\mu$.

United States: Pennsylvania, Iowa, New York, Massachusetts, Ohio, Illinois, Connecticut, New Hampshire, Michigan.

57a. Form *cylindricum* Hirn.

(Pl. XXVIII, fig. 244.)

1900, p. 129, Pl. XV, fig. 85; *Oe. boscii* (Le Cl.) Wittr. in P. B. A. No. 118, 1895.

Oogonium and oospore cylindric-oblong or more rarely ellipsoid or obovoid-ellipsoid; ribs of oospore scarcely crenulate; vegetative cell $11-16 \times 44-150\mu$; oogonium $30-36 \times 42-81\mu$; oospore $27-34 \times 40-65\mu$.

United States: Massachusetts, Connecticut, Missouri, Illinois, New Hampshire.

57b. Var. *aureum* Tilden.

(Pl. XXVIII, fig. 245.)

In Amer. Algae, Nos. 123 and 186, 1896; 1898, p. 90, Pl. 8, figs. 1-3; Hirn 1900, Pl. XV, p. 130, fig. 87.

Oospore obovoid to globose-ellipsoid, not completely filling oogonium, middle spore wall with dentate ribs (sometimes nearly areolate; vegetative cell $10-13 \times 35-110\mu$; oogonium $30-35 \times 38-50\mu$; oospore $29-33 \times 35-43\mu$; antheridium $10 \times 10\mu$.

United States: Colorado.

57c. Var. *longiarticulatum* Hansgirg.

(Pl. XXVIII, figs. 242 and 243.)

1888, p. 46; *Oe. calosporum* Hirn 1895, p. 20, Pl. 1, fig. 3; Hirn 1900, p. 130, Pl. XV, fig. 86.

Oogonium single; oospore obovoid to nearly ellipsoid, not filling oogonium, ribs of median layer of spore wall distinctly crenate; vegetative cell $12-15 \times 60-90\mu$; oogonium $27-30 \times 58-60\mu$; oospore $24-27 \times 44-52\mu$.

United States: Michigan, Massachusetts; Austria, Finland; Brazil.

There exists considerable variation in the members of the *crenulatocostatum* group. The crenate or crenulate ribs (sometimes nearly smooth), the usually seriate oogonia, and small size are characteristic. Cf. *Oe. exocostatum* (No. 50). The variety *aureum* has dentate ribs which sometimes appear nearly areolate. It is felt, after examining hundreds of specimens of this group, that the above form and varieties are usually sufficiently distinct to be recognized. Ecological variants are common and to give varietal distinction to each of these is merely to make a difficult situation worse. Plants of *longiarticulatum* from Brazil are slightly smaller in all dimensions than those given above (Borge).

58. *Oe. arcysporum* Nordstedt and Hirn.

(Pl. XXVIII, fig. 247.)

In Hirn 1900, p. 104, Pl. VII, fig. 44.

Monoecious; oogonium 1-5, obovoid to ellipsoid-globose, pore superior; oospore ellipsoid-globose to subglobose, not usually filling oogonium, spore wall of three layers: outer smooth, median areolate, inner smooth; antheridium 1-4, subepigynous to subhypogynous; vegetative cell $13-20 \times 50-200\mu$; oogonium $41-55 \times 43-68\mu$; oospore $38-50 \times 40-55\mu$; antheridium $11-15 \times 8-12\mu$.

Brazil.

The only monoecious form with an areolate oospore wall.

59. *Oe. areolatum* Lagerheim.

(Pl. XXVIII, figs. 248-250.)

1890, p. 80 and 89; Hirn 1900, p. 105, Pl. VII, fig. 45; Tiffany 1926, p. 91.

Dioecious, macrandrous; oogonium 1-4, obovoid or obovoid-globose, with superior pore; oospore subellipsoid or ellipsoid-globose, filling or not filling oogonium, middle layer of spore wall areolate; antheridium 3-12, sperms 2; basal cell elongate; terminal cell apically obtuse; female vegetative cell $16-21 \times 65-165\mu$, male $15-19 \times 60-140\mu$; oogonium $48-60 \times 60-75\mu$; oospore $45-57 \times 48-60\mu$; antheridium $14-17 \times 8-10\mu$.

United States: Iowa, Ohio; Ecuador, Brazil.

This species is to be compared with *Oe. hoehnei* (No. 202). It is one of the few dioecious macrandrous forms with areolate oospores.

60. *Oe. dictyosporum* Wittrock.

(Pl. XXVIII, fig. 246.)

1874, p. 13; Hirn 1900, p. 103, Pl. VII, fig. 43; Tiffany 1926, p. 99.

Monoecious; oogonium 1-2, obovoid-globose, with superior pore; oospore globose to ellipsoid-globose, usually not filling oogonium, outer layer of spore wall reticulate, inner smooth; antheridium 1-3, subepigynous; sperms 2; vegetative cell $11-16 \times 25-95\mu$; oogonium $33-40 \times 38-46\mu$; oospore $28-38 \times 30-40\mu$; antheridium $8-13 \times 5-10\mu$.

United States: Iowa; Brazil; Africa.

60a. Form *westii* Tiffany.1929, p. 74; *Oe. dictyosporum* Wittr. forma West 1907, p. 98.

Oogonium and oospore ovoid-ellipsoid; a little larger; vegetative cell $14-16 \times 50-80\mu$; oogonium $42 \times 52\mu$; oospore $40 \times 50\mu$; antheridium $13 \times 8-9\mu$.

Africa.

The only monoecious representatives of the genus with reticulate oospores are *Oe. dictyosporum* and its f. *westii*. The latter is a little larger than the type and is better recognized as a form until we have more data on its distinctive characteristics.

61. *Oe. foveolatum* Wittrock.

(Pl. XXIX, fig. 252.)

1878, p. 133; Hirn 1900, p. 106, Pl. VII, fig. 46.

Monoecious; oogonium 1-2, obovoid to subellipsoid-globose, with superior pore; oospore globose to subellipsoid-globose, filling or not filling oogonium, outer spore wall scrobiculate; antheridium 1-7; sperms two, division horizontal; basal cell elongate, terminal cell apically obtuse; vegetative cell $14-23 \times 35-115\mu$; oogonium $37-49 \times 38-57\mu$; oospore $33-46 \times 34-48\mu$; antheridium $15-19 \times 8-12\mu$.

Brazil; St. Thomas Island.

The only monoecious representative of the genus with scrobiculate oospores.

62. *Oe. scrobiculatum* Wittrock.

(Pl. XXIX, figs. 258 and 259.)

In Wittr. and Nordst. Alg. Exs. No. 1018, 1893; Hirn 1900, p. 133, Pl. XV, fig. 90.

Dioecious, macrandrous; oogonium 1-3, obovoid or subellipsoid, pore superior; oospore of same form as oogonium and nearly filling it, outer spore wall scrobiculate, inner smooth; antheridium 1-?, sperms 2, division horizontal; female vegetative cell $16-24 \times 50-144\mu$, male $15-19 \times 45-110\mu$; suffultory cell $21-30 \times 34-90\mu$; oogonium $40-48 \times 60-88\mu$; oospore $39-45 \times 48-57\mu$; antheridium $13-15 \times 8-12\mu$.

Ecuador.

Cf. *Oe. verrucosum* (No. 63) and *Oe. tiffanii* (No. 64).

63. *Oe. verrucosum* Hallas.

(Pl. XXXI, figs. 272-274.)

1905, p. 408, fig. 17; Hirn 1906, p. 24, Pl. II, fig. 8.

Dioecious, macrandrous; oogonium 1-2, broadly ellipsoid or obovoid-globose, pore superior; oospore globose or ellipsoid-globose, not filling oogonium, middle layer of spore wall scrobiculate; suffultory cell tumid; antheridium 4-?; sperms 2, division horizontal; basal cell elongate; terminal cell, frequently an oogonium, obtuse; vegetative cell, female $11-26 \times 45-200\mu$; oogonium $56-68 \times 56-94\mu$; oospore $52-64 \times 44-68\mu$; antheridium $16 \times 16\mu$; suffultory cell $19-49 \times 30-145\mu$; basal cell $30-35 \times 45-160\mu$.

United States: Michigan; Denmark.

Cf. *Oe. scrobiculatum* above and *Oe. tiffanii* below. The antheridial filaments of *Oe. verrucosum* are very short, usually consisting only of a holdfast cell and a few antheridial cells.

64. *Oe. tiffanii* Ackley.

(Pl. XXX, figs. 261-263.)

1929, p. 304, Pl. XXXVI, figs. 15 and 16.

Dioecious, macrandrous; oogonium 1-2, subdepressed-globose or broadly pyriform-globose, pore superior; oospore globose, nearly filling

oogonium, median layer of spore wall scrobiculate; suffultory cell enlarged; antheridium 1-7, division horizontal, sperms 2; basal cell ellipsoid-elongate; vegetative cell $21-22 \times 100-240\mu$; suffultory cell $25-32 \times 100-230\mu$; oogonium $64-76 \times 65-69\mu$; oospore $54-65 \times 55-66\mu$; antheridium $19-21 \times 14-20\mu$.

United States: Michigan (Muskegon Lake).

Oe. tiffanii is to be compared with *Oe. scrobiculatum* (No. 62) and *Oe. verrucosum* above. These three normally have smaller suffultory cells and are thus separated from Nos. 65-69.

65. *Oe. punctatum* Wittrock.

(Pl. XXIX, figs. 255 and 256.)

1878, p. 142; De Toni 1889, p. 84; Hirn 1900, p. 132, Pl. XV, fig. 89.

Dioecious, macrandrous; oogonium 1-4, obovoid (rarely globose-obovoid), pore superior; oospore obovoid, nearly filling oogonium (rarely subglobose and not filling oogonium), outer spore wall scrobiculate; antheridium 1-5, often alternating with vegetative cell; sperms 2, division horizontal; basal cell elongate terminal cell (often oogonium) apically obtuse; vegetative cell $15-22 \times 42-128\mu$; oogonium $38-45 \times 52-65\mu$; oospore $37-43 \times 43-55\mu$; antheridium $15-17 \times 6-10\mu$.

Mexico.

Oe. punctatum, *Oe. argenteum* (No. 66), *Oe. wyliei* (No. 67), and *Oe. americanum* (No. 68) should be compared in identifying forms with scrobiculate oospores and non-tumid suffultory cells. They are easily separable on the basis of size.

66. *Oe. argenteum* Hirn.

(Pl. XXIX, figs. 253 and 254.)

Hirn 1900, p. 289, Pl. VII, fig. 47; Ackley 1929, p. 302.

Dioecious, macrandrous; oogonium 1, obovoid-globose to globose, pore superior (rarely supramedian); oospore ovoid to globose, outer layer of spore wall scrobiculate; antheridia 3-4, sperms 2, division horizontal; basal cell elongate; female vegetative cell (14-) $20-28 \times 80-160\mu$, male $20-22 \times 70-160\mu$; oogonium $44-52 \times 48-62\mu$; oospore $43-48 \times 44-50\mu$; antheridium $22 \times 8\mu$.

United States: Michigan; Brazil.

The figure of *Oe. argenteum* in Pl. XXIX is that of Hirn and represents also one of the variations occurring in the Michigan material. The outer oospore wall of the alga seen by Hirn (1900) is undoubtedly scrobiculate. Ornamentations on oospore walls are quite regularly constant for a given species, appearing definitely on one layer or another. The Michigan specimens (Ackley 1929) contained oospores whose middle wall layer was scrobiculate with an apparently smooth and very thin outer layer. It is perhaps best to include, for the present at least,

only those specimens that conform to the above description. If it develops that these forms with a middle scrobiculate wall may also have the outer wall scrobiculate, the description will have to be emended accordingly. The former had best be considered a form:

66a. Form **michiganese** Tiffany nov. f.

Oe. argenteum Hirn (in part) in Ackley 1929, p. 302, Pl. XXXV, fig. 3-6.

Middle wall of oospore scrobiculate; pore of oogonium suprmedian; otherwise as in the type.

United States: Michigan.

67. **Oe. wyliei** Tiffany.

(Pl. XXX, figs. 264-267.)

1926, p. 90, Pl. I, figs. 1-4.

Dioecious, macrandrous; oogonium 1-4, globose to ovoid, pore superior; oospore globose to ovoid, filling or not filling oogonium, outer spore wall irregularly scrobiculate; antheridium 1-4; sperms 2, division horizontal; basal cell elongate; terminal cell, often an oogonium, apically obtuse or broadly apiculate; vegetative cell $16-24 \times 80-170\mu$; oogonium $52-64 \times 68-112\mu$; oospore $48-60 \times 52-64\mu$; antheridium $16-19 \times 8-18\mu$.

United States: Iowa, Michigan.

See *Oe. punctatum* (No. 65).

68. **Oe. americanum** Transeau.

(Pl. XXXI, figs. 275 and 276.)

1917, p. 231.

Dioecious, macrandrous; oogonium 1, globose to depressed-globose, pore superior; oospore globose, ellipsoid-globose, or depressed-globose, filling oogonium or not, spore wall of three layers: median layer scrobiculate; antheridium 1-5, frequently alternating with vegetative cell; sperms 2, division horizontal; basal cell elongate; terminal cell obtuse; female vegetative cell $28-48 \times 40-100\mu$, male vegetative cell $24-30 \times 40-100\mu$; oogonium $40-76 \times 48-70\mu$; oospore $38-74 \times 46-56\mu$; antheridium $20-28 \times 4-12\mu$.

United States: Michigan, Illinois.

See *Oe. punctatum* (No. 65).

69. **Oe. taphrosporum** Nordstedt and Hirn.

(Pl. XXXI, figs. 270 and 271.)

In Hirn 1900, p. 133, Pl. XVI, fig. 91; Collins 1909, p. 239; P. B. A. No. 813.

Dioecious, macrandrous; oogonium 1-6, obovoid or obovoid-ellipsoid, pore superior; oospore globose or ellipsoid globose, not filling oogonium, outer layer of spore wall scrobiculate, inner layer smooth; antheridium 2-?; terminal cell, sometimes an oogonium, obtuse; basal cell elongate;

vegetative cell $25-38 \times 100-375\mu$; oogonium $70-83 \times 81-113\mu$; oospore $58-65 \times 62-70\mu$; antheridium $24-32 \times 8-12\mu$.

United States: Massachusetts, Michigan, Illinois; Brazil; India; Columbia.

This species is readily recognized by its large size and its scrobiculate oospore, the latter not filling the oogonium. West (1914) recorded the alga from Columbia with vegetative cells $19-22\mu$ in diameter, and with oogonia and oospores slightly smaller than the dimensions given above.

70. *Oe. pseudacrosporum* Wittrock.

(Pl. XXX, figs. 268 and 269.)

In Hirn 1900, p. 193, Pl. XXXII, fig. 196.

Monoecious; oogonium 1, ellipsoid, operculate, division supreme, lid very small and often deciduous; oospore ellipsoid, completely filling oogonium, spore wall coalescing with that of oogonium, longitudinally ribbed, ribs evidently crenulate; antheridium 1-4, hypogynous or sub-epigynous; sperms 2, division horizontal; basal cell elongate; terminal cell long, setiform; vegetative cell $9-13 \times 32-110\mu$; oogonium $(27-32-37 \times (40-45-56\mu$; antheridium $8-10 \times 6-10\mu$.

United States: Iowa, Florida; Sweden.

Its monoecious habit, supreme operculum, and ribbed oospore are characteristic. It should be compared with the nannandrous *Oe. acrosporum* (No. 195).

71. *Oe. paucocostatum* Transeau.

(Pl. XXXII, fig. 277.)

1914, p. 300, Pl. XXVIII, fig. 5; Collins 1918, p. 66; Tiffany 1926, Pl. V, figs. 55 and 56.

Dioecious, macrandrous; oogonium 1, ellipsoid, operculate, division superior; oospore ellipsoid, nearly filling the oogonium, outer wall smooth, median wall longitudinally ribbed, ribs 15-19 in number, inner wall smooth; antheridium 2-8, sperms 2, division horizontal; terminal cell obtuse, basal cell usually elongate; vegetative cell, female $(15-19-25 \times 70-160\mu$; oogonium $54-60 \times 70-104\mu$; oospore $50-56 \times 66-90\mu$; antheridium $18-23 \times 8-12\mu$.

United States: Illinois, Iowa, Ohio.

71a. Var. *gracilis* Tiffany.

(Pl. XXXII, fig. 278.)

1921, p. 273; 1926, p. 99.

Somewhat smaller than the species, oospore ellipsoid or occasionally globose-ellipsoid, completely filling the oogonium or not filling the oogonium; otherwise similar to the type; vegetative cell $15-20 \times 66-120\mu$; oogonium $48-52 \times 70-88\mu$; oospore $44-48 \times 60-70\mu$; antheridium $17-20 \times 8-12\mu$.

United States: Ohio, Iowa.

Oe. paucocostatum is characterized among the species with ellipsoid oogonia and longitudinally ribbed oospores by the small number of ribs and by its relatively smaller dimensions. Its nearest relative is the larger *Oe. australianum* (No. 72).

72. *Oe. australianum* Hirn.

(Pl. XXXII, figs. 283 and 284.)

1900, p. 192, Pl. XXXI, fig. 192.

Dioecious, macrandrous; oogonium 1-2, ellipsoid, operculate, division superior; oospore ellipsoid or globose-ellipsoid, practically filling oogonium, median layer of spore wall longitudinally ribbed, ribs 17-23, sometimes anastomosing; antheridium to 6-seriate; vegetative cell $17-22 \times 70-170\mu$; oogonium $59-74 \times 74-96\mu$; oospore $56-70 \times 59-74\mu$; antheridium $14-16 \times 10-15\mu$.

Australia.

Cf. *Oe. paucocostatum*, above.

73. *Oe. tumidulum* (Kuetzing) Wittrock.

(Pl. XXXII, figs. 281 and 282.)

1874, p. 35; *Conferva tumidula* English Botany No. 1670 in Kuetz. Dec. Alg. No. 60, 1833; Hirn 1900, p. 191, Pl. XXXI, fig. 191.

Dioecious, macrandrous; oogonium 1, subellipsoid, operculate, division superior; oospore ellipsoid or globose-ellipsoid, quite filling oogonium, outer layer of spore wall with 40-50 longitudinal ribs granulate and sometimes anastomosing, inner layer smooth; antheridium to 45-seriate; sperms 2, division horizontal; vegetative cell, female $18-25 \times 65-125\mu$, male $15-20 \times 60-120\mu$; oogonium $52-63 \times 78-90\mu$; oospore $49-58 \times 61-75\mu$; antheridium $15-20 \times 9-15\mu$.

Germany.

Oe. tumidulum is the only dioecious, macrandrous member of the genus having the combination of subellipsoid oogonia with superior pore and more than 40 longitudinal ribs on the oospore. Cf. the monoecious *Oe. insigne* (No. 75).

74. *Oe. nobile* Wittrock.

(Pl. XXXII, fig. 279.)

1874, p. 14; Hirn 1900, p. 189, Pl. XXX, fig. 188.

Monoecious; oogonium 1 (very rarely 2), ellipsoid or subobovoid-ellipsoid, operculate, division superior; oospore ellipsoid-globose or globose, not filling oogonium, spore wall of three layers: outer layer smooth, middle layer with 30-35 continuous longitudinal ribs, rarely anastomosing, inner layer smooth; antheridium 1-3, hypogynous; sperms 2, division horizontal; vegetative cell $16-20 \times 80-180\mu$; oogonium $57-65 \times 67-90\mu$; oospore $48-55 \times 50-58\mu$; antheridium $15-19 \times 9-13\mu$.

Norway.

74a. Var. *minus* Hirn.

(Pl. XXXII, fig. 280.)

1900, p. 190, Pl. XXX, fig. 189; *Oe. insignis* Hirn var. *minus* Hirn 1895, p. 15.

Larger oospore, ellipsoid, filling oogonium; vegetative cell $13-20 \times 80-240\mu$; oogonium $60-63 \times 85-100\mu$; oospore $56-59 \times 68-80\mu$; antheridium $16-19 \times 12-17\mu$.

United States: Massachusetts; Finland.

Oe. nobile is near the larger *Oe. insignis* (No. 75) and differs from *Oe. tumidulum* (No. 73) in being monoecious. The variety *minus* sometimes has smaller vegetative cells and always a larger oospore than the type.

75. *Oe. insignis* Hirn.

(Pl. XXXIII, fig. 285.)

1895, p. 14, Pl. 1, fig. 2; 1900, p. 191, Pl. XXX and XXXI, fig. 190.

Monoecious; oogonium 1, ellipsoid-obovoid, operculate, division superior; oospore ellipsoid, quite filling oogonium, spore wall of three layers: outer smooth, inner longitudinally ribbed, the ribs quite entire, anastomosate, $40-45$ in number, inner smooth; antheridium $1-7$, subepigynous, rarely scattered; sperms 2, division horizontal; vegetative cell ($18-$) $25-38 \times 75-210\mu$; oogonium $70-78 \times 100-120\mu$; oospore $65-75 \times 88-104\mu$; antheridium $23-27 \times 15-23\mu$.

Finland.

Cf. *Oe. tumidulum* (No. 73) and *Oe. nobile* (No. 74).

76. *Oe. excisum* Wittrock and Lundell.

(Pl. XXXIV, fig. 317.)

In Wittr. 1872, p. 3, Pl. 1, figs. 1-4; Hirn 1900, p. 153, Pl. XXIV, fig. 126.

Monoecious; oogonium 1, subconically oblong, medianly plicate with about 9 longitudinal undulations, operculate, division median and broad; oospore ellipsoid with a median constriction, not filling oogonium, spore wall smooth; antheridium $1-2$, subepigynous or hypogynous, often terminal; sperm (?) 1; basal cell subhemispherical; terminal cell apically obtuse; vegetative cell $3-6 \times 14-44\mu$; oogonium $13-15 \times 18-26\mu$; oospore $9-12 \times 15-18\mu$; antheridium $3-4 \times 6-7\mu$; basal cell $8-9 \times 4-5\mu$.

Finland, Sweden, Latvia.

Oe. excisum is one of the very few species of the genus having an oospore with a median constriction. It is near *Oe. pusillum* (No. 221), from which it differs in having undulate folds in the oogonium.

77. *Oe. inversum* Wittrock.

(Pl. XXXIII, figs. 295-297.)

1876, p. 47, Pl. 13, figs. 22-24; (?) *Oe. monticchii* Fiorini-Mazzanti 1860, p. 259, Pl. 1, figs. 3, 4, 4a, 4b; Hirn 1900, p. 179, Pl. XXVIII, fig. 171; *Oe. inversum* Wittr. var. *subclusum* Wittr. in Wittr. and Nordst. Alg. Exs. No. 26, 1877; *Oe. inversum* Wittr. f. *subclusum* (Wittr.) Hirn 1900, p. 180, Pl. XXVIII, fig. 172; Heering 1914, p. 203, fig. 290; Tiffany 1926, p. 95, Pl. VI, figs. 59-62.

Dioecious, macrandrous; oogonium 1, globose, operculate, division inferior; oospore globose, quite filling the oogonium, spore wall smooth; antheridium 1-8; sperm 1; vegetative cell capitellate; basal cell subhemispherical, not elongate; filament not infrequently incrustated with lime; female vegetative cell $12-14 \times 25-100\mu$, male $9-11 \times 20-80\mu$; oogonium (28-) $32-35 \times 30-34\mu$; oospores (27-) $30-32 \times 27-30\mu$; antheridium $10-12 \times 9-12\mu$; basal cell $13-20 \times 7-14\mu$.

United States: Iowa; Australia; Austria, France, Germany, Sweden, Switzerland; Africa.

Easily distinguished among the macrandrous species by the inferior position of the operculum. It is in some respects similar to *Oe. infimum* below. The f. *subclusum*, according to the Iowa material (Tiffany 1926), is merely an ecological variant and is not tenable. The variations of the species proper include the form.

78. *Oe. infimum* Tiffany.

(Pl. XXXIII, figs. 289-292.)

1924, p. 183, Pl. II, figs. 6-9; 1926, p. 95, Pl. VI, figs. 59-62.

Dioecious, macrandrous; oogonium 1, globose or subglobose (or the basal part extended, appearing subpyriform-globose), operculate, division at the lowest extremity of the oogonium; oospores globose or subglobose, membrane smooth; male plants a little larger than the female; antheridium 1-10; sperms 2; vegetative cells distinctly capitellate; basal cell of filament subhemispherical, not elongated; filaments not infrequently incrustated with lime; female vegetative cells $12-18 \times 60-140\mu$, male $16-20 \times 60-140\mu$; oogonium $40-48 \times 41-50\mu$; oospore $40-44 \times 38-42\mu$; antheridium $14-20 \times 8-12\mu$; basal cell $30-42 \times 16-24\mu$.

United States: Iowa.

Oe. infimum is the only known member of the genus with the operculum located at the lowest extremity of the oogonium. It is further characterized by its evidently capitellate vegetative cells.

79. *Oe. howardii* West.

(Pl. XXXIII, fig. 293.)

1904a, p. 281, Pl. 464, figs. 1-5; Hirn 1906, p. 16, Pl. III, fig. 9.

Dioecious, macrandrous; oogonium 1-2, globose (or subglobose), operculate, division median, distinct; oospore globose (or subdepressed-globose) filling oogonium; antheridium to 16-seriate, sperm (?) 1; vegetative cell broadly capitellate; basal cell subhemispherical, scarcely elongate; vegetative cell, female $9-12 \times 20-44\mu$; male $7-9 \times 16-35\mu$; oogonium $29-33 \times 29-33\mu$; oospore $25-29 \times 25-29\mu$; antheridium $7-9 \times 8-14\mu$; basal cell $14-16 \times 10-11\mu$.

West Indies: Barbados.

79a. Var. *minus* Tiffany.

(Pl. XXXIII, fig. 294.)

1927, p. 204, Pl. IX, figs. 6 and 7 (as v. *minor*).

Smaller than the type; vegetative cell $7-9 \times 27-48\mu$; oogonium $26-29 \times 28-36\mu$; oospore $24-26 \times 24-26\mu$; antheridium $7-8 \times 10-12\mu$; basal cell $12-14 \times 10-12\mu$.

United States: Illinois.

Oe. howardii has broadly capitellate vegetative cells as in *Oe. infimum* (No. 78) and *Oe. inversum* (No. 77), but differs from both in the median position of its operculum. It is perhaps nearest *Oe. latiusculum* (No. 81), the latter being larger and possessing a very wide division of the oogonium. The variety *minus* is characterized chiefly by its smaller size and subspherical basal cell.

80. *Oe. pratense* Transeau.

(Pl. XXXIV, figs. 298 and 299.)

1914, p. 297, Pl. XXIX, figs. 9-12; Collins 1918, p. 66; Tiffany 1926, p. 96, Pl. I, figs. 10 and 11.

Dioecious, macrandrous; oogonium 1 (rarely 2), subdepressed-globose or broadly pyriform-globose, operculate, division median, narrow but distinct; oospore depressed-globose or subglobose, filling or nearly filling oogonium, spore wall smooth; antheridium 1-2, usually alternating with vegetative cell; sperm 1; basal cell elongate; vegetative cell, female $10-17 \times 35-95\mu$, male $8-15 \times 32-82\mu$; oogonium $33-40 \times 35-50\mu$; oospore $32-38 \times 28-35\mu$; antheridium $10-14 \times 13-18\mu$.

United States: Illinois, Iowa, Ohio, Alabama, Mississippi, Michigan.

Oe. pratense is distinguished from *Oe. latiusculum* below by the narrow opening of the operculum and by its cylindrical vegetative cells. The latter character together with its larger size separates it from *Oe. howardii* (No. 79). The basal cell of *Oe. pratense* is uniformly elongate. Among the monoecious forms it bears some resemblance to *Oe. acmandrium* (No. 82) and *Oe. psaegetosporum* (No. 83).

81. *Oe. latiusculum* Tiffany.

(Pl. XXXIII, figs. 287 and 288.)

1924, p. 182, Pl. III, figs. 4-6; 1926, p. 96, Pl. VI, figs. 72-74.

Dioecious, macrandrous; oogonium 1-2, globose to ellipsoid-globose; oogonium operculate, division median, very wide; oospore of the same form as the oogonium which it very nearly or completely fills, wall smooth; male plant somewhat larger than the female; antheridium 1, usually alternating with a single vegetative cell, sperm 1; vegetative cell distinctly capitellate; basal cell of filament subhemispherical, not elongated; filament not infrequently incrustated with lime; female veg-

etative cell $10-18 \times 16-40\mu$, male $14-20 \times 16-40\mu$; oogonium $32-36 \times 32-40\mu$; oospore $28-32 \times 28-34\mu$; antheridium $14-18 \times 12-20\mu$; basal cell $16-24 \times 12-16\mu$.

United States: Iowa, Mississippi, Alabama, Kentucky.

The wide operculum, the capitellate vegetative cells, the single antheridium alternating with a vegetative cell, and the subhemispherical basal cell are characteristic of the species.

82. *Oe. acmandrium* Elfving.

(Pl. XXXIV, fig. 300.)

In Hirn 1895, p. 13; Hirn 1900, p. 150, Pl. XXIII, fig. 120.

Monoecious; oogonium 1-2, depressed-globose or subglobose, operculate, division median and narrow; oospore depressed-globose or subglobose, filling oogonium, wall smooth; antheridium 1-3, subepigynous, epigynous, or hypogynous, often terminal; sperm single; basal cell elongate; vegetative cell $7-10 \times 30-80\mu$; oogonium $30-35 \times 28-38\mu$; oospore $28-33 \times 25-29\mu$; antheridium $8-10 \times 10-15\mu$.

United States: Illinois; Finland, Sweden; India.

Cf. *Oe. pratense* (No. 80) and *Oe. psaegmatosporum* below.

83. *Oe. psaegmatosporum* Nordstedt.

(Pl. XXXIV, fig. 301.)

1877, p. 24, Pl. 3, fig. 1-3; Hirn 1900, p. 150, Pl. XXIII, fig. 121.

Monoecious; oogonium 1-5, broadly pyriform-globose, operculate, division median, narrow but distinct; oospore depressed-globose to globose, inflating the middle part of oogonium, spore wall (?) smooth; antheridium to 15-seriate, hypogynous, cell a little tumid; sperm single; vegetative cell $9-10 \times 56-80\mu$; oogonium $28-33 \times 33-40\mu$; oospore $27-31 \times 24-27\mu$; antheridium $9-12 \times 6-10\mu$.

Denmark, Sweden.

Near *Oe. acmandrium* above, differing in its pyriform-globose and medianly inflated oogonia and many-seriate antheridia.

84. *Oe. minus* Wittrock.

(Pl. XXXIV, fig. 302.)

1874, p. 9; *Oe. punctato-striatum* var. *minor* Wittr. 1870, p. 123; *Oe. spirogranulatum* Schmidle 1894, p. 43, Pl. 7, fig. 1; Hirn 1900, p. 151, Pl. XXIII, fig. 122.

Monoecious; oogonium 1, pyriform-globose or depressed-globose, operculate, division median, wide; oospore depressed-globose, not filling oogonium, spore wall smooth; antheridium to 10-seriate, subepigynous or subhypogynous or rarely scattered, cells a little tumid; sperm 1; vegetative cell somewhat capitellate; wall of vegetative cell and of oogonium spirally punctate; basal cell depressed-globose or subhemispherical, wall vertically plicate; vegetative cell $9-13 \times 30-78\mu$; oogonium $34-46 \times 28-42\mu$; oospore $30-42 \times 26-36\mu$; antheridium $9-13 \times 3-5\mu$.

United States: Massachusetts, Michigan; Finland, Germany, Spain, Norway, Sweden, Latvia, Russia.

Oe. minus is one of the few species of the genus with spirally punctate vegetative cell and oogonial walls. It is easily separable from the nannandrous *elegans* (No. 155) and the macrandrous *punctatostriatum* (No. 85) not only by its monoecious habit but also by its size.

85. *Oe. punctatostriatum* De Bary.

(Pl. XXXIV, figs. 303 and 304.)

1854, p. 47, Pl. 2, figs. 15 and 16; Hirn 1900, p. 152, Pl. XXIII, fig. 123.

Dioecious, macrandrous; oogonium 1, depressed-globose, operculate, division median, rather wide and distinct; oospore depressed globose, not filling oogonium, spore wall smooth; antheridium to 10-seriate; sperm 1; wall of vegetative cell and oogonium spirally punctate; basal cell depressed-globose or subhemispherical, wall vertically plicate; female vegetative cell $18-22 \times 38-128\mu$, male $16-19 \times 33-108\mu$; oogonium $48-55 \times 38-48\mu$; oospore $40-51 \times 35-42\mu$; antheridium $16-19 \times 6-12\mu$; basal cell $28-31 \times 21-25\mu$.

United States: Iowa, Illinois, Florida, Michigan; England, Finland, France, Ireland, Germany, Norway, Russia, Latvia, Sweden; Brazil; Greenland; Australia; British Columbia.

Cf. *Oe. minus* above.

86. *Oe. capitellatum* Wittrock.

(Pl. XXXIV, fig. 315.)

1874, p. 7; *Oe. piliferum* Wittr. 1870, p. 122, and 1872, p. 23; Hirn 1900, p. 149, Pl. XXIII, fig. 18.

Monoecious; oogonium 1, subdepressed or depressed globose, operculate, division median, narrow but distinct; oospore depressed globose, completing (or nearly so) oogonium, spore wall smooth; antheridium 1-2, subepigynous or hypogynous or rarely scattered; sperm (?) 1; vegetative cell capitellate; basal cell subhemispherical; terminal cell piliform; vegetative cell $6-9 \times 20-60\mu$; oogonium $20-25 \times 17-23\mu$; oospore $18-23 \times 15-19\mu$; antheridium $6-7 \times 8-9\mu$; basal cell $16-18 \times 6-8\mu$.

United States: Ohio, Illinois, Mississippi, Alabama; British Columbia; Denmark, Finland, Sweden; Turkestan; Caucasus; Burma.

Among the species with capitellate vegetative cells it is characterized by its monoecious habit and small size. It differs from *Oe. quadratum* below in the absence of angular-globose oogonia.

87. *Oe. quadratum* Hallas.

(Pl. XXXIV, figs. 306-308.)

1905, p. 405, fig. 12; Hirn 1906, p. 19, Pl. III, fig. 10.

Monoecious; oogonium 1-4, angular-globose, operculate, division median; oospore angular-globose, filling oogonium, spore wall smooth; antheridium 1, alternating with vegetative cell; vegetative cell capitellate; basal cell subhemispherical; vegetative cell $5-12 \times 25-140\mu$; oogonium $22-28 \times 28-33\mu$; oospore $20-26 \times 26\mu$; antheridium $10-12 \times 12-14\mu$; basal cell $7-12 \times 6-9\mu$.

Denmark.

Cf. *Oe. capitellatum* above.88. *Oe. bohemicum* Hirn.

(Pl. XXXVI, fig. 344.)

1900, p. 169, Pl. XXVII, fig. 154; Heering 1914, p. 216, fig. 327; Tiffany 1926, p. 101, Pl. III, fig. 30.

Monoecious; oogonium 1, globose, operculate, division superior; oospore globose, filling oogonium, spore walls smooth; antheridium 1-4, subepigynous; sperms 2, division horizontal; vegetative cell capitellate, $10-16 \times 21-66\mu$; oogonium $42-45 \times 46-49\mu$; oospore $40-43 \times 40-43\mu$; antheridium $9-10 \times 5-7\mu$.

United States: Iowa; Austria.

This is the largest of the monoecious species combining the median location of the operculum and the capitellate vegetative cell. The dimensions above, from the Iowan material, also include those of the Austrian specimens.

89. *Oe. mitratum* Hirn.

(Pl. XXXVI, figs. 334 and 335.)

1895, p. 22, Pl. 1, fig. 7; 1900, p. 302, Pl. XXIV, fig. 132.

Dioecious, macrandrous; oogonium 1-4, globose or subglobose, operculate, division suprmedian or superior, narrow but distinct; oospore globose (rarely subglobose), filling oogonium, spore wall smooth; antheridium 1-5, subepigynous, sperms 2, division horizontal; vegetative cell often broadly capitellate; vegetative cell $5-10 \times 18-80\mu$; oogonium $18-24 \times 20-28\mu$; oospore $17-23 \times 17-22\mu$; antheridium $6-9 \times 6-8\mu$.

United States: Illinois; Austria, Finland, Sweden.

Hirn (1900) records this species as doubtfully monoecious, but material from Illinois (Transeau) shows it to be definitely dioecious, macrandrous. The emended description was supplied from the Illinois material. Cf. *Oe. petri* below.

90. *Oe. petri* Wittrock.

(Pl. XXXVI, fig. 332.)

1874, p. 6; Hirn 1900, p. 154, Pl. XXIV, fig. 127.

Monoecious; oogonium 1 (rarely 2), pyriform or pyriform-globose, operculate, division a little above median; oospore subdepressed or

depressed-globose, filling oogonium, spore wall smooth; antheridium 1-2, hypogynous or epigynous or subepigynous; sperm 1; basal cell elongate; terminal cell apically obtuse; vegetative cell $6-7 \times 32-48\mu$; oogonium $21-24 \times 22-29\mu$; oospore $20-23 \times 16-19\mu$; antheridium $5-7 \times 9-11\mu$.

Germany, Spain, Sweden; Ceylon.

91. *Oe. trioicum* Woronichin.

1923, p. 99.

Dioecious macrandrous, or monoecious; oogonium 1-3, sometimes terminal, ellipsoid, operculate, division superior; oospore ellipsoid or subovoid, spore wall smooth, faintly violet; antheridium 1-6, hypogynous in monoecious filament, scattered or terminal in male filament; terminal cell obtusely rounded; basal cell attenuate; male plants $130-330\mu$ long; vegetative cell $4-5 \times 12-22\mu$; oogonium $25-27 \times 13-16\mu$; oospore $20-21 \times 14\mu$; antheridium $4 \times 3\mu$.

Near Baku, in Azerbaijan (west coast of the Caspian).

This plant according to Woronichin is both dioecious, macrandrous and monoecious, in this respect similar to *Oe. varians* (No. 10). It should be compared with the macrandrous *pringsheimii* (No. 95) and the monoecious *gracillimum* (No. 114). Unfortunately no figure is available.

92. *Oe. welwitschii* West and West.

(Pl. XXXVII, figs. 351 and 352.)

1897, p. 5; Hirn 1900, p. 174, Pl. XXVIII, fig. 162; Tiffany 1926, p. 97, Pl. IV, figs. 41 and 42.

Dioecious, macrandrous; oogonium 1-3, subovoid-globose, operculate, division superior; oospore globose, filling or not filling oogonium, spore wall smooth; antheridium 2; female vegetative cell $20-28 \times 25-84\mu$, male $17-20 \times 35-80\mu$; oogonium $43-50 \times 43-52\mu$; oospore $35-43 \times 35-42\mu$; antheridium $16-18 \times 7-9\mu$.

United States: Iowa, Illinois, Alabama; Africa.

This species is near *Oe. iowense* below, but differs in having at once both larger vegetative cells and smaller oogonia. It may be near the imperfectly known *dioicum* (No. 94) and is larger than *Oe. pringsheimii* (No. 95).

93. *Oe. iowense* Tiffany.

(Pl. XXXVII, figs. 353-355.)

1924, p. 181, Pl. III, figs. 1-3; 1926, p. 97, Pl. VI, figs. 69-71.

Dioecious, macrandrous; oogonium 1-2, globose, or ellipsoid-globose, operculate, division superior; oospore of the same form as the oogonium, which it completely fills or not, spore wall smooth; antheridium 1-25, sperms 2, division horizontal; basal cell of filament commonly elongated; vegetative cell $10-16 \times 44-100\mu$; oogonium $52-60 \times 60-80\mu$; oospore $45-56 \times 50-64\mu$; antheridium $10-12 \times 10-20\mu$; basal cell $16-24 \times 60-80\mu$.

United States: Iowa, Ohio, Kentucky.

This species is easily recognized by the considerable difference between the diameter of the vegetative cells and that of the relatively much larger oogonia. Its oogonia are larger than those of *Oe. welwitschii* above.

94. *Oe. dioicum* Carter.

(Pl. XXXVII, figs. 356 and 357.)

1858, p. 30, Pl. 3, figs. 1-2, 5-8, 13-16; Hirn 1900, p. 175, Pl. XXVIII, fig. 163.

Dioecious, macrandrous; oogonium 1, obovoid-globose, operculate, division superior; oospore globose, not filling oogonium, spore wall smooth; male plants a little smaller than female; antheridium 10-20, sperms (?) 2, division vertical; female vegetative cell about 32×80 - 110μ , male about 28×70 - 95μ ; oogonium about $82 \times 92\mu$.

India.

No data beyond those of Carter are available, and *Oe. dioicum* is retained here largely because it cannot be definitely discarded upon our present knowledge of the species.

95. *Oe. pringsheimii* Cramer; Wittrock.

(Pl. XXXV, figs. 325 and 326.)

Cramer 1859, p. 17, Pl. 1, fig. C 1-4; Wittr. 1870, p. 135; Wittr. 1874, p. 33, Pl. 1, figs. 16 and 17; Hirn 1900, p. 170, Pl. XXVII, fig. 155; *Oe. pachydermatosporum* Nordst. 1878, p. 21, Pl. 2, fig. 13-15; *Oe. pringsheimii* Cram.; Wittr. var. *Nordstedtii* Wittr. f. *pachydermatosporum* (Nordst.) Hirn 1900, p. 173, Pl. XXVII, fig. 160; Collins 1909, p. 246; Heering 1914, p. 202, fig. 284; Tiffany 1926, p. 98, Pl. V, figs. 45 and 46.

Dioecious, macrandrous; oogonium 1-6, subovoid-globose, operculate, division superior; oospore globose, nearly filling oogonium, spore wall smooth, often thickened; antheridium to 10-seriate, often alternating with vegetative cell; sperms 2, division horizontal; basal cell elongate, terminal cell broadly apiculate or obtuse; female vegetative cell 14 - 20×28 - 100μ , male 12 - 16×24 - 64μ ; oogonium 35 - 43×36 - 46μ ; oospore 30 - 37×30 - 37μ ; antheridium 10 - 15×6 - 9μ .

United States: Iowa, New York, Pennsylvania, Florida, Ohio, Utah; Africa, Afghanistan; Australia; Austria, Denmark, England, Finland, Italy, Norway, Latvia, Sweden, Switzerland.

95a. Var. *abbreviatum* Hirn.

(Pl. XXXV, fig. 331.)

1900, p. 173, Pl. XXVII, fig. 161.

Smaller; vegetative cell broader; oogonium 1, obovoid-globose, nearly filled by oospore; vegetative cell 10 - 13×15 - 39μ ; oogonium 28 - 32×30 - 35μ ; oospore 27 - 30×27 - 30μ .

Brazil.

95b. Var. *nordstedtii* Wittrock.

(Pl. XXXV, figs. 327-330.)

In Wittr. and Nordst. Alg. Exs. No. 8, 1877; *Oe. ciliare* De Notaris 1868, p. 120; *Oe. franklinianum* Wittr. in Tilden Amer. Alg. No. 2, 1894; *Oe. nordstedtii* Wittr. 1872, p. 6, Pl. I, figs. 7 and 8; *Oe. pringsheimii* var. *varians* Nordst. 1888, p. 11, Pl. I, fig. 9; *Oe. pachydermatosporum* Nordst. in Lewin 1888, p. 18, Pl. 3, fig. 56; Hirn 1900, p. 171, Pl. XXVII, figs. 156-158; *Oe. pringsheimii* Cram.; Wittr. var. *nordstedtii* Wittr. f. *euganeorum* (Wittr.) Hirn 1900, p. 172, Pl. XXVII, fig. 159; *Oe. euganeorum* Wittr. 1874, p. 39; Tiffany 1926, p. 98, Pl. V, fig. 47.

Smaller than the type, oogonium 1-2; oospore not filling oogonium; female vegetative cell 10-16×20-76μ, male 9-15×18-68μ; oogonium 28-39×36-45μ; oospore 26-34×27-34μ; antheridium 9-12×8-9μ.

United States: Minnesota, California, Iowa, Illinois, Ohio, Michigan; Austria, England, Finland, Latvia, Germany, Italy, Switzerland, Spain; India; Patagonia; Greenland; Samoa.

Oe. pringsheimii is characterized by its subovoid-globose oogonia and superior position of the operculum, among the dioecious species. Its oogonia are smaller than either *Oe. iowense* (No. 93) or *Oe. welwitschii* (No. 92). The variety *abbreviatum* has shorter vegetative cells and var. *nordstedtii* is more slender than the type.

96. *Oe. nodulosum* Wittrock.

(Pl. XXXIX, figs. 374 and 375.)

1872a, p. 22, Pl. I, figs. 8-10; 1874, p. 13, Pl. I, figs. 2 and 3; Borge 1899, p. 6, Pl. I, fig. 3; Hirn 1900, p. 187, Pl. XXIX, fig. 184.

Monoecious; oogonium 1-2, obovoid-globose or more rarely obovoid-ellipsoid, operculate, division superior; oospore globose or subglobose or more rarely globose-ellipsoid, nearly filling oogonium, spore wall smooth, often thick; antheridium 1-3, subepigynous or hypogynous; sperms 2, division horizontal; basal cell elongate, not constricted; terminal cell obtuse or apiculate; vegetative cell, with two undulate constrictions, 20-29×30-140μ; oogonium 48-57×56-73μ; oospore 46-53×49-56μ; antheridium 18-25×7-9μ.

Australia; China; Patagonia; England, Sweden.

96a. Var. *commune* Hirn.

(Pl. XXXIX, fig. 376.)

1900, p. 187, Pl. XXX, fig. 185; *Oe. nodulosum* Wittr. in P. B. A. No. 74, 1895, and in Hirn 1895, p. 13; Collins 1909, p. 248.

Oogonium subobovoid-ellipsoid to ellipsoid or more rarely globose-ellipsoid; vegetative cell 22-29×35-140μ; oogonium 64-74×70-90μ; oospore 56-70×67-80μ; antheridium 18-26×7-10μ.

United States: Massachusetts; Finland, Sweden.

The nodulose character of the vegetative cells is very distinctive. *Oe. sphaerandrium* (No. 105) sometimes has nodulose cells, but it is much smaller. Var. *commune* has larger

oogonia and oospores. *Oe. undulatum* (No. 119), a nannandrous species, has vegetative cells four times undulate-constricted while *Oe. nodulosum* has two undulate constrictions.

97. *Oe. porrectum* Nordstedt and Hirn.

(Pl. XXXVIII, figs. 361 and 362.)

In Hirn 1900, p. 186, Pl. XXIX, fig. 183.

Dioecious, macrandrous; oogonium 1, oblong, operculate, division superior; oospore ellipsoid or globose-ellipsoid, not filling oogonium longitudinally, spore wall smooth; antheridium to 4-seriate; sperms 2, division horizontal; female vegetative cell $7-10 \times 25-55\mu$, male $6-9 \times 25-62\mu$; oogonium $24-27 \times 44-53\mu$; oospore $23-24 \times 25-28\mu$; antheridium $6-7 \times 6-8\mu$.

Brazil.

98. *Oe. nanum* Wittrock; Tiffany.

(Pl. XXXVI, figs. 342 and 343.)

Wittrock 1874, p. 37; Hirn 1900, p. 305, Pl. XXIX, fig. 174; Tiffany 1926, Pl. VI, figs. 63-65.

Dioecious, macrandrous; oogonium 1-3, ovoid to broadly ellipsoid, operculate, division superior; oospore ovoid to globose-ellipsoid, usually filling the oogonium, spore wall smooth; antheridium 1-3, sperm 1; basal cell subhemispherical; terminal cell, often an oogonium, apically obtuse; vegetative cell, often irregularly swollen, $6-10 \times 15-33\mu$; oogonium $24-28 \times 30-36\mu$; oospore $21-27 \times 23-30\mu$; antheridium $7-10 \times 8-11\mu$; basal cell $14-17 \times 12-16\mu$.

United States: Iowa; India.

Originally found in India, this species was only completely described from the Iowan material. It bears some resemblance to *Oe. pisanum* below. It is found almost entirely epiphytic on submerged leaves, on Pithophora, on Cladophora, and on other species of Oedogonium.

99. *Oe. pisanum* Wittrock.

(Pl. XXXVIII, figs. 363 and 364.)

1876, p. 50, Pl. 13, fig. 28; *Oe. subpisanum* Lewin 1888, p. 17, Pl. 3, figs. 49-52; Hirn 1900, p. 181, Pl. XXIX, fig. 175; Heering 1914, p. 203, fig. 291; Tiffany 1926, p. 97.

Dioecious, macrandrous; oogonium 1 (rarely 2-3), ellipsoid-ovoid to ovoid, operculate, division superior; oospore ovoid to ellipsoid, nearly filling oogonium, spore wall smooth; antheridium 1-4; sperms 2, division horizontal; basal cell elongate; terminal cell piliferous; vegetative cell $5-12 \times 12-72\mu$; oogonium $23-29 (-32) \times 34-43 (-45)\mu$; oospore $21-25 \times 27-37\mu$; antheridium $4-9 \times 5-9\mu$.

United States: Iowa, Illinois, Ohio, Mississippi; Austria, Italy, Spain; Africa.

99a. Var. **gracilis** Transeau and Tiffany.
(Pl. XXXVIII, figs. 365 and 366.)

1919, p. 241; Tiffany 1926, p. 98, Pl. III, figs. 31-34.

Oogonium and oospore ellipsoid to ellipsoid-ovoid; vegetative cell, female $6-9 \times 18-48\mu$, male $5-8 \times 16-45\mu$; oogonium $16-20 \times 20-30\mu$; oospore $15-18 \times 18-28\mu$; antheridium $5-7 \times 5-10\mu$.

United States: Iowa, Illinois, Mississippi.

The combination of small size, elongate basal cell, and piliferous terminal cell is sufficient to characterize the species. The presence of subhemispherical basal cells in *Oe. nanum* (No. 98) easily separates it from *Oe. pisanum*. The variety *gracilis* is smaller. Often epiphytic on other species of algae.

100. **Oe. pyriforme** Wittrock.
(Pl. XXXVII, fig. 348.)

1874, p. 39; Hirn 1900, p. 303, Pl. XXV, fig. 137.

Monoecious; oogonium 1, pyriform, operculate, division superior; oospore pyriform, filling or not filling oogonium, spore wall smooth; antheridium 1-3, subepigynous, epigynous, hypogynous, or scattered; sperms 2, division horizontal; vegetative cell $13-16 \times 48-90\mu$; oogonium $40-46 \times (44-48-60\mu)$; oospore $36-42 \times 36-44\mu$; antheridium $10-12 \times 8-12\mu$.

Tasmania; British Columbia.

The pyriform oogonium is characteristic. The dimensions of *Oe. pyriforme* separate it from *Oe. simplex* (No. 102) and *Oe. pyrulum* (No. 103). Originally found in Tasmania, the above complete description is furnished from material collected in British Columbia by Mr. G. H. Wailes.

101. **Oe. gunnii** Wittrock.
(Pl. XXXIV, fig. 318.)

1874, p. 37; Hirn 1900, p. 298, Pl. XXIII, fig. 119.

Monoecious; oogonium 1-4, subdepressed- or depressed-globose, operculate, division median, narrow but distinct; oospore same form as oogonium and filling it, spore wall smooth, outer layer thick and hyaline, inner brown; antheridium subepigynous; vegetative cell $6-9 \times 30-85\mu$; oogonium $23-29 \times 19-29\mu$; oospore $22-27 \times 17-23\mu$; antheridium $6 \times 12\mu$.

United States: Alabama; Australia.

Material collected in Alabama by Professor E. N. Transeau shows *Oe. gunnii* to be definitely monoecious. Its distinctly narrow median division of the oogonium separates it from the imperfectly known *Oe. poecilosporum* (No. 224).

102. *Oe. simplex* Hirn.

(Pl. XXXVI, fig. 340.)

1900, p. 158, Pl. XXIV, fig. 135.

Monoecious; oogonium 1, obovoid-pyriform, operculate, division superior; oospore globose, ellipsoid or obovoid-globose, not filling oogonium, spore wall smooth; antheridium 1-?, subepigynous; sperms 2, division horizontal; vegetative cell $11-13 \times 35-58\mu$; oogonium $26-32 \times 33-40\mu$; oospore (22-) $24-30 \times 23-30\mu$; antheridium $8-10 \times 7-10\mu$.

Brazil.

Cf. *Oe. pyriforme* (No. 100).103. *Oe. pyrulum* Wittrock.

(Pl. XXXVII, fig. 349.)

1872, p. 2; Hirn 1900, p. 158, Pl. XXV, fig. 136.

Monoecious; oogonium 1, globose-pyriform, operculate, division superior; oospore subdepressed-globose or subglobose, not filling oogonium, spore wall smooth; antheridium 1, epigynous or hypogynous or rarely subepigynous; basal cell elongate; terminal cell apically broadly acute; vegetative cell $8-11 \times 30-75\mu$; oogonium $30-33 \times 31-35\mu$; oospore $26-29 \times 24-28\mu$; antheridium $8-9 \times 10-11\mu$.

Sweden; Ceylon.

103a. Var. *amplius* W. R. Taylor.

(Pl. XXXVII, fig. 350.)

1928, p. 107, Pl. 13, figs. 1-4 (as v. *amplior*).

Larger oogonium; vegetative cell $9-12 \times 40-64\mu$; oogonium $34-40 \times 30-44\mu$; oospore $26-28 \times 26-28\mu$, not filling oogonium; antheridium subepigynous $10 \times 6\mu$.

British Columbia.

Oe. pyrulum is separated from the species of the genus having pyriform or subpyriform oogonia largely by dimensions. The variety *amplius* has larger oogonia than the type and more nearly approaches the size of *Oe. pyriforme* (No. 100).

104. *Oe. loricaum* Hirn.

(Pl. XXXVI, fig. 339.)

1895, p. 22, Pl. 1, fig. 6; 1900, p. 156, Pl. XXIV, fig. 133.

Monoecious; oogonium 1 (rarely 2), subpyriform-globose, sometimes subglobose, operculate, division superior; oospore subglobose or subdepressed-globose, quite filling oogonium, spore wall smooth; antheridium 1-2, subepigynous; sperms 2, division horizontal; vegetative cell $8-11 \times 30-75\mu$; oogonium $23-28 \times 23-40\mu$; oospore $22-26 \times 21-24\mu$; antheridium $7-8 \times 5-7\mu$.

Finland.

This species is similar to *Oe. pithophorae* (No. 106) below, but the latter has globose and slightly larger oospores.

105. *Oe. sphaerandrium* Wittrock and Lundell.

(Pl. XXXVI, fig. 337.)

In Wittr. 1874, p. 7; *Oe. subcapitellatum* Hirn 1895, p. 13, Pl. 1, fig. 1; Hirn 1900, p. 155, Pl. XXIV, figs. 129 and 130; Borge 1913, p. 31, fig. 2.

Monoecious; oogonium 1-4, subpyriform- to subdepressed-globose, operculate, division a little above median; oospore subdepressed- or depressed-globose, nearly filling oogonium, spore wall smooth, antheridium to 6-seriate, subepigynous or scattered; sperm 1; vegetative cell capitellate, sometimes nodulose; basal cell subhemispherical, terminal cell piliform; vegetative cell $4-10 \times 6-40 \mu$; oogonium $18-23 \times 18-23 \mu$; oospore $16-21 \times 14-19 \mu$; antheridium $6-8 \times 5-6 \mu$.

Austria, Finland, Germany, Sweden; Siberia.

The capitellate (sometimes nodulose) vegetative cells, the small size, and suprmedian operculum are characteristic.

106. *Oe. pithophorae* Wittrock.

(Pl. XXXVI, fig. 338.)

1878, p. 141; Hirn 1900, p. 157, Pl. XXIV, fig. 134.

Monoecious; oogonium 1, pyriform-globose, operculate, division superior; oospore globose, almost filling oogonium, spore wall smooth, often thickened; antheridium 1-?, subepigynous; basal cell elongate; vegetative cell $9-11 \times 24-48 \mu$; oogonium $26-30 \times 27-35 \mu$; oospore $25-29 \times 24-29 \mu$; antheridium $8-10 \times 7-9 \mu$.

St. Thomas, West Indies.

Epiphytic on *Pithophora cleveana* Wittrock.

Cf. *Oe. porrectum* (No. 97).

107. *Oe. autumnale* Wittrock.

(Pl. XXXVI, fig. 341.)

1874, p. 11; Hirn 1900, p. 167, Pl. XXVI, fig. 151.

Monoecious; oogonium 1, obovoid-globose, operculate, division superior; oospore globose or subglobose, completing oogonium (or nearly so), spore wall smooth; antheridium 1-2, subepigynous, hypogynous, or scattered; sperms 2, division horizontal; basal cell elongate; terminal cell broadly apically acute; vegetative cell $16-20 \times 25-50 \mu$; oogonium $39-45 \times 45-51 \mu$; oospore $37-42 \times 37-44 \mu$; antheridium $15-18 \times 9-10 \mu$.

United States: Pennsylvania; Finland, Italy, Sweden, Latvia.

Cf. *Oe. crispum* (No. 109).

108. *Oe. obesum* (Wittrock) Hirn.

(Pl. XXXVII, fig. 347.)

1900, p. 166, Pl. XXVI, fig. 148; *Oe. pyrulum* Wittr. var. *obesum* Wittr. 1876, p. 44, Pl. 13, fig. 20.

Monoecious; oogonium 1, obovoid-globose, operculate, division superior; oospore globose, not quite filling oogonium, spore wall smooth, often thickened; antheridium 1-2, subepigynous or more rarely sub-

hypogynous; sperms 2, division horizontal; vegetative cell $12-15 \times 35-75\mu$; oogonium $40-43 \times 38-44\mu$; oospore $33-35 \times 33-35\mu$; antheridium $11-14 \times 10-15\mu$.

United States: Massachusetts; Austria, Bulgaria, Finland, France; Ceylon; Brazil.

Cf. *Oe. crispum* (No. 109).

109. *Oe. crispum* (Hassall) Wittrock.

(Pl. XXXV, fig. 319.)

Vesiculifera crispa Hass. 1845, p. 203, Pl. 52, fig. 8; *Oe. crispum* Wittr. 1874, p. 10; 1876, p. 45; *Oe. rostellatum* Pringsh. 1858, p. 69, Pl. 5, fig. 1; (?) *Vesiculifera vernalis* Hass. 1843, p. 434; (?) *V. candollei* Hass. 1845, p. 208, Pl. 52, fig. 9; *Oe. vernale* Wittr. 1874, p. 10; *Oe. crispum* (Hass.) Wittr. var. *elongatum* Wittr. 1876, p. 45; *Oe. hispanicum* Lewin 1888, p. 16, Pl. 2, figs. 40-42; Hirn 1900, p. 159, Pl. XXV, figs. 138 and 139; *Oe. crispum* (Hass.) Wittr. f. *vernale* (Hass.; Wittr.) Hirn 1900, p. 161, Pl. XXV, fig. 141; Heering 1914, p. 214, fig. 334; Tiffany 1926, p. 101, Pl. III, fig. 29.

Monoecious; oogonium usually 1, obovoid-globose, operculate, division superior; oospore globose or subglobose, quite filling the oogonium, spore wall smooth; antheridium 1-5, subepigynous or hypogynous; sperms 2, division horizontal; basal cell elongate, terminal cell apically obtuse; vegetative cell (10-) $12-16 \times 35-80\mu$; oogonium $37-45 \times 41-53\mu$; oospore $35-43 \times 37-43\mu$; antheridium $8-14 \times 7-12\mu$.

United States: Colorado, Pennsylvania, California, Michigan, Mississippi, Illinois, Ohio, Iowa, Alaska; Austria, England, Denmark, Finland, France, Germany, Italy, Spain, Sweden; Africa; Asia; Australia; Bolivia; Patagonia; Greenland; Vancouver, British Columbia.

109a. Form *granulosum* (Nordstedt) Hirn.

(Pl. XXXV, fig. 322.)

1900, p. 162, Pl. XXV, fig. 142; *Oe. crispum* (Hass.) Wittr. var. *granulosum* Nordst. 1877, p. 24; De Toni 1889, p. 37.

Oogonium subglobose, filled by oospore; antheridium subepigynous; spore wall of oospore punctate-granulate; vegetative cell $14-17 \times 35-63\mu$; oogonium $40-43 \times 42-45\mu$; oospore $37-40 \times 37-40\mu$; antheridium $11-13 \times 8-11\mu$.

Sweden.

109b. Form *inflatum* Hirn.

(Pl. XXXV, fig. 323.)

1900, p. 161, Pl. XXV, fig. 140; *Oe. vesicatum* Link var. *flavescens* Kuetz. in Rabenh. Alg. Sachs. No. 271, 1853; *Oe. rostellatum* Pringsh. in *ibid.* No. 2275, 1872.

Oogonium obovoid-globose; vegetative cell $12-16 \times 35-95\mu$; oogonium $40-50 \times 45-53\mu$; oospore $37-45 \times 38-45\mu$; antheridium $8-12 \times 9-12\mu$.

United States: Michigan, Florida, Mississippi; Switzerland, Sweden, Germany; South Africa.

109c. Var. *gracilescens* Wittrock.

(Pl. XXXV, fig. 321.)

In Wittr. and Nordst. Alg. Exs. No. 509, 1883; *Oe. Lagerheimii* Wittr. in *ibid.* No. 1018, 1893; Hirn 1900, p. 162, Pl. XXV, fig. 143; P. B. A. No. 518.

More slender than the type; oogonium varying from obovoid-globose to subellipsoid; oospore globose to ovoid; vegetative cell $10-14 \times 30-70\mu$; oogonium $33-39 \times 33-51\mu$; oospore $32-37 \times 33-42\mu$; antheridium $9-10 \times 7-9\mu$.

United States: Massachusetts, Iowa, Missouri, Ohio, Illinois, Mississippi, Louisiana, Minnesota; Austria; Brazil; Morocco, Angola.

109d. Var. *hawaiense* Nordstedt.

(Pl. XXXV, fig. 320.)

1878, p. 20, Pl. 2, figs. 9 and 10; Hirn 1900, p. 165, Pl. XXVI, fig. 147.

Oogonium subobovoid-globose to pyriform-globose, not filled by oospore; terminal cell mucronate or shortly setigerous; vegetative cell $10-16 \times 20-64\mu$; oogonium $30-38 \times 31-38\mu$; oospore $27-32 \times 27-33\mu$; antheridium $8-11 \times 6-8\mu$.

Hawaii.

109e. Var. *uruguayense* Magnus and Wille.

(Pl. XXXV, fig. 324.)

In Wille 1884, p. 39, Pl. 2, fig. 63; *Oe. crispum* (Hass.) Wittr. f. *gracilis* in Wittr. and Nordst. Alg. Exs. No. 704, 1886; Hirn 1900, p. 164, Pl. XXVI, fig. 145; *Oe. crispum* (Hass.) Wittr. var. *uruguayense* M. & W. f. *proprium* Hirn 1900, p. 164, Pl. XXVI, fig. 146.

Oogonium subobovoid-globose, usually filled by oospore; antheridium subepigynous or hypogynous or scattered; vegetative cell $10-14 \times 16-51\mu$; oogonium $30-38 \times 30-45\mu$; oospore $27-35 \times 27-37\mu$; antheridium $8-13 \times 6-12\mu$.

United States: Pennsylvania; Argentina, Brazil.

Oe. crispum is perhaps the commonest monoecious member of the genus among the operculate forms. Its vegetative cells are more slender than those of *Oe. autumnale* (No. 107) and there is less variation in the relative size of its oogonium and oospore than in *Oe. obesum* (No. 108). The forms and varieties of the species are separable from one another and the type largely on the basis of relative shapes and sizes of vegetative and reproductive cells. (See key).

110. *Oe. rupestre* Hirn.

(Pl. XXXVI, fig. 345.)

1900, p. 168, Pl. XXVI, fig. 152.

Monoecious; oogonium 1, suboboviform-globose, operculate, division superior; oospore globose, not filling oogonium, spore wall smooth; antheridium 1-3, subepigynous or hypogynous; sperms 2, division

horizontal; vegetative cell $20-27 \times 26-67\mu$; oogonium $48-58 \times 48-60\mu$; oospore $43-50$ (-54) $\times 43-50$ (-54) μ ; antheridium $18-22 \times 9-11\mu$.

Austria, France, Ireland, Latvia.

110a. Form **pseudautumnale** Hirn.

(Pl. XXXVI, fig. 346.)

1900, p. 169, Pl. XXVII, fig. 153.

A little smaller; vegetative cell $20-25 \times 30-65\mu$; oogonium $44-49 \times 45-55\mu$; oospore $43-47 \times 43-47\mu$; antheridium $19-22 \times 10-14\mu$.

United States: Mississippi; France, Germany.

Oe. rupestre has larger vegetative cells than any of the other monoecious operculate species with obovoid-globose to subglobose oogonia. It is to be compared with *Oe. crispum* (No. 109). *F. pseudautumnale* has smaller fruiting cells than the type.

111. **Oe. ahlstrandii** Wittrock.

(Pl. XXXVIII, fig. 358.)

In Witt. and Nordst. Exs. No. 401, 1882; Hirn 1900, p. 183, Pl. XXIX, fig. 179; Collins 1909, p. 247.

Monoecious; oogonium 1, ellipsoid, operculate, division superior; oospore ellipsoid, filling oogonium, wall smooth; antheridium 1-2, hypogynous; sperms 2, division horizontal; terminal cell apically obtuse; vegetative cell $10-18 \times 30-180\mu$; oogonium $35-42 \times 57-69\mu$; oospore $34-41 \times 53-62\mu$; antheridium $13-17 \times 9-12\mu$.

United States: New York; Sweden, England.

The large oogonium of this species separates it from all other ellipsoid, operculate, monoecious forms, except the still larger *Oe. pachydermum*, below.

112. **Oe. pachydermum** Wittrock and Lundell.

(Pl. XXXIX, figs. 377 and 378.)

In Witt. 1870, p. 125; Hirn 1900, p. 188, Pl. XXX, fig. 187.

Monoecious; oogonium 1 (rarely 2), ellipsoid, operculate, division superior, wall frequently thickened; oospore ellipsoid, not filling oogonium, spore wall smooth; antheridium 1-3, hypogynous or subepigynous, often terminal; sperms 2, division horizontal; basal cell elongate; terminal cell apiculate; vegetative cell $21-27 \times 34-120\mu$; oogonium $50-70 \times 75-100\mu$; oospore $40-60 \times 50-80\mu$; antheridium $18-21 \times 10-12\mu$.

Finland, Sweden.

Cf. *Oe. ahlstrandii* (No. 111).

113. *Oe. kirchneri* Wittrock.

(Pl. XXXVIII, fig. 371.)

1882, p. 104; *Oe. alternans* Kirch. 1878, p. 53; Hirn 1900, p. 183, Pl. XXIX, fig. 178.

Monoecious; oogonium 1, ellipsoid or obovoid-ellipsoid, operculate, division superior; oospore of same form as oogonium, nearly filling oogonium, spore wall smooth; antheridium 1-2, hypogynous and alternating with oogonium; sperms 2, division horizontal; terminal cell obtuse; vegetative cell $8-15 \times 16-60\mu$; oogonium $20-24 \times 34-48\mu$; oospore $19-23 \times 32-41\mu$; antheridium $9-13 \times 4-7\mu$.

Germany.

Chiefly characterized among those with ellipsoid, operculate oogonia by the alternation of oogonium and antheridium.

114. *Oe. gracillimum* Wittrock and Lundell.

(Pl. XXXVIII, fig. 373.)

In Wittr. 1874, p. 15; Wolle 1887, p. 74, Pl. 75, fig. 2; Hirn 1900, p. 184, Pl. XXIX, fig. 180; Collins 1909, p. 247; Heering 1914, p. 218, fig. 318; Tiffany 1926, p. 100, Pl. I, fig. 9.

Monoecious; oogonium 1, oblong, operculate, division superior; oospore oblong-ellipsoid, not filling oogonium, spore wall smooth; antheridium 1; sperms 2, division horizontal; basal cell elongate; vegetative cell $4-7 \times 16-42\mu$; oogonium $14-19 \times 34-40\mu$; oospore $13-17 \times 24-32\mu$; antheridium $3-5 \times 4-7\mu$.

United States: Iowa, Pennsylvania, Michigan, Illinois, Ohio, Mississippi, Kentucky; Sweden; Angola.

114a. Form *majus* West and West.

1897, p. 4; Hirn 1900, p. 184.

Vegetative cell $6-7 \times 20-40\mu$; oogonium $20-23 \times 32-35\mu$; oospore $17 \times 24\mu$.

Africa.

Oe. gracillimum almost invariably has some oogonia with imperfectly developed oospores.

115. *Oe. oblongellum* Kirchner.

(Pl. XXXVIII, fig. 372.)

In Hirn 1900, p. 182, Pl. XXIX, fig. 177.

Monoecious; oogonium 1, ellipsoid-obovoid, operculate, division superior; oospore ellipsoid, nearly filling oogonium, spore wall smooth; antheridium 1-2, hypogynous or scattered; sperms 2, division horizontal; terminal cell obtuse; vegetative cell $7-9 \times 14-35\mu$; oogonium $20-24 \times 25-33\mu$; oospore $18-22 \times 24-30\mu$; antheridium $6-8 \times 4-12\mu$.

United States: Illinois, Mississippi; Germany; Burma.

Oe. oblongellum is near *Oe. oblongum*, below, but the latter almost invariably has oogonia not filled lengthwise by oospores; the former has oogonia nearly completely filled.

116. *Oe. oblongum* Wittrock.

(Pl. XXXVIII, fig. 367.)

1872, p. 2; 1874, p. 15; Hirn 1900, p. 185, Pl. XXIX, fig. 181; Heering 1914, p. 218, fig. 319; Collins 1918, p. 66; Tiffany 1926, p. 101, Pl. III, fig. 28.

Monoecious; oogonium 1, oblong, operculate, division superior; oospore ellipsoid to ovoid, not filling oogonium longitudinally, spore wall smooth; antheridium 1-3, sperms 2, division horizontal; basal cell elongate; vegetative cell $6-11 \times 20-86\mu$; oogonium $20-26 \times 41-60\mu$; oospore $19-23 \times 30-36\mu$; antheridium $6-9 \times 7-9\mu$.

United States: Iowa, Illinois, Mississippi; Siberia; New Caledonia; India; Austria, Denmark, Finland, Germany, Russia, Norway, Latvia, Sweden.

116a. Form *majus* (Nordstedt) Hirn.

(Pl. XXXVIII, fig. 370.)

1900, p. 185, Pl. XXIX, fig. 182; *Oe. oblongum* Wittr. var. *majus* Nordst. in Wittr. 1876, p. 45, Pl. 13, fig. 21.

Vegetative cell $8-11 \times 35-75\mu$; oogonium $26-28 \times 42-50\mu$; oospore $22-26 \times 31-34\mu$; antheridium $8-9 \times 9-10\mu$.

United States: Michigan; Finland, Italy, Sweden; Morocco.

116b. Form *sphaericum* (Hallas) Hirn.

(Pl. XXXVIII, figs. 368 and 369.)

1906, p. 18, Pl. III, fig. 12; *Oe. sphaericum* Hallas 1905, p. 406, fig. 13.

Oospore globose; vegetative cell $6-11 \times 20-85\mu$; oogonium $21-28 \times 30-52\mu$; oospore $16-26 \times 16-26\mu$; antheridium $8 \times 6\mu$; basal cell $9 \times 72\mu$.

Denmark.

Cf. *Oe. oblongellum* (No. 115). The f. *sphaericum* is easily separated from the species by its globose oospore; f. *majus* has slightly larger oogonia.

117. *Oe. itzigsohnii* De Bary.

(Pl. XXXIX, figs. 379 and 380.)

1854, p. 56, Pl. 3, figs. 29-32; Wittr. 1870, p. 123; *Oe. platygynum* Wittr. in Eichler 1893, p. 2, Pl. 9, fig. 2; Hirn 1900, p. 177, Pl. XXVIII, fig. 167.

Monoecious; oogonium 1, ellipsoid, with median, conically obtuse projections, appearing stellate in vertical view, with 7-10 radiations, operculate, division inframedian; oospore globose, not completing oogonium, spore wall smooth; antheridium 1-2; terminal cell apically obtuse or apiculate; vegetative cell $8-10 \times 25-60\mu$; oogonium $34-40 \times 32-40\mu$; oospore $20-23 \times 20-23\mu$; antheridium $8-9 \times 8-15\mu$.

United States: Massachusetts; Germany, Poland, Sweden, Austria, England, Bulgaria, Denmark, Latvia, Russia, France.

117a. Form *heteromorphum* Hirn.

(Pl. XXXIX, fig. 381.)

1900, p. 178, Pl. XXVIII, fig. 168.

Processes on oogonium intermediate between conically obtuse projections and mammiform protuberances; vegetative cell $6-9 \times 30-110\mu$; oogonium $33-40 \times 35-40\mu$; oospore $20-25 \times 20-25\mu$.

Germany, Sweden.

117b. Var. *minus* West.

(Pl. XXXIX, fig. 382.)

1893, p. 97; 1897, p. 4; Hirn 1900, p. 178, Pl. XXVIII, fig. 169.

Smaller; processes of oogonium acutely rounded; vegetative cell $5-7 \times 25-70\mu$; oogonium $28-30 \times 28-30\mu$; oospore $16-20 \times 16-20\mu$.

England.

Oe. itzigsohnii is near *Oe. mammiferum* below, but differs in having conically obtuse projections, instead of mammiform projections, on the oogonia. Var. *minus* is slightly smaller than the type and f. *heteromorphum* has oogonial projections somewhat intermediate between the species proper and *Oe. mammiferum*.

118. *Oe. mammiferum* Wittrock; Nordstedt.

(Pl. XXXIX, figs. 383-385.)

Witr. 1874, p. 16; Nordst. 1877, p. 25, Pl. 3, figs. 4-6; *Oe. huillense* West 1897, p. 5, Pl. 365, figs. 7 and 8; Hirn 1900, p. 175, Pl. XXVIII, figs. 165 and 166.

Monoeocious; oogonium 1 (rarely 2-3), ellipsoid, with mammiform projections appearing as 7-9 radiations seen vertically, operculate, division inframedian, very narrow; oospore globose or subdepressed-globose (rarely depressed-globose), not filling oogonium, spore wall smooth; antheridium 1, subepigynous; basal cell elongate; vegetative cell $5-8 \times 20-56\mu$; oogonium $20-28 \times 20-30\mu$; oospore $12-17$ (-20) $\times 12-16\mu$; antheridium $6 \times 7\mu$.

United States: Massachusetts; Norway, Sweden; India; Africa.

Cf. *Oe. itzigsohnii* above.

119. *Oe. undulatum* (Brebisson) Al. Braun; Wittrock.

(Pl. XLII, fig. 407.)

Conferva undulata Breb.; *Oe. undulatum* Al. Braun in De Bary 1854, p. 94; Witr. 1870, p. 130; *Cymalonema confervaceum* Kuetz. 1849, p. 375 and 1853, p. 15, Pl. 47, fig. 1; Borge 1899, p. 4, Pl. 1, fig. 2; *Oe. undulatum* A. Br. var. *interrupte-incisum* Schroeder 1897, p. 10, Pl. 2, fig. 1; *Oe. sp.* Moebius 1894, p. 320, Pl. 1, fig. 21; *Oe. undulatum* A. Br. var. *Moebiusii* Schmidle 1896, p. 297, Pl. 9, fig. 1; *Oe. undulatum* (Breb.) Al. Br. var. *Moebiusii* Schmidle in Borge 1896, p. 5; Hirn 1900, p. 257, Pl. XLV, figs. 272-275.

Dioecious, nannandrous, gynandrosporous or idioandrosporous; oogonium 1-2, subglobose or ellipsoid-globose, operculate, division inferior, wide; oospore globose or subglobose, quite filling oogonium,

spore wall smooth, sometimes thick; androsporangium to 7-seriate; vegetative cell undulate (4 undulate constrictions); basal cell elongate, not undulate; terminal cell (sometimes oogonium) apically obtuse; dwarf male elongate-obconic, usually on suffultory cell, rarely on other vegetative cell near oogonium, antheridium interior; vegetative cell $15-22 \times 45-110\mu$; oogonium $48-56 \times 50-75\mu$; oospore $42-50 \times 42-52 (-60)\mu$; androsporangium $15-21 \times 7-14\mu$; dwarf male $8-10 \times 48-65 (-70)\mu$.

United States: Michigan, Florida, Indiana; Uruguay; Australia; Manchuria, China; Austria, England, Finland, France, Germany, Norway, Poland, Sweden, Russia, Latvia; Columbia; Africa (Victoria).

119a. Var. **americanum** Transeau.

(Pl. XLII, fig. 408.)

1917, p. 232.

Vegetative cell as in type; oogonium much larger; vegetative cell $15-22 \times 45-110\mu$; oogonium $58-68 \times 60-80\mu$; oospore $48-60 \times 48-56\mu$.

United States: Michigan.

119b. Form **senegalense** (Nordstedt) Hirn.

(Pl. XLII, fig. 409.)

1900, p. 261, Pl. XLV, figs. 276 and 277; *Oe. undulatum* (Breb.) A. Br. var. *senegalense* Nordst. 1880, p. 13; P. B. A. No. 73.

Oogonium and oospore smaller than the type; dwarf male shorter; oogonium to 5-seriate; vegetative cell with three median swellings repand; the terminal swelling entire, $15-22 \times 45-110\mu$; oogonium $42-52 \times 44-63\mu$; oospore $37-44 \times 36-44\mu$; androsporangium $15-19 \times 9-18\mu$; dwarf male $6-8 \times 37-44\mu$.

United States: Massachusetts; Africa: Senegal.

The undulate vegetative cells are characteristic. It bears some relation in the vegetative state to *Oe. nodulosum* (No. 96). The var. *americanum* has considerably larger oogonia than the type and f. *senegalense* has slightly smaller fruiting cells. A number of variations of this species are on record in algological literature, but the two above seem the only really tenable ones outside the species proper.

120. **Oe. nebraskense** Ohashi.

(Pl. XLII, figs. 410-413.)

Oe. nebraskensis Ohashi 1926, p. 213, figs. 1-20.

Dioecious, nannandrous, gynandrosporous; oogonium 1-4, flask-shaped or obovoid, pore supramedian; oospore subspherical or ovoid, not quite filling oogonium, outer layer of spore wall smooth, middle layer punctate, inner layer smooth; androsporangium 1-4, basal cell elongate; terminal cell attenuate-obtuse; dwarf male curved, on suffultory cell,

antheridium exterior, 1-4; vegetative cell slightly capitellate; filament 20-41 (-50) celled; vegetative cell, terminal $9-12 \times 170-305\mu$, median $20-27 \times 57-160\mu$, basal $22-35 \times 79-230\mu$; suffultory cell $41-58 \times 93-130\mu$; oogonium $60-67 \times 70-79\mu$; oospore $53-64 \times 60-76\mu$; androsporangium $20-25 \times 22-40\mu$; dwarf male stipe $12-19 \times 47-68\mu$; antheridium $9-13 \times 17-20\mu$.

United States: Nebraska (near Lincoln).

The only nannandrous species combining the characters of supramedian pore and punctate oospore.

121. *Oe. aster* Wittrock.

(Pl. XLIII, fig. 416.)

1872, p. 4; 1874, p. 29; De Toni 1889, p. 62; Hirn 1900, p. 198, Pl. XXXII, fig. 202; Fremy and Meslin 1926, p. 124, fig. 9.

Dioecious, nannandrous; oogonium 1, subdepressed-globose, pore median; oospore subdepressed-globose or subglobose, not completely filling oogonium; outer layer of spore wall echinate; suffultory cell sometimes a little swollen; terminal cell obtuse; dwarf male a little curved, on suffultory cell, stipe sometimes 2-celled, antheridium exterior, 1-2; vegetative cell $7-13 \times 50-110\mu$; oogonium $33-35 \times 34-39\mu$; oospore (with spines) $31-32 \times 25-30\mu$; dwarf male stipe $6-7 \times 20-25\mu$; antheridium $5-6 \times 7-8\mu$.

United States: Illinois; Sweden, Germany, Latvia, France; Ceylon.

Oe. aster is near *Oe. pugens* (No. 125) but the latter is larger in all dimensions.

122. *Oe. hystrix* Transeau and Tiffany.

(Pl. XLIII, fig. 414.)

1919, p. 240, Pl. XIV, figs. f-i; Tiffany 1926, p. 102, Pl. VIII, figs. 85 and 86.

Dioecious, nannandrous, idioandrosporous; oogonium 1, globose or somewhat obovoid, pore median; oospore globose to subglobose, nearly filling the oogonium, outer spore wall densely covered with spines; suffultory cells swollen; androsporangium 3-6; terminal cell obtuse; basal cell elongated; dwarf male nearly straight, resting on the suffultory cell, antheridium exterior, 1; female vegetative cell $8-15 \times 42-100\mu$, male $6-9 \times 50-67\mu$; suffultory cell $16-19 \times 42-70\mu$; oogonium $30-40 \times 36-53\mu$; oospore (including spines) $23-38 \times 28-43\mu$; androsporangium $6-8 \times 8-15\mu$; dwarf male stipe $6-10 \times 20-32\mu$; antheridium $5-6 \times 6-10\mu$.

United States: Illinois, Iowa.

This species is near *Oe. hystrix*, below, but differs in its smaller size and in its swollen suffultory cells.

123. *Oe. hystrix* Wittrock.

(Pl. XLIII, fig. 417.)

1870, p. 133; Wolle 1887, p. 87, Pl. 84, fig. 8; Hirn 1900, p. 213, Pl. XXV, fig. 218.

Dioecious, nannandrous, gynandrosporous (or possibly idioandrosporous); oogonium 1, ellipsoid, pore median; oospore ellipsoid,

nearly filling oogonium, outer spore wall echinate; androsporangium 1-3; terminal cell obtuse; dwarf male slightly curved, on suffultory cell; antheridium exterior, 1; vegetative cell $17-28 \times 30-120\mu$; oogonium $38-48 \times 45-65\mu$; oospore $37-46 \times 43-55\mu$; androsporangium $17-18 \times 13-18\mu$; dwarf male stipe $10-11 \times 22-25\mu$; antheridium $6-8 \times 9-14\mu$.

United States: Pennsylvania; Finland, Germany, Sweden.

123a. Var. **canadense** Tiffany.

(Pl. XLIII, fig. 418.)

1929, p. 74; *Oe. hystrix* Wittr. in Skuja 1927, p. 99, Pl. II, fig. 13.

Pore inframedian; suffultory cell enlarged; vegetative cell $14-25 \times 80-125\mu$; suffultory cell $24-30 \times 64-72\mu$; oogonium $40-53 \times 52-64\mu$; oospore $38-48 \times 40-60\mu$; androsporangium $16-20 \times 15-30\mu$; antheridium $7-8 \times 8-14\mu$; dwarf male stipe $9-12 \times 28-40\mu$.

British Columbia; Latvia.

123b. Var. **subglobosum** Wittrock.

(Pl. XLIII, fig. 419.)

1872a, p. 22, Pl. 1, fig. 7; Hirn 1900, p. 214, Pl. XXXV, fig. 219.

Smaller; oogonium and oospore subglobose; vegetative cell $17-21 \times 35-93\mu$; oogonium $36-38 \times 41-43\mu$; oospore $34-36 \times 39-40\mu$; androsporangium $16-18 \times 9-14\mu$; dwarf male stipe $10 \times 23\mu$; antheridium $7 \times 7\mu$.

Sweden.

The species is to be compared with *Oe. hystricinum* (No. 122). The variety *canadense* differs from the type in its inframedian pore and slightly swollen suffultory cell; var. *subglobosum* is smaller with subglobose oogonium.

124. **Oe. echinospermum** Al. Braun.

(Pl. XLIII, figs. 420 and 421.)

In Kuetzing 1849, p. 366; Kuetzing 1853, p. 12, Pl. 36, fig. 2; Wittr. 1870, p. 129; Hirn 1900, p. 199, Pl. XXXIII, fig. 204; P. B. A. No. 1673.

Dioecious, nannandrous, gynandrosporous or idioandrosporous; oogonium 1, ellipsoid-globose or subglobose, pore median; oospore globose, quite filling oogonium, outer layer of spore wall echinate; androsporangium 1-5; dwarf male a little curved, on suffultory cell, antheridium exterior, 1-2; vegetative cell $18-30 \times 45-130\mu$; oogonium $39-50 \times 41-57\mu$; oospore (with spines) $38-47 \times 38-49\mu$; androsporangium $21-25 \times 9-15\mu$; dwarf male stipe $10-15 \times 26-35\mu$; antheridium $6-12 \times 6-15\mu$.

United States: Illinois, Ohio, Alabama, Mississippi, Massachusetts, New York, New Jersey, Pennsylvania, Michigan; Austria, Scotland, England, Germany, Latvia, Finland, France, Sweden; Australia; British Columbia.

124a. Var. *horridum* Hirn.

(Pl. XLIII, fig. 422.)

1900, p. 201, Pl. XXXIII, fig. 205; *Oe. echinospermum* A. Br. var. Wolle 1887, p. 86, Pl. 85, figs. 6-9.

Larger with longer and more dense spines; antheridium pluricellular.

United States: Florida.

The species is one of the most easily recognized among the nannandrous forms by its echinate oospore, median pore, and large dimensions (as compared with *Oe. hystrix*, *Oe. aster*, and *Oe. pungens*). The variety *horridum* has no record other than that of Wolle, and its retention in the genus is open to considerable question.

125. *Oe. pungens* Hirn.

(Pl. XLIV, figs. 423 and 424.)

1900, p. 199, Pl. XXXII, fig. 203.

Dioecious, nannandrous, gynandrosporous; oogonium 1, subdepressed-globose (or nearly subglobose), pore median or a little above median; oospore subglobose, almost filling oogonium, outer spore wall echinate; androsporangium 1-?, hypogynous; dwarf male a little curved, on suffultory cell, antheridium exterior, 1-2; vegetative cell $12-16 \times 50-94\mu$; oogonium $40-48 \times 40-50\mu$; oospore (with spines) $37-44 \times 35-43\mu$; androsporangium $14-16 \times 10-15\mu$; dwarf male stipe $9-12 \times 20-30\mu$; antheridium $6-7 \times 8-12\mu$.

United States: South Carolina, Illinois.

Cf. *Oe. aster* (No. 121).

126. *Oe. exospirale* Tiffany.

(Pl. XLIV, figs. 425 and 426.)

1924, p. 184, Pl. II, figs. 1-3; 1926, p. 102, Pl. IX, figs. 100-102.

Dioecious, nannandrous, idioandrosporous; oogonium 1 (rarely 2), subglobose or obovoid-globose (rarely subhexagonal), pore median; oospore globose or subglobose, nearly filling the oogonium, spore wall of two layers: outer layer marked by 5-8 spiral ribs uniting at the poles, the polar axis always placed in a transverse position, never parallel with the filament, the inner layer smooth; suffultory cell swollen; androsporangium 1-6; basal cell elongate; dwarf male a little curved, situated on the suffultory cell, with exterior antheridium 1-3; vegetative cell $8-12 \times 60-88\mu$; suffultory cell $16-28 \times 60-88\mu$; oogonium $40-44 \times 44-48\mu$; oospore $32-38 \times 38-40\mu$; androsporangium $12-16 \times 14-20\mu$; dwarf male stipe $14-16 \times 30-40\mu$; antheridium $8-12 \times 14-16\mu$.

United States: Iowa, Ohio, Mississippi.

This species should be compared with *Oe. spirale* (No. 128) and *Oe. illinoisense* (No. 127). It differs from the former in having smaller oogonia and oospores and in its tumid suffultory

cells; it is separated from the latter on the basis of smaller dimensions.

127. *Oe. illinoisense* Transeau.

(Pl. XLIV, figs. 431-433.)

1914, p. 299, Pl. XXIX, figs. 6-8.

Dioecious, nannandrous, gynandrosporous; oogonium 1, subglobose to obovoid-globose, pore median; oospore globose or subglobose, nearly filling oogonium, outer spore wall with 4-7 spiral ribs, uniting at poles, polar axis always transverse to axis of filament; suffultory cell enlarged; androsporangium 1-5; basal cell elongate; dwarf male slightly curved, on suffultory cell; antheridium exterior, 1-4; vegetative cell $13-18 \times 80-140\mu$; suffultory cell $32-40 \times 50-80\mu$; oogonium $51-60 \times 60-70\mu$; oospore $45-56 \times 48-66\mu$; androsporangium $13-17 \times 17-22\mu$; dwarf male stipe $14-17 \times 37-57\mu$; antheridium $9-12 \times 15-23\mu$.

United States: Illinois, Mississippi, Kentucky.

Cf. *Oe. exospirale*, above.

128. *Oe. spirale* Hirn.

(Pl. XLIV, figs. 427 and 428.)

1900, p. 201, Pl. XXXIII, fig. 206.

Dioecious, nannandrous, idioandrosporous; oogonium 1, subglobose or obovoid-globose, pore median; oospore globose or subglobose, not filling oogonium, spore wall double: outer layer with 4-7 spiral ribs, anastomosate, united at the poles, inner layer smooth; androsporangium 1-3; basal cell elongate; dwarf male a little curved, near oogonium, antheridium exterior, 1; vegetative cell $20-33 \times 40-130\mu$; oogonium $52-60 \times 52-60\mu$; oospore $46-56 \times 46-56\mu$; androsporangium $17-22 \times 11-14\mu$; dwarf male stipe $11-16 \times 41-59\mu$; antheridium $10-11 \times 18-20\mu$.

Asia (Java).

128a. Var. *acutum* West and West.

(Pl. XLIV, fig. 430.)

1902, p. 128, Pl. 17, figs. 3-5; Hirn 1906, p. 22, Pl. III, fig. 14.

Spiral ribs larger, irregularly and acutely dentate; vegetative cell $20-26 \times 70-106\mu$; oogonium $46-56 \times 54-60\mu$; oospore $37-45 \times 37-45\mu$; dwarf male stipe $9-12 \times 34-44\mu$; antheridium $8-9 \times 14-18\mu$.

United States: Mississippi; Ceylon, India.

128b. Var. *latviense* Tiffany.

Pl. XLIV, fig. 429.)

1929, p. 75; *Oe. spirale* Hirn f. in Skuja 1927, p. 102, Pl. II, fig. 14 a-b.

Pore supramedian or superior; vegetative cell $16-20 \times 65-120\mu$; oogonium $49-60 \times 65\mu$; oospore (without ribs) $35-45 \times 35-45\mu$, not filling oogonium, outer wall with 5-6 spirally arranged ribs; dwarf male $12 \times 50\mu$.

Latvia.

The species should be compared with *Oe. exospirale* (No. 126). The variety *acutum* has smaller oospores and larger ribs; var. *latviense* has a distinctly above-median pore which is sometimes nearly superior.

129. *Oe. depressum* Pringsheim.

(Pl. XLVI, fig. 446.)

1858, p. 69, Pl. 5, fig. 5; Hirn 1900, p. 249, Pl. XLII, fig. 261.

Dioecious, nannandrous, gynandrosporous; oogonium 1 (rarely 2), depressed globose, pore median; oospore depressed globose, not filling oogonium; androsporangium 2-celled; dwarf male oblong-obovoid, on oogonium, antheridium interior; vegetative cell $8-9 \times 25-54\mu$; oogonium $28 \times 26\mu$; oospore $23 \times 17-18\mu$; dwarf male $4-5 \times 14-16\mu$.

Germany; Greenland.

Cf. *Oe. semiapertum*, below.

130. *Oe. semiapertum* Nordstedt and Hirn.

(Pl. XLVI, fig. 445.)

In Hirn 1900, p. 250, Pl. XLII, fig. 262.

Dioecious, nannandrous, gynandrosporous; oogonium 1 (rarely 2), subhexagonal-globose to subpyriform-globose, pore median, rimiform, extending nearly half way round the oogonium; oospore globose or subglobose, not filling oogonium, spore wall smooth; androsporangium 1-3, hypogynous or subepigynous; terminal cell apically obtuse; dwarf male subobovoid, on oogonium, antheridium interior; vegetative cell $9-12 \times 38-105\mu$; oogonium $32-35 \times 33-40\mu$; oospore $25-29 \times 25-30\mu$; androsporangium $9-10 \times 10-14\mu$; dwarf male $8-9 \times 14-15\mu$.

French Guiana.

This species and *Oe. depressum*, above, are the smallest of nannandrous, medianly poriferous members of the genus having smooth and nearly globose oospores. *Oe. depressum* is separated by its smaller size and non-rimiform pore.

131. *Oe. gallicum* Hirn.

(Pl. XLVII, fig. 453.)

1900, p. 197, Pl. XXXII, fig. 201; *Oe. flavescens* Hass. in Roumeguere Alg. Exs. No. 1173.

Dioecious, nannandrous, (?) idioandrosporous; oogonium 1-2, subhexagonally globose, pore a little above median; oospore of same form as oogonium and quite filling it, spore wall smooth; dwarf male suberect, on suffultory cell (rarely oogonium), antheridium exterior, 1-2; vegetative cell $17-22 \times 51-120\mu$; oogonium $43-48 \times 44-54\mu$; oospore $41-46 \times 39-44\mu$; dwarf male stipe $11-16 \times 26-30\mu$; antheridium $8-9 \times 7-11\mu$.

France.

Cf. *Oe. braunii*, below.

132. *Oe. braunii* Kuetzing; Pringsheim.

(Pl. XLVII, fig. 458.)

Kuetzing 1849, p. 366; 1853, p. 12, Pl. 36, fig. 3; Pringsheim 1858, p. 70, Pl. 5, fig. 6; Wille 1887, p. 79, Pl. 79, figs. 6 and 7; *Oe. calcareum* Cleve var. *gaditanum* Lewin 1888, p. 18, Pl. 3, fig. 53; Hirn 1900, p. 194, Pl. XXXII, fig. 197; Collins 1909, p. 249; Heering 1914, p. 172; Tiffany 1926, p. 102.

Dioecious, nannandrous, gynandrosporous; oogonium 1, ovoid to subglobose, pore median; oospore globose, not filling oogonium, spore wall smooth; androsporangium 1-2; basal cell elongate; terminal cell apically obtuse; dwarf male usually on suffultory cell, stipe somewhat curved; antheridium exterior, 1; vegetative cells $13-15 \times 25-60\mu$; suffultory cell $16-20 \times 28-50\mu$; oogonium $30-37 \times 33-43\mu$; oospore $27-33 \times 27-33\mu$; androsporangium $13-15 \times 11-12\mu$; dwarf male stipe $7-12 \times 20-28\mu$; antheridium $5-8 \times 9-10\mu$.

United States: Iowa, Illinois, New Jersey, Pennsylvania, Michigan; Austria, England, Denmark, Finland, France, Germany, Spain, Ireland, Latvia, Norway, Sweden; Africa; British Columbia.

132a. Var. *hafniense* (Hallas) Hirn.

(Pl. XLVII, figs. 460-462.)

1906, p. 10, Pl. III, fig. 13; *Oe. hafniense* Hallas 1905, p. 398, fig. 1.

A little smaller; idioandrosporous; oospore globose or angular-globose; vegetative cell $12-15 \times 25-60\mu$; oogonium $23-30 \times 28-35\mu$; oospore $21-28 \times 21-28\mu$; dwarf male stipe $6-8 \times 12-21\mu$; antheridium $7 \times 7\mu$.

Denmark.

132b. Var. *zehneri* Tiffany.

(Pl. XLVII, fig. 459.)

1927, p. 204, Pl. IX, fig. 2.

Somewhat larger in all dimensions; oospore ovoid; vegetative cell $12-24 \times 34-72\mu$; suffultory cell $21-32 \times 48-52\mu$; oogonium $40-50 \times 48-60\mu$; oospore $34-44 \times 36-50\mu$; dwarf male stipe $8-10 \times 16-24\mu$; antheridium $7-8 \times 8-12\mu$.

United States: Indiana.

The species is near, but smaller than, *Oe. gallicum* (No. 131) and *Oe. flavescens*, below. Var. *hafniense* is smaller than the type and is idioandrosporous; var. *zehneri* is somewhat larger than the species and has ovoid oospores.

133. *Oe. flavescens* (Hassall) Wittrock).

(Pl. XLIX, fig. 471.)

(?) *Vesiculifera flavescens* Hass. 1845, p. 206, Pl. 53, fig. 9; *Oe. flavescens* Wittr. 1870, p. 127; *Oe. flavescens* (Hass.) Wittr. var. *gynandrosporum* Hirn 1895, p. 16; Hirn 1900, p. 196, Pl. XXXII, figs. 199 and 200; Collins 1909, p. 249; Heering 1914, p. 174, fig. 232; Tiffany 1926, p. 103, Pl. IX, fig. 95.

Dioecious, nannandrous, idioandrosporous or gynandrosporous; oogonium 1, ellipsoid-globose to subglobose, pore median; oospore

globose, not quite filling oogonium, spore wall smooth; androsporangium 1-9; dwarf male somewhat curved, on the suffultory cell; antheridium 1-2; vegetative cell $18-23 \times 72-140\mu$; oogonium $49-52 \times 51-60\mu$; oospore $45-49 \times 45-49\mu$; androsporangium $17-20 \times 8-18\mu$; dwarf male stipe $11-12 \times 36-45\mu$; antheridium $9-10 \times 15-20\mu$.

United States: Ohio, Mississippi, Alabama, Massachusetts, Minnesota, Iowa; England, Sweden, Finland.

Cf. *Oe. braunii* (No. 132).

134. *Oe. manschuricum* Skvortzow.
(Pl. XLVI, fig. 442.)

1926, p. 434, fig. 8.

Dioecious, nannandrous; oogonium 1, globose, pore supramedian; oospore globose or nearly so, not quite filling oogonium, outer spore wall echinate; dwarf male broadly egg-shaped, unicellular, on oogonium; vegetative cell $14-17 \times 56-85\mu$; oogonium $30-31 \times 37-38\mu$; oospore $26-28 \times 26-30\mu$; dwarf male $13-24 \times 20-37\mu$.

North Manchuria.

Characterized by its echinate oospores and supramedian pores. To be compared with *Oe. hystricinum* (No. 122) and *Oe. echinospermum* (No. 124).

135. *Oe. sexangulare* Cleve.
(Pl. XLVII, fig. 454.)

In Wittr. 1870, p. 131; Wolle 1887, p. 82, Pl. 79, figs. 8 and 9; *Oe. hexagonum* Kuetz. in Roumeguere Alg. Exs. No. 294 and *Oe. vesicalum* Link f. *fuscescens* Dupray in loc. cit. No. 687; Hirn 1900, p. 211, Pl. XXXV, fig. 216; P. B. A. No. 1679.

Dioecious, nannandrous, gynandrosporous; oogonium 1 (rarely 2), sexangular-ellipsoid, pore a little above median; oospore same form as oogonium and nearly filling it, spore wall smooth; suffultory cell sometimes slightly tumid; androsporangium 1-3; dwarf male on suffultory cell, a little curved, stipe of 2-3 cells, antheridium exterior, 1; vegetative cell $9-16 \times 30-110\mu$; oogonium $29-33 \times 33-39\mu$; oospore $27-31 \times 31-36\mu$; androsporangium $13-14 \times 10-14\mu$; dwarf male stipe $7-9 \times 21-30\mu$; antheridium $6-7 \times 9-12\mu$.

United States: Indiana, Mississippi, New York, Louisiana, Pennsylvania, Kentucky, Michigan; Austria, France, Norway, Sweden.

135a. Var. *angulosum* (Hallas) Hirn.
(Pl. XLVII, figs. 455 and 456.)

1906, p. 21, Pl. III, fig. 17; *Oe. angulosum* Hallas 1905, p. 402, fig. 7.

Smaller, oospore not filling oogonium; vegetative cell $10-14 \times 15-54\mu$; oogonium $19-30 \times 30-42\mu$; oospore $17-28 \times 21-30\mu$; dwarf male stipe $6-7 \times 18-21\mu$; antheridium $6 \times 7\mu$.

Denmark.

135b. Var. *majus* Wille.

(Pl. XLVII, fig. 457.)

1880, p. 68; Hirn 1900, p. 212, Pl. XXV, fig. 217.

Larger, pore median; vegetative cell $15-23 \times 30-68\mu$; oogonium $36-42 (-48) \times (36-) 41-45\mu$; oospore $34-40 \times 39-43\mu$; androsporangium $14-18 \times 8-10\mu$; dwarf male stipe $7-9 \times 18-30\mu$; antheridium $6-8 \times 6-10\mu$.

United States: Mississippi, Massachusetts; Austria, Finland, Germany, Norway, Sweden.

The species is readily identified among the nannandrous forms by its sexangular-ellipsoid oogonia and oospores. Var. *angulosum* is smaller and var. *majus* is larger than the species.

136. *Oe. stellatum* Wittrock.

(Pl. XLV, fig. 441.)

1870, p. 129; 1874, p. 27, Pl. I, fig. 15; (?) *Oe. spinospermum* Reinsch 1875, p. 81, Pl. 15, fig. 1; Hirn 1900, p. 205, Pl. XXXIV, fig. 219.

Dioecious, nannandrous, gynandrosporous; oogonium 1-3, obovoid-globose, pore superior; oospore globose, quite filling oogonium, outer spore wall with 4-7 somewhat spirally arranged ribs, appearing definitely dentate on the margin, sometimes anastomosate; androsporangium 1-3, usually subepigynous; basal cell elongate; terminal cell slender, subhyaline, apically obtuse; dwarf male suberect, on suffultory cell, antheridium 1-2; vegetative cell $15-35 \times 40-225\mu$; oogonium $51-64 \times 56-70\mu$; oospore (with ribs) $50-58 \times 50-58\mu$; androsporangium $14-19 \times 13-20\mu$; dwarf male stipe $11-13 \times 45-52\mu$; antheridium $6-9 \times 8-13\mu$.

United States: Alabama, Florida; Africa; Brazil; Australia; Finland, Germany, Italy, Sweden; Ceylon.

This alga is often placed with the species having spirally arranged ribs on the oospore. The ribs are really irregular rows of dentate spines, and I have thus separated it in the key.

137. *Oe. silvaticum* Hallas.

(Pl. XLIX, fig. 472.)

1905, p. 404; fig. 9; Hirn 1906, p. 21, Pl. III, fig. 16.

Dioecious, nannandrous, gynandrosporous; oogonium 1, ellipsoid-globose, pore superior; oospore globose, not filling oogonium; suffultory cell subtumid; dwarf male suberect, on suffultory cell, antheridium exterior; vegetative cell $12-15 \times 65-120\mu$; suffultory cell $20 \times 60\mu$; oogonium $37 \times 50\mu$; oospore $28 \times 28\mu$; androsporangium $7 \times 17\mu$; dwarf male stipe $11 \times 39\mu$; antheridium $7 \times 7\mu$.

Denmark.

In general appearance this species resembles *Oe. flavescens* (No. 133), but differs in its smaller size and superior pore. The suffultory cells are not nearly so much enlarged relatively as in *Oe. borisianum*, below.

138. *Oe. borisianum* (Le Clerc) Wittrock.

(Pl. XLVIII, fig. 469.)

1874, p. 25; *Prolifera borisiana* Le Clerc 1817, p. 475, Pl. 23, fig. 6; *Oe. borisianum* Wittr. 1870, p. 132; *Vesiculifera aurea* Hassall 1842, p. 392; *V. borissii* Hassall 1845, p. 201, Pl. 52, fig. 7; *Oe. apophysatum* Al. Braun in Kuetzing 1849, p. 366, and 1853, p. 12, Pl. 35, fig. 5; *Oe. seligerum* Vaupell 1859, p. 17, Pl. 1. (?) *Androgynia mirabilis* Wood 1872, p. 196, Pl. 18, fig. 2; *Oe. birmanicum* Wittrock 1874, p. 24; Hirn 1900, p. 217, Pl. XXXVI, fig. 223; Collins 1909, p. 255; Heering 1914, p. 178, fig. 243; Tiffany 1926, p. 103, Pl. VIII, fig. 87; P. B. A. No. 517.

Diocious, nannandrous, gynandrosporous or idioandrosporous; oogonium 1-5, obovoid or quadrangular-ellipsoid, pore superior; oospore ovoid to obovoid (often quadrangular-ovoid), not filling oogonium, spore wall smooth; androsporangium 1-7, usually in the upper part of the filament; basal cell elongate; terminal cell, often an oogonium, broadly apiculate, obtuse, or sometimes setiferous; dwarf male, somewhat curved, on suffultory cell; antheridium 2; vegetative cell $15-23 \times 45-140\mu$; suffultory cell $31-38 \times 50-92\mu$; oogonium (33-) $40-50 \times 55-90\mu$; oospore $35-46 \times 48-60\mu$; androsporangium $16-19 \times 15-23\mu$; dwarf male stipe $12-18 \times 35-47\mu$; antheridium $7-10 \times 11-16\mu$.

United States: Iowa, Massachusetts, Utah, Michigan, California, Illinois, Ohio, Pennsylvania; British Columbia; Austria, England, Denmark, Finland, France, Germany, Spain, Sweden, Latvia; Brazil, Columbia, Paraguay; Australia; India; South Africa.

138a. Var. *westii* Tiffany and Brown.

(Pl. XLVIII, fig. 470.)

In Tiffany, 1929, p. 74; *Oe. borisianum* (Le Cl.) Wittr. in West 1903, p. 36.

Larger dimensions throughout; vegetative cell $17-19 \times 130-180\mu$; suffultory cell $31-42 \times 105-168\mu$; oogonium $53-60 \times 67-88\mu$; oospore $49-58 \times 49-58\mu$; antheridium $17 \times 12-13\mu$.

United States: Michigan, Mississippi; England.

Oe. borisianum is one of the commonest nannandrous species. It is easily identified by the striking difference in size between the cells above the oogonia and the suffultory cells. *F. westii* is larger than the type.

139. *Oe. alternans* Wittrock and Lundell.

(Pl. XLVIII, fig. 463.)

In Wittr. 1874, p. 20; Hirn 1900, p. 251, Pl. XLII, fig. 263.

Diocious, nannandrous, gynandrosporous; oogonium 1, subglobose, often alternating with vegetative cell, pore superior; oospore globose, filling oogonium, wall smooth, often thick; suffultory cell often sub-tumid; androsporangium 1-2, subepigynous or epigynous; basal cell elongate; terminal cell obtuse; dwarf male oblong, with an inferior constriction, on suffultory cell; vegetative cell $21-35 \times 45-160\mu$; suffultory cell $26-40 \times 52-160\mu$; oogonium $65-80 \times 65-86\mu$; oospore $59-75 \times 59-75\mu$; androsporangium $18-25 \times 16-25\mu$; dwarf male $12-17 \times 38-46\mu$.

Sweden.

Readily characterized by its short barrel-shaped vegetative cells, often alternating with oogonia.

140. *Oe. kitutæ* G. S. West.

1907, p. 98.

Dioecious, nannandrous; oogonium 1-2, subglobose or obovate-globose, pore superior; oospore globose, not filling oogonium, spore wall smooth; terminal cell obtuse; dwarf male elongate, slightly curved, smaller below, situated on oogonium or suffultory cell; antheridium interior (?); vegetative cell $26 \times 130-180\mu$; oogonium $76-78 \times 75-80\mu$; oospore $72 \times 72\mu$; suffultory cell $44 \times 132\mu$; dwarf male $12-13 \times 58-66\mu$.

Africa.

West has left no figure for this species. It is to be compared with *Oe. borisianum* (No. 138) and *Oe. silvaticum* (No. 137).

141. *Oe. victoriense* G. S. West.

(Pl. XLIX, fig. 473.)

In Hardy 1906, p. 22; West 1909, p. 47, Pl. 6C.

Dioecious, nannandrous, gynandrosporous; oogonium 1, ellipsoid-globose to ellipsoid-doliiform, pore superior; oospore exactly globose, not filling oogonium, spore wall smooth; androsporangium 5-6 celled; dwarf male on suffultory cell, stipe curved, antheridia 2; suffultory cell broadly dilated; vegetative cell $25-29 \times 50-87\mu$; oogonium $49 \times 57\mu$; oospore $39 \times 39\mu$; androsporangium $25-26 \times 4-8\mu$; antheridium $7-8 \times 4-5\mu$.

Africa (Victoria).

Resembles *Oe. crassiusculum*, below, but has single oogonia, nearly exactly globose oospores, and smaller oogonia.

142. *Oe. crassiusculum* Wittrock.

(Pl. XLIX, fig. 474.)

1870, p. 132, 1874, p. 24; Cooke 1884, p. 162, Pl. 60, fig. 5; Hirn 1900, p. 215.

Dioecious, nannandrous, gynandrosporous; oogonium 1-2, globose-obovoid or subglobose, pore superior; oospore ellipsoid-globose or globose, spore wall smooth and thickened, quite filling oogonium; androsporangium 2-5; dwarf male suberect, on or near suffultory cell, antheridium exterior, 1-?; vegetative cell $27-30 \times 95-105\mu$; oogonium $54-60 \times 60-75\mu$; oospore $51-57 \times 52-63\mu$; androsporangium $26-28 \times 10-18\mu$; dwarf male $13 \times 60\mu$; antheridium $7-9 \times 9-16\mu$.

United States: Ohio, Mississippi, Connecticut; Australia; England, Sweden.

142a. Var. **arechavaletæ** (Wittrock) Hirn.

(Pl. L, fig. 478.)

1900, p. 216, Pl. XXXVI, fig. 222; *Oe. arechavaletæ* Wittr. in Wittr. and Nordst. Exs. 1886 and in De Toni 1889, p. 55.

(?) Idioandrosporous; vegetative cell $21-29 \times 32-110\mu$; oogonium $39-54 \times 42-65\mu$; oospore $37-52 \times 40-57\mu$; dwarf male $14-16 \times 40-57\mu$; antheridium $9-14 \times 10-16\mu$.

Uruguay.

142b. Var. **idioandrosporum** Nordstedt and Wittrock.

(Pl. XLIX, figs. 475 and 476.)

In Wittr. & Nordst. Exs. 1879; *Oe. huntii* Wood in Tilden Amer. Algae 1894; *Oe. crassiusculum* Wittr. in P. B. A. 1895; *Oe. crassipellitum* West 1899, p. 55, Pl. 394, figs. 3-5; Hirn 1900, p. 215, Pl. XXXV, fig. 220; Collins 1909, p. 255; Heering 1914, p. 178, fig. 242; Tiffany 1926, p. 103, Pl. VIII, fig. 89; P. B. A. Nos. 72 and 716.

Idioandrosporous; oogonium 1-3, globose-obovoid to globose, pore superior; oospore ellipsoid-globose, ovoid, angular-globose (rarely globose), nearly filling the oogonium, spore wall smooth, thick; androsporangium 1-4; dwarf male straight or slightly curved, on or near the suffultory cell; antheridium exterior, 1-2; vegetative cell $25-36 \times 65-200\mu$; oogonium $48-59 \times 57-90\mu$; oospore $42-57 \times 50-66\mu$; androsporangium $30-34 \times 12-21\mu$; dwarf male stipe $14-16 \times 60-70\mu$; antheridium $8-10 \times 10-18\mu$.

United States: Massachusetts, Maine, Michigan, Pennsylvania, New Jersey, Minnesota, Connecticut, South Dakota, Iowa, Ohio, New York, Illinois, Mississippi; England, Denmark, France, Germany, Sweden.

This variety is one of the commonest of the nannandrous species of the genus. In fact, in comparison to it, the type seems rare. Characterized by the thick-walled oospore, globose-obovoid oogonia, and relatively long and straight vegetative cells. It differs from the species largely in its idioandrosporous habit.

142c. Var. **cataractum** (Wolle) Tiffany nov. comb.

(Pl. XLIX, fig. 477.)

Oe. cataractum Wolle 1887, p. 77, Pl. 85, figs. 10-12; *ibid.*, Hirn 1900, p. 308, Pl. XLII, fig. 264; *ibid.*, Collins 1909, p. 266.

Oogonium 1-2, often terminal, obovoid-globose or subglobose or broadly ovate, pore superior; oospore globose or obovoid-globose, nearly filling oogonium; androsporangium 2-6; dwarf male curved, on suffultory cell or below; vegetative cell $28-38 \times 42-112\mu$; oogonium $55-60 \times 60-75\mu$; oospore $50-55 \times 50-60\mu$; androsporangium $26-30 \times 10-15\mu$; dwarf male $10 \times 65\mu$.

United States: Pennsylvania, Ohio.

From Wolle's description this plant is certainly very close to *Oe. crassiusculum*, as suggested by Hirn (1900). Wolle's

figure does not help much and his confusing statements regarding an interior and a unicellular antheridium probably mean that he did not attach much significance to this character. The Ohio material with short vegetative cells seemed close to *Oe. crassiusculum* in other characteristics. I have therefore placed Wille's species as a variety of *crassiusculum*, with an emended description based on the Ohio specimens.

143. *Oe. multisporum* Wood.

(Pl. XLVI, figs. 450 and 451.)

1869, p. 141; *Androgynia multispora* Wood 1872, p. 196, Pl. 17, fig. 3; Hirn 1900, p. 232, Pl. XXXIX, fig. 239; Collins 1909, p. 257; Fritsch and Rich 1913, p. 76, fig. 1 A; P. B. A. No. 1784.

Dioecious, nannandrous; oogonium 1-3, subovoid or subglobose, pore superior; oospore globose, nearly filling oogonium, spore wall smooth; dwarf male a little curved or nearly erect, near or on oogonium, antheridium exterior, 1-4; vegetative cell $10-15 \times 10-30\mu$; oogonium $24-35 \times 27-33\mu$; oospore $27-30 \times 24-30\mu$; dwarf male stipe $10-11 \times 26-30\mu$; antheridium $7-9 \times 7-9\mu$.

United States: Pennsylvania, Illinois; England.

143a. Var. *magnum* Ackley.

(Pl. XLVI, fig. 452.)

1929, p. 305, Pl. XXXVI, figs. 20-22.

Larger than the species; idioandrosporous; oogonium 1, subglobose, pore supramedian (rarely median); androsporangium 1-2; dwarf male on or near oogonium or scattered; vegetative cell $14-18 \times 30-90\mu$; suffultory cell $17-21 \times 28-90\mu$; oogonium $40-43 \times 33-38\mu$; oospore $38-41 \times 32-36\mu$; androsporangium $17-18 \times 16-22\mu$; dwarf male stipe $12-14 (-20) \times 24-26 (-29)\mu$; antheridium $8-10 (-12) \times 8-10\mu$.

United States: Michigan (Muskegon Lake).

It is sometimes difficult to decipher the pore in the species proper. The plant is usually irregularly formed with numerous and often scattered dwarf males and frequently gives the appearance of not being mature. The variety is larger.

144. *Oe. irregulare* Wittrock.

(Pl. XLVI, fig. 447.)

1870, p. 128; 1874, p. 22; Hirn 1900, p. 202, Pl. XXXIII, fig. 207.

Dioecious, nannandrous; oogonium 1, globose or rarely subdepressed-globose, pore superior; oospore globose, completing oogonium, spore wall smooth; dwarf male erect, on or near oogonium, antheridium exterior, 1-4; vegetative cell $15-20 \times 40-80\mu$; oogonium $37-45 \times 36-47\mu$; oospore $36-42 \times 34-41\mu$; dwarf male stipe $12-15 \times 20-24\mu$; antheridium $10-12 \times 6-8\mu$.

United States: Illinois, Ohio, Florida, Michigan; British Columbia; Denmark, Sweden.

144a. Var. **condensatum** (Hallas) Hirn.

(Pl. XLVI, figs. 448 and 449.)

1906, p. 17, Pl. III, fig. 15; *Oe. condensatum* Hallas 1905, p. 400, fig. 4.

Idioandrosporous; oogonium rarely in two's; oospore depressed-globose; androsporangium 1-13; vegetative cell $14-18 \times 20-80\mu$; oogonium $35-46 \times 28-38\mu$; oospore $32-44 \times 25-36\mu$; androsporangium $12-14 \times 7-21\mu$; dwarf male stipe $11-13 \times 21-24\mu$; antheridium $9-10 \times 7-12\mu$.

United States: Michigan; Denmark.

The species is characterized by erect dwarf males, globose oospores, and smooth spore walls. It is larger than *Oe. multisporum* (No. 143) and there is little tendency toward a cluster of dwarf males. The var. *condensatum* has depressed-globose oospores and the dwarf males sometimes appear more or less in clusters. In superficial appearance there is considerable resemblance between *Oe. multisporum* var. *magnum* (No. 143a) and *Oe. irregulare* var. *condensatum*.

145. **Oe. donnellii** Wolle; Hirn.

(Pl. XLV, fig. 437.)

Wolle 1880, p. 48; Wolle 1887, p. 85, Pl. 84, figs. 3-6; Hirn 1900, p. 206, Pl. XXXIV, fig. 211.

Dioecious, nannandrous, idioandrosporous; oogonium 1 (rarely 2), obovoid-globose, pore superior; oospore globose, not filling oogonium, outer layer of spore wall echinate, spirals 5-7, anastomosate; androsporangium 4-10; dwarf male a little curved, on suffultory cell or rarely on oogonium, antheridium exterior, 1-2-?; vegetative cell $41-59 \times 54-175\mu$; oogonium $63-78 \times 70-93\mu$; oospore (with spines) $60-70 \times 60-70\mu$; androsporangium $40-45 \times 10-12\mu$; dwarf male stipe $16-21 \times 63-74\mu$; antheridium $14-15 \times 8-22\mu$.

United States: Florida.

145a. Var. **wittrockianum** Hirn.

(Pl. XLV, figs. 438 and 439.)

1900, p. 207, Pl. XXXIV, fig. 212; *Oe. wittrockianum* Hirn 1896 in Wittr., Nordst. and Lagerh. Alg. Exs. No. 1251.

Smaller; spines a little smaller; vegetative cell $(32-)35-45 \times 70-135\mu$; oogonium $63-73 \times 68-75\mu$; oospore (with spines) $55-66 \times 56-68\mu$; androsporangium $30-38 \times 11-26\mu$; dwarf male stipe $11-15 \times 50-65\mu$; antheridium $9-10 \times 20-23\mu$.

Brazil, Paraguay.

The species is characterized by the echinate oospore wall, the spines being arranged spirally. The variety *wittrockianum* is very similar to the type, having slightly more slender filaments and smaller spines.

146. *Oe. armigerum* Hirn.

(Pl. XLIII, fig. 415.)

1900, p. 203, Pl. XXXIII, fig. 208; (?) *Oe. echinatum* Wood in Wille 1887, p. 86, Pl. 85, figs. 13-16.

Dioecious, nannandrous; oogonium 1, subglobose, pore superior; oospore globose, nearly filling oogonium, outer layer of spore wall echinate; dwarf male curved, on suffultory cell, stipe sometimes 2-4 celled; antheridium exterior, 1-?; vegetative cell $9-11 \times 36-100\mu$; oogonium $29-33 \times 32-35\mu$; oospore $26-29 \times 26-29\mu$; dwarf male, lower cell $7-8 \times 20-24\mu$, upper cell $4-6 \times 21-30\mu$; antheridium $5-6 \times 7-8\mu$.

Brazil.

Characterized by its echinate oospore wall with the spines not spirally arranged. It is smaller than either *Oe. echinatum* (No. 147) or *Oe. lindmanianum* (No. 148).

147. *Oe. echinatum* (Wood) Wittrock.

(Pl. XLV, fig. 436.)

Androgynia echinata Wood 1872, p. 198, Pl. 18, fig. 3; *Oe. echinatum* Wittr. 1878, p. 137; Hirn 1900, p. 204.

Dioecious, nannandrous; oogonium (?) 1, globose, usually depressed, pore superior; oospore globose, distending oogonium, outer layer of spore wall echinate; dwarf male suberect, near oogonium, (?) antheridium exterior, 1; vegetative cell $8-13 \times 50-138\mu$; oogonium $35-36 \times 35-36\mu$; oospore (without echinations) $25-26 \times 25-26\mu$.

United States: Pennsylvania.

Wood's figure of the echinate oospore is the only record of this species, other than the description above.

148. *Oe. lindmanianum* Wittrock.

(Pl. XLV, figs. 434 and 435.)

In Wittr., Nordst., and Lagerh. Alg. Exs. No. 1216, 1896; Hirn 1900, p. 204, Pl. XXXIV, fig. 209.

Dioecious, nannandrous, idioandrosporous; oogonium 1, obovoid-globose or subglobose, pore superior; oospore globose or subglobose, quite filling oogonium; outer spore wall echinate; androsporangium to 7-seriate; dwarf male a little curved, on suffultory cell, antheridium exterior, 1; vegetative cell $22-30 \times 40-120\mu$; oogonium $46-57 \times 46-57\mu$; oospore (with spines) $45-56 \times 45-54\mu$; androsporangium $22-28 \times 17-27\mu$; dwarf male stipe $12-15 \times 38-46\mu$; antheridium $6-8 \times 11-16\mu$.

Paraguay.

Cf. *Oe. armigerum* (No. 146).

149. *Oe. cyathigerum* Wittrock.

(Pl. LI, figs. 484 and 485.)

1870, p. 131, Pl. 1, figs. 6 and 7; *Oe. cyathigerum* Wittr. var. *rumelica* Istvanffy 1890, p. 77; *Oe. ornatum* Hirn 1896, p. 3, fig. (p. 4); Hirn 1900, p. 252, Pl. XLIII, figs. 265 and 266; Heering 1914, p. 184, fig. 256; Tiffany 1926, p. 103, Pl. IX, figs. 104 and 105.

Dioecious, nannandrous, idioandrosporous; oogonium 1-3, subovoid or quadrangular-ellipsoid, pore superior; oospore same form as oogonium,

filling it, outer layer of spore wall smooth, median layer with 16–25 longitudinal, continuous, rarely anastomosing, often curved, ridges—inner layer smooth; basal cell elongate; terminal cell, often an oogonium, obtuse; dwarf male goblet-shaped, curved, on the suffultory cell or oogonium; antheridium interior; vegetative cell $21-30 \times 40-300\mu$; suffultory cell $42-48 \times 75-110\mu$; oogonium $57-66 \times 70-100\mu$; oospore $51-62 \times 60-75\mu$; androsporangium $23-30 \times 12-30\mu$; dwarf male cell $12-15 \times 50-58\mu$.

United States: Iowa, Illinois, Ohio, Mississippi, Michigan; Denmark, France, Germany, Sweden, Rumania, Switzerland, England.

149a. Form **americanum** Wolle.

1887, p. 77, Pl. 81, figs. 20-22; Hirn 1900, p. 256.

Smaller; dwarf male on oogonium; vegetative cell $17-21 \times 22-62\mu$; oogonium $45-50 \times 55-63\mu$; oospore $39-40 \times 44-46\mu$; dwarf male $12-15 \times 50-54\mu$.

United States: Pennsylvania.

149b. Form **ornatum** (Wittrock) Hirn.

(Pl. LI, figs. 488-490.)

1900, p. 254, Pl. XLIII, fig. 267; *Oe. cyathigerum* Wittr. var. *ornatum* Wittr. 1878, p. 134.

Idioandrosporous; elongated dwarf males; vegetative cell $21-30 \times 60-270\mu$; suffultory cell $39-48 \times 60-116\mu$; oogonium $56-65 \times 69-102\mu$; oospore $54-60 \times 65-83\mu$; androsporangium $21-27 \times 13-27\mu$; dwarf male $12-16 \times 60-75\mu$.

Mexico (Vera Cruz).

149c. Form **perfectum** Hirn.

(Pl. LI, figs. 486 and 487.)

1900, p. 254; Pl. XLIII, fig. 268; *Oe. cyathigerum* Wittr. in Borge 1896, p. 5.

Larger; androsporangium to 23-seriate; vegetative cell $22-35 \times 65-350\mu$; suffultory cell $37-56 \times 56-140\mu$; oogonium $65-81 \times 72-111\mu$; oospore $61-78 \times 65-85\mu$; androsporangium $21-30 \times 15-30\mu$; dwarf male $12-16 \times 55-69\mu$.

United States: Mississippi, Ohio; Brazil; Australia; India.

149d. Var. **ellipticum** Magnus and Wille.

(Pl. LII, fig. 493.)

In Wille 1884, p. 40, Pl. 2, fig. 67; Hirn 1900, p. 255, Pl. XLIV, fig. 271; Fritsch and Rich 1924, p. 320, fig. 4 A-C.

Oospore frequently not filling oogonium, quadrangular-ellipsoid or nearly ellipsoid or rarely subglobose; vegetative cell $19-30 \times 30-120\mu$; suffultory cell $26-43 \times 40-125\mu$; oogonium $50-63 \times 68-94\mu$; oospore $48-57 \times 45-66\mu$; dwarf male $15-18 \times 43-55\mu$.

United States: Ohio; Uruguay, Brazil; South Africa.

149e. Var. **hormosporum** (West) Hirn.

(Pl. LII, figs. 491 and 492.)

1900, p. 255, Pl. XLIV, figs. 269 and 270; *Oe. hormosporum* West 1897, p. 5.

Oogonium and oospore smaller; oogonia 2–10 (rarely 1), androsporangium up to 15-seriate; vegetative cell $22-30 \times 76-180\mu$; suffultory cell $35-45 \times 52-112\mu$; oogonium $48-57 \times 56-95\mu$; oospore $46-55 \times 53-65\mu$; androsporangium $18-22 \times 12-24\mu$; dwarf male $9-13 \times 47-54\mu$.

Africa.

The species is readily identified among the nannandrous group by the irregular longitudinal ridges appearing on the median oospore wall. The suffultory cells are very much enlarged. The varieties and forms are separated largely on the basis of size. It differs from *Oe. wolleanum* (No. 150) in its smaller number of ribs. Fritsch and Rich (1924) record a form of *Oe. cyathigerum* with suffultory cell not always swollen and with frequently globose oospores. This is perhaps merely a slight variation of var. *ellipticum*.

150. **Oe. wolleanum** Wittrock.

(Pl. I, figs. 479 and 480.)

1878, p. 137; Wille 1887, p. 82, Pl. 80, figs. 4 and 5; *Oe. aphophysatum* A. Br. in Rabenh. Alg. Sach. No. 291, 1853; *Oe. concatenatum* (Hass.) Kuetz. f. *luxurians* Breb. in Roumeg. Alg. Exs. Nos. 683 and 1062; Hirn 1900, p. 220, Pl. XXXVII, fig. 226.

Dioecious, nannandrous, gynandrosporous or idioandrosporous; oogonium 1–4, subobovoid or quadrangular-ellipsoid, pore superior, wall with raised longitudinal lines on the inner surface; oospore of same form as oogonium, quite filling it, outer layer of spore wall with 25–35 longitudinal ribs, rarely anastomosing, inner layer smooth; androsporangium 1–3, often subepigynous or scattered in the upper part of the filament; basal cell elongate; terminal cell, sometimes oogonium, short-acute or acuminate; dwarf male on suffultory cell, stipe slightly curved; antheridium exterior, 1–3; vegetative cell $21-30 \times 65-235\mu$; suffultory cell $45-56 \times 68-110\mu$; oogonium $58-68 \times 69-89\mu$; oospore $56-66 \times 65-83\mu$; androsporangium $21-30 \times 18-25\mu$; dwarf male stipe $15-24 \times 54-60\mu$; antheridium $9-14 \times 7-11\mu$.

United States: Massachusetts, Connecticut, New Jersey, Iowa, Pennsylvania, Minnesota, Florida, Illinois; Brazil; India; France, Latvia, Germany; Australia; Greenland.

150a. Var. **concinnum** Hirn.

(Pl. I, fig. 481.)

1900, p. 222, Pl. XXXVII, fig. 228.

Smaller; oospore not quite filling oogonium; ribs of outer layer of spore wall finely crenulate; vegetative cell $19-27 \times 80-180\mu$; suffultory cell $41-52 \times 72-150\mu$; oogonium $54-63 \times 63-93\mu$; oospore $52-61 \times 62-74\mu$; dwarf male stipe $16-20 \times 50-30\mu$; antheridium $10-13 \times 10-15\mu$.

United States: Minnesota.

150b. Form **insigne** (Nordstedt) Hirn.

(Pl. L, figs. 482 and 483.)

1900, p. 222, Pl. XXXVII, fig. 227; *Oe. wolleanum* Wittr. var. *insigne* Nordst. in Wittr. and Nordst. Alg. Exs. No. 207, 1879.

Stouter; ribs on the oogonium and oospore 35–40; oogonium 1–10; androsporangium to 10-seriate; antheridium 1–4; vegetative cell 18–36 × 53–250 μ ; suffultory cell 57–66 × 100–150 μ ; oogonium 67–80 × 78–92 μ ; oospore 64–76 × 74–90 μ ; androsporangium 19–30 × 18–30 μ ; dwarf male stipe 18–24 × 54–68 μ ; antheridium 12–14 × 10–12 μ .

United States: New Jersey; Sweden.

This species is characterized by regular, longitudinal ribs on the outer layer of the oospore wall. It differs from *Oe. cyathigerum* (No. 149) in this respect and also in having a larger number of ribs. Var. *concinnum* has crenulate ribs, and f. *insigne* differs in its slightly larger size and in the larger number of ribs on the oospore.

151. **Oe. concatenatum** (Hassall) Wittrock.

(Pl. LII, fig. 494.)

Vesiculifera concatenata Hass. 1845, p. 201, Pl. 51, fig. 3; *Oe. concatenatum* Wittr. 1874, p. 25; *Oe. apophysatum* Al. Br. in Pringsh. 1858, p. 71, Pl. 5, fig. 9; *Oe. apophysatum* Pringsh. in Wittr. 1872a, p. 22; *Oe. concatenatum* (Hass.) Wittr. var. *lagenarioides* Pilarszky 1899, p. 65, Pl. 3; Hirn 1900, p. 223, Pl. XXXVIII, fig. 230; Heering 1914, p. 180, fig. 245; Tiffany 1926, p. 104, Pl. VIII, fig. 92.

Dioecious, nannandrous, gynandrosporous; oogonium 1–6, subovoid or quadrangular-ellipsoid, pore superior; oospore of same form as oogonium, nearly filling it, outer layer of spore wall smooth, middle layer with pits, more or less distinctly arranged in 30–35 longitudinal series, inner wall smooth; suffultory cell swollen; androsporangium 1–4; basal cell elongate; terminal cell obtuse; dwarf male curved, on suffultory cell; antheridium exterior, 1–4; vegetative cell 25–40 × 75–400 μ ; suffultory cell 50–62 × 88–155 μ ; oogonium 63–83 × 76–105 μ ; oospore 60–75 × 67–95 μ ; androsporangium 25–28 × 15–36 μ ; dwarf male stipe 17–25 × 50–75 μ ; antheridium 13–15 × 12–25 μ .

United States: Illinois, Iowa, Massachusetts, New Jersey, Pennsylvania, New York, Michigan; Austria, England, Finland, Germany, Spain, Hungary, Sweden, Denmark; Alaska; British Columbia.

151a. Form **hutchinsiae** (Wittrock) Hirn.

(Pl. LII, fig. 495.)

1900, p. 225, Pl. XXXVIII, fig. 231; *Oe. hutchinsiae* Wittr. 1874, p. 42; Heering 1914, p. 180; Tiffany 1926, p. 104.

Form a little smaller, oospore more nearly globose, the pits of the middle layer of spore wall not arranged in evident longitudinal series; vegetative cells 26–35 × 75–210 μ ; suffultory cell 37–50 × 65–200 μ ; oogonium 52–75 × 67–95 μ ; oospore 50–73 × 55–77 μ .

United States: Iowa; England, France, Spain.

151b. Var. *rectangulare* Rich.

(Pl. LIII, figs. 496 and 497.)

1925, p. 72, fig. 1.

Oospore quadrangular-ellipsoid, membrane thick, middle layer of spore wall with its pits in transverse and longitudinal rows (about 15 transverse rows); idioandrosporous; vegetative cell $32-36 \times 75-400\mu$; suffultory cell $50 \times 88-155\mu$; oogonium $56 \times 76-105\mu$; oospore $48-56 \times 67-95\mu$; androsporangium $27 \times 15-36\mu$.

England.

Oe. concatenatum is unique among the poriferous, nannandrous forms of the genus for its prominently pitted oospore. *F. hutchinsiae* and var. *rectangulare* are separable largely on the arrangement of the pits on the median layer of the oospore.

152. *Oe. huntii* Wood.

(Pl. XLVIII, figs. 465-468.)

1869, p. 333; Wittrock 1878, p. 136; *Androgynia huntii* Wood 1872, p. 197, Pl. 17, fig. 2; Hirn 1900, p. 208, Pl. XXXIV, fig. 213; Collins 1908, p. 57; P. B. A. No. 1471.

Dioecious, nannandrous; oogonium usually 1, subglobose or subobovoid-globose (rarely subhexagonal), pore inferior; oospore globose, not filling oogonium, outer spore wall with four raised spiral ribs; basal cell elongate; terminal cells tapering, setiferous (consisting of 10 or more superimposed long cylindrical cells, each smaller in diameter than the one below); dwarf male nearly straight, on suffultory cell, antheridium exterior, 1-2 (?); vegetative cell $15-25 \times 36-82\mu$; oogonium $50-60 \times 52-60\mu$; oospore $38-42 \times 38-42\mu$; dwarf male stipe $11 \times 52\mu$; antheridium $10 \times 30\mu$.

United States: Massachusetts, Pennsylvania.

Fruiting material showing the 4 spiral ribs has been seen only by Wood. Collins (1908) records the species showing the peculiar tapering setiferous terminal cells (see description above). This is the only species in the nannandrous group having ribbed oospores and an inferior pore.

153. *Oe. hispidum* Nordstedt.

(Pl. XLV, fig. 440.)

In Wittr. 1870, p. 128, and 1874, p. 28; Hirn 1900, p. 210, Pl. XXXV, fig. 215.

Dioecious, nannandrous, gynandrosporous; oogonium 1, terminal, subellipsoid or ellipsoid-globose, pore inferior; oospore globose to globose-ellipsoid, not filling oogonium or rarely enlarging it, outer spore wall echinate; androsporangia 2; terminal cell obtuse; dwarf male a little curved, on suffultory cell, antheridium exterior, 1; vegetative cell $9-14 \times 36-130\mu$; oogonium $35-44 \times 42-56\mu$; oospore (with spines) $32-39 \times 32-40\mu$; androsporangium $10-12 \times 6-8\mu$; dwarf male stipe $7-8 \times 17-18\mu$; antheridium $5-6 \times 7-9\mu$.

France, Sweden.

The only nannandrous species with echinate oospores and inferior position of the oogonial pore.

154. *Oe. cleveanum* Wittrock.

(Pl. XLVI, figs. 443 and 444.)

1870, p. 128; 1872a, p. 22; 1874, p. 28; *Oe. echinospermum* Al. Br. in Pringsh. 1858, p. 70, Pl. 5, fig. 7; *Oe. berolinense* Wittr. 1872, p. 6; *Oe. cleveanum* Wittr. var. *arvensis* Istvanffy 1887, p. 242; Hirn 1900, p. 209, Pl. XXXV, fig. 214; *Oe. cleveanum* Wittr. f. *exoticum* Hirn 1900, p. 210 (not Hirn 1906, p. 33).

Dioecious, nannandrous, gynandrosporous; oogonium 1, subobovoid-globose or subglobose, pore inferior; oospore globose, filling oogonium, outer layer of spore wall dentate, teeth in 4-6 spiral rows, anastomosate; suffultory cell sometimes tumid; androsporangium 1-7; dwarf male a little curved, on suffultory cell, antheridium exterior, 1; vegetative cell (14-) $18-26 \times 40-110\mu$; oogonium $45-60 \times 48-63\mu$; oospore (with teeth) $44-57 \times 45-59\mu$; androsporangium $18-22 \times 9-18\mu$; dwarf male stipe $10-11 \times 29-32\mu$; antheridium $5-9 \times 12-16\mu$.

England, France, Germany, Hungary, Sweden; Brazil.

154a. Form *exoticum* Hirn.

1906, p. 33, (not 1900, p. 210); Schmidle 1901, p. 344.

Oogonium 1-3; vegetative cell $14-25 \times 44-200\mu$; oogonium $39-46 \times 40-47\mu$; oospore (with teeth) $38-44 \times 38-44\mu$; dwarf male stipe $10-11 \times 27-30\mu$; antheridium $6-7 \times 11-13\mu$.

Caroline Islands.

Oe. cleveanum is characterized by the inferior position of the oogonial pore and the dentate oospore, the teeth being arranged somewhat spirally. The form *exoticum* as described by Hirn (1900) is so nearly identical with the species as to be untenable even as a form. The specimens from the Caroline Islands referred to f. *exoticum* by Schmidle (1901) and Hirn (1906) have distinctly smaller oogonia and oospores and are thus perfectly tenable.

155. *Oe. elegans* West and West.

(Pl. LIII, figs. 498 and 499.)

1902, p. 128, Pl. 17, figs. 6-7; Hirn 1906, p. 13, Pl. IV, fig. 22.

Dioecious, nannandrous; oogonium 1, depressed-globose, operculate, division median; oospore depressed-globose, quite filling oogonium, spore wall smooth; dwarf male small, unicellular, on oogonium; vegetative cell densely and minutely granulate, the deposits spirally arranged; vegetative cell $6-9 \times 25-85\mu$; oogonium $27-31 \times 25-26\mu$; oospore $25-28 \times 18-19\mu$; dwarf male $5-6 \times 8-9\mu$.

Ceylon.

The only nannandrous member of the genus with granulate vegetative cells. Cf. *Oe. minus* (No. 84).

156. *Oe. rigidium* Hirn.

(Pl. LIII, figs. 506 and 507.)

1900, p. 237, Pl. XL, fig. 244.

Dioecious, nannandrous, gynandrosporous; oogonium 1, obovoid-globose, operculate, division superior; oospore globose, nearly filling oogonium, spore wall smooth, often thick; androsporangium 1-2, epigynous, subepigynous or hypogynous; vegetative cell broadly capitellate; terminal cell obtuse; dwarf male curved, on oogonium, antheridium exterior, 1-?; vegetative cell $12-14 \times 38-58\mu$; oogonium $35-39 (-42) \times 38-45\mu$; oospore (30-) $34-38 \times 34-38\mu$; androsporangium $10-11 \times 11-12\mu$; dwarf male stipe $7-9 \times 18-22\mu$; antheridium $5-7 \times 6-7\mu$.

Brazil; South Africa.

The only nannandrous species having the combination of superior position of the oogonial operculum, and capitellate vegetative cells.

157. *Oe. bengalense* Hirn.

(Pl. LIII, figs. 504 and 505.)

1900, p. 268, Pl. XLVI, fig. 287.

Dioecious, nannandrous, idioandrosporous; oogonium 1-4, subdepressed-globose to ovoid, operculate, division median, rather wide; oospore subdepressed-globose or globose; quite filling oogonium, spore wall smooth; androsporangium 1-4; vegetative cell capitellate; basal cell elongate; terminal cell, not rarely the oogonium, apically obtuse; dwarf male obovoid, unicellular, on oogonium; vegetative cell $14-19 \times 45-110\mu$; oogonium $46-54 \times 44-54\mu$; oospore $43-48 \times 43-48\mu$; androsporangium $14-15 \times 13-18\mu$; dwarf male $12-13 \times 16-18\mu$.

India: Bengal.

Cf. *Oe. indicum*, below.158. *Oe. indicum* Hirn.

(Pl. LIV, fig. 509.)

1900, p. 269, Pl. XLVI, fig. 288.

Dioecious, nannandrous, gynandrosporous; oogonium 1-2, depressed-globose or depressed obovoid-globose, operculate, division median, narrow but distinct; oospore depressed-globose, completing oogonium, spore wall smooth; androsporangium 1-2-?; subepigynous; vegetative cell broadly capitellate; basal cell elongate; dwarf male obovoid, unicellular, on oogonium; vegetative cell $20-25 \times 50-110\mu$; oogonium $57-64 \times 48-53\mu$; oospore $55-61 \times 43-50\mu$; androsporangium $19-20 \times 8-10\mu$; dwarf male $11-14 \times 18\mu$.

India.

This species together with *Oe. bengalense* above combine the characters of capitellate vegetative cells and median position of a narrow operculum. *Oe. indicum* is the larger of the two. Cf. *Oe. areschougii* (No. 160), *Oe. brasiliense* (No. 159), and *Oe. confertum* (No. 161).

159. *Oe. brasiliense* Borge.

(Pl. LIV, fig. 508.)

1899, p. 4, Pl. 1, fig. 1; Hirn 1900, p. 272, Pl. XLVI, fig. 292.

Dioecious, nannandrous, idioandrosporous; oogonium 1-3, subdepressed-globose or subpyriform-globose, operculate, division median, broad; oospore subdepressed-globose, not quite filling oogonium, spore wall smooth; androsporangium to 5-seriate; vegetative cell capitellate; dwarf male obovoid, unicellular, on oogonium; vegetative cell $16-22 \times 35-95\mu$; oogonium $53-63 \times 52-59\mu$; oospore $48-53 \times 45-50\mu$; androsporangium $15-16 \times 11-14\mu$; dwarf male $10-12 \times 14-19\mu$.

Brazil.

Characterized by its broad operculum and capitellate vegetative cells.

160. *Oe. areschougii* Wittrock.

(Pl. LIII, fig. 500.)

1870, p. 122, Pl. 1, figs. 1 and 2; 1874, p. 19, Pl. 1, figs. 10 and 11; Hirn 1900, p. 270, Pl. XLVI, fig. 289.

Dioecious, nannandrous, gynandrosporous; oogonium 1-6, subdepressed- or depressed pyriform-globose, operculate, division median, broad; oospore globose, rarely subdepressed-globose, not completing oogonium longitudinally, spore wall smooth; androsporangium 1-6, subepigynous or hypogynous or rarely scattered; vegetative cell capitellate; basal cell elongate; terminal cell, usually the oogonium, apically obtuse; dwarf male obovoid, unicellular, on oogonium; vegetative cell $8-13 \times 35-75\mu$; oogonium $34-39 \times 36-40\mu$; oospore $22-26 \times 22-25\mu$; androsporangium $9-11 \times 10-12\mu$; antheridium $6-7 \times 13-15\mu$.

United States: New Jersey; Brazil; Finland, Germany, Sweden, Denmark; Greenland; Siberia.

160a. Var. *americanum* Tiffany.

(Pl. LIII, fig. 501.)

1927, p. 205, Pl. IX, figs. 8 and 9.

Idioandrosporous; oogonium smaller; androsporangium to 11-seriate; vegetative cell $8-12 \times 40-80\mu$; oogonium $29-36 \times 26-40\mu$; oospore $23-26 \times 22-26\mu$; androsporangium $7-9 \times 7-11\mu$.

United States: Illinois.

160b. Form *robustum* Hirn.

(Pl. LIII, figs. 502 and 503.)

1900, p. 271, Pl. XLVI, fig. 290; (?) *Oe. obtruncatum* var. *oblatum* Tilden, Amer. Algæ No. 3, 1894; *Oe. areschougii* f. *major* Collins 1912, p. 88.

Idioandrosporous; oospore larger; oogonium to 8-seriate; vegetative cell $12-17 \times 36-120\mu$; oogonium $36-40 \times 36-53\mu$; oospore $30-32 \times 27-31\mu$; androsporangium $9-12 \times 10-13\mu$; dwarf male $6-8 \times 14-15\mu$.

England; (?) Minnesota (U. S.).

This species is characterized by its capitellate vegetative cells, wide operculum, small size, and small oospores in comparison with oogonia. Var. *americanum* differs from the species in being idioandrosporous; f. *robustum* in addition has larger oospores.

161. **Oe. confertum** Hirn.

(Pl. LIV, fig. 510.)

1900, p. 272, Pl. XLVI, fig. 291.

Dioecious, nannandrous, (?) idioandrosporous; oogonium 1-4, depressed-globose or depressed pyriform, operculate, division median, wide; oospore globose or subdepressed-globose, not filling oogonium longitudinally, spore wall smooth; vegetative cell capitellate; dwarf male obovoid, on oogonium; vegetative cell $19-26 \times 66-105\mu$; oogonium $56-63 \times 44-56\mu$; oospore $42-48 \times 40-44\mu$; dwarf male $10-12 \times 13-16\mu$.

Australia.

To be compared with *Oe. areschougii* (No. 160), which is smaller and has less depressed oogonia. It is also similar in appearance to the larger *Oe. perspicuum*, below.

162. **Oe. perspicuum** Hirn.

(Pl. LIV, fig. 511.)

1900, p. 273, Pl. XLVI, fig. 293; *Oe. dioicum* Carter in West and West 1901, p. 175, Pl. IV, fig. 42.

Dioecious, nannandrous; oogonium 1-8, depressed or subdepressed-globose, operculate, division median, broad; oospore globose or subglobose, not filling oogonium longitudinally, spore wall smooth; dwarf male obovoid, unicellular, on oogonium; vegetative cell (30-) $35-43 \times 95-175\mu$; oogonium $89-100 \times 64-104\mu$; oospore $55-70 \times 52-70\mu$; dwarf male $15-18 \times 19-22\mu$.

Australia; Siam.

Cf. *Oe. confertum*, above.

163. **Oe. clavatum** Hallas.

(Pl. LIV, figs. 512 and 513.)

1905, p. 399, fig. 3; Hirn 1906, p. 11, Pl. IV, fig. 23.

Dioecious, nannandrous, gynandrosporous; oogonium 1-3, subpyriform to subdepressed-globose, operculate, division a little above medium; oospore subdepressed, depressed-globose, or globose, quite filling oogonium (rarely not), spore wall smooth; androsporangium 1-2, scattered; dwarf male obovoid, unicellular, on oogonium; vegetative cell capitellate or nearly clavate; basal cell subhemispherical; vegetative cell $2-9 \times 5-80\mu$; oogonium $14-26 \times 16-21\mu$; oospore $12-21 \times 12-14\mu$; androsporangium $4-6 \times 5-7\mu$; dwarf male $6 \times 14-15\mu$.

United States: New York (Long Island); Denmark.

One of the smallest nannandrous species with capitellate or nearly clavate vegetative cells.

164. *Oe. oelandicum* Wittrock; Hirn.

(Pl. LVI, fig. 539.)

Wittrock 1874, p. 17; Hirn 1900, p. 273, Pl. XLVII, fig. 297.

Dioecious, nannandrous, gynandrosporous; oogonium 1-7, depressed-globose, with 12-16 longitudinal ridges, operculate, division supramedian, rather broad; oospore depressed-globose, nearly filling oogonium, spore wall smooth; androsporangium to 6-seriate; vegetative cell capitellate; terminal cell apically obtuse; dwarf male obovoid, unicellular, on oogonium; vegetative cell $10-15 \times 25-125\mu$; oogonium $31-40 \times 25-32\mu$; oospore $25-36 \times 23-30\mu$; androsporangium $7-12 \times 10-18\mu$; dwarf male $7-8 \times 12-15\mu$.

United States: Michigan; British Columbia; Sweden.

164a. Form *minus* Borge.

(Pl. LVI, fig. 540.)

1911, p. 205, Pl. 2, fig. 10.

Smaller; vegetative cell $9-10 \times 36-50\mu$; oogonium $27-30 \times 19-25\mu$; dwarf male $6-7 \times 11-12\mu$.

The species is characterized by its longitudinally ridged oogonia, supramedian operculum, and capitellate vegetative cells. It is smaller than the similarly appearing, larger *Oe. megaporum*, below. *F. minus* has shorter oogonia.

165. *Oe. megaporum* Wittrock.

(Pl. LVI, fig. 543.)

1872, p. 3, Pl. 1, figs. 5 and 6; Hirn 1900, p. 274, Pl. XLVII, figs. 298 and 299; *Oe. oelandicum* Wittr. var. *subpyriforme* Wittr. in Wittr., Nordst., and Lagerh. Alg. Exs. No. 1218, 1896.

Dioecious, nannandrous, idioandrosporous; oogonium 1-6, broadly pyriform, with 12-16 rounded longitudinal ridges, operculate, division supramedian and wide; oospore subdepressed-globose, not filling oogonium, spore wall smooth; androsporangium to 4-seriate; vegetative cell capitellate, basal cell elongate; terminal cell apically obtuse; dwarf male unicellular, broadly obovoid, on oogonium; vegetative cell $13-17 \times 40-100\mu$; oogonium $37-42 \times 40-45\mu$; oospore $31-35 (-38) \times 27-30\mu$; androsporangium $10-11 \times 12-22\mu$; dwarf male $8-12 \times 13-16\mu$.

United States: Michigan; Finland, Germany, Sweden.

Cf. *Oe. oelandicum* (No. 164).

166. *Oe. bahusiense* Nordstedt.

(Pl. LV, figs. 530 and 531.)

1877, p. 26, Pl. 3, figs. 7-11; Hirn 1900, p. 279, Pl. XLVIII, fig. 307.

Dioecious, nannandrous, gynandrosporous; oogonium 1 (rarely 2), depressed pyriform, with 9-12 longitudinal ridges appearing as a median whorl, operculate, division inframedian; oospore depressed-obovoid or depressed-globose, nearly filling oogonium, spore wall smooth; androsporangium to 4-seriate, scattered, often terminal; vegetative cell

capitellate; dwarf male unicellular, obovoid, small, on oogonium; vegetative cell $12-15 \times 25-60\mu$; oogonium $30-36 \times 22-29\mu$; oospore $23-28 \times 20-24\mu$; androsporangium $10-13 \times 5-7\mu$; dwarf male $5-6 \times 8-9\mu$.

Sweden.

This species combines the inframedian position of the operculum, ridged oogonium, capitellate vegetative cell, and non-tumid suffultory cell. It differs from *Oe. platygynum*, below, in its larger dimensions.

167. *Oe. platygynum* Wittrock.

(Pl. LVI, fig. 544.)

1872, p. 1; 1874, p. 17, Pl. 1, figs. 5-9; *Oe. platygynum* Wittr. f. *major* West 1891, p. 109, Pl. 18, fig. 1; Hirn 1900, p. 276, Pl. XLVII, figs. 301 and 302; P. B. A. No. 1677.

Dioecious, nannandrous, gynandrosporous and idioandrosporous; oogonium 1-2, depressed obovoid, with 7-12 (more often 8) rounded projections arranged in a whorl around the middle, operculate, division inframedian; oospore depressed- or subdepressed-globose, not quite filling oogonium, spore wall smooth; suffultory cell sometimes tumid; androsporangium 1-3; vegetative cell evidently capitellate; terminal cell apically obtuse; dwarf male unicellular, obovoid, small, on oogonium; vegetative cell $5-11 \times 14-50\mu$; oogonium $21-30 \times 16-24\mu$; oospore $17-24 \times 13-20\mu$; androsporangium $6-8 \times 7-8\mu$; dwarf male $4-5 \times 8-10\mu$.

United States: New Jersey, Pennsylvania, Florida, Minnesota; Brazil; England, Denmark, Finland, Germany, Ireland, Norway, Latvia, Sweden; Africa; Ceylon.

167a. Form *obtusum* Hirn.

(Pl. LVI, fig. 545.)

1900 p. 277, Pl. XLVII, fig. 303.

Idioandrosporous, oogonium a little less depressed, processes obtusely rounded; vegetative cell $6-10 \times 14-50\mu$; oogonium $22-26 \times 18-24\mu$; oospore $17-19 \times 14-15\mu$; androsporangium $6-8 \times 8-10\mu$; dwarf male $4-5 \times 7-9\mu$.

England, Sweden.

167b. Var. *continuum* Nordstedt.

(Pl. LVI, fig. 546.)

1888, p. 12, Pl. 1, figs. 16 and 17; Hirn 1900, p. 278, Pl. XLVII, fig. 305.

Oogonium to .5-seriate, projections more truncately rounded; suffultory cell enlarged; vegetative cell $8 \times 48\mu$; suffultory cell $12-14 \times 25-40\mu$; oogonium $28-32 \times 21-26\mu$; oospore $22 \times 20\mu$; ? androsporangium $15-17 \times 12\mu$.

New Zealand.

167c. Var. *novæ zelandiæ* Hirn.

(Pl. LVI, fig. 547.)

1900, p. 278, Pl. XLVII, fig. 304; *Oe. platygynum* Wittr. *forma* Nordst. 1888, p. 12, Pl. I, figs. 14 and 15.

Gynandrosporous; oogonium subpyriform, projections obtuse or truncately rounded; vegetative cells very broadly capitellate, $6-9 \times 20-50\mu$; oogonium $24-28 \times 22-26\mu$; oospore $20-22 \times 18-20\mu$; androsporangium $7-8 \times 7-11\mu$.

New Zealand.

Cf. *Oe. bahusiense* (No. 166).

168. *Oe. longicolle* Nordstedt.

(Pl. LV, figs. 516 and 517.)

1878, p. 20, Pl. 2, figs. 11 and 12; Hirn 1900, p. 263, Pl. XLV, fig. 278.

Diocious, nannandrous; oogonium 1-7, pyriform, often elongated longitudinally, operculate, division median and narrow; oospore subdepressed-globose or subglobose, inflating oogonium medianly, not filling it longitudinally, spore wall smooth; basal cell subhemispherical; dwarf male unicellular, obovoid, minute, on oogonium; vegetative cell $4-6 \times 16-45\mu$; oogonium $13-16 \times 16-32\mu$; oospore $12-15 \times 10-16\mu$; dwarf male $2 \times 4\mu$; basal cell $14 \times 6\mu$.

Australia; Ceylon.

168a. Var. *senegalense* Nordstedt.

(Pl. LV, figs. 518 and 519.)

1880, p. 13, Pl. 1, fig. 23; (?) *Oe. delicatulum* Kuetz. in Wolle 1887, Pl. S1, figs. 12 and 13; Hirn 1900, p. 264, Pl. XLV, figs. 279 and 280. *Oe. longicolle* Nordst. var. *senegalense* Nordst. f. *afghanicum* Schaarsch. 1884, p. 249, Pl. 5, fig. 26, and in Hirn 1900, p. 264, Pl. XLV, fig. 281.

Larger; vegetative cell $5-8 \times 12-40\mu$; oogonium $16-20 \times 14-24\mu$; oospore $14-20 \times 11-17\mu$; basal cell $13-14 \times 5-7\mu$.

Africa; Brazil; Ceylon; Australia; Afghanistan; Burma.

The species is characterized by its small size, subhemispherical basal cell, nannandrous habit, and smooth oogonium. It is the smallest of similar species that follow: *Oe. rothii* (No. 169) and *Oe. decipiens* (No. 171).

169. *Oe. rothii* (Le Clerc) Pringsheim.

(Pl. LV, fig. 526.)

(?) *Prolifera rothii* Le Clerc 1817, p. 476, Pl. 23, fig. 8; *Oe. rothii* Pringsh. 1858, p. 69, Pl. 5, fig. 4; *Oe. cryptoporum* Wittr. var. *vulgare* Wittr. f. *abbreviata* Gutw. 1897, p. 126; *Oe. cryptoporum* Wittr. var. *subdepressum* Wittr. in Wittr. and Nordst. Alg. Exs. No. 152, 1878; Hirn 1900, p. 265, Pl. XLV, fig. 282; Collins, 1909, p. 262; P. B. A. No. 520.

Diocious, nannandrous, gynandrosporous; oogonium 1-3, subdepressed-globose, operculate, division median, narrow; oospore depressed-globose, almost filling oogonium, spore wall smooth; androsporangium 1-4; subhypogynous, hypogynous, subepigynous, or scattered; vegetative cell $6-10 \times 20-76\mu$; oogonium $20-27 \times 16-27\mu$; oospore $17-25 \times 14-20\mu$; androsporangium $6-8 \times 5-10\mu$; dwarf male $4 \times 11-12\mu$.

United States: Massachusetts; Austria, Denmark, England, Germany, Norway, Sweden.

Cf. *Oe. longicolle*, above.

170. *Oe. danicum* Hallas.

(Pl. LIV, fig. 515.)

1905, p. 404, fig. 10; Hirn 1906, p. 12, Pl. IV, fig. 24.

Dioecious, nannandrous, gynandrosporous; oogonium 1, pyriform or pyriform-globose, operculate, division supramedian; oospore globose or depressed-globose, filling oogonium or not, spore wall smooth; androsporangium 1-3, hypogynous or scattered; dwarf male obovoid, unicellular, on suffultory cell or scattered; vegetative cell $4-7 \times 20-82\mu$; oogonium $21-23 \times 23-35\mu$; oospore $19-20 \times 15-20\mu$; androsporangium $4-7 \times 7-12\mu$; dwarf male $6-7 \times 14-16\mu$.

Denmark.

171. *Oe. decipiens* Wittrock.

(Pl. LV, fig. 520.)

1870, p. 126; *Oe. vesicatum* Link in De Bary 1856, p. 224, Pl. 5; *Oe. rothii* (Le Cl.) Pringsh. f. *major* West and West 1897, p. 5; Hirn 1900, p. 266, Pl. XLVI, figs. 283 and 284; Collins 1909, p. 262; Heering 1914, p. 188, fig. 258; Tiffany 1926, p. 104, Pl. VIII, fig. 88.

Dioecious, nannandrous, gynandrosporous; oogonium 1-3, subdepressed globose, operculate, division median, rather narrow; oospore subdepressed or depressed-globose, almost filling the oogonium, spore wall smooth; suffultory cell not swollen; androsporangium 1-6, subepigynous, hypogynous, or scattered; dwarf male unicellular, usually on the oogonium; vegetative cell $9-12 \times 28-80\mu$; oogonium $30-38 \times 27-40\mu$; oospore $25-34 \times 23-28\mu$; androsporangium $9-10 \times 8-15\mu$; dwarf male $6-7 \times 13-15\mu$.

United States: Michigan, Iowa, New Jersey; France, Germany, Sweden, England; Africa.

171a. Form *dissimile* Hirn.

(Pl. LV, figs. 521 and 522.)

1900, p. 267, Pl. XLVI, fig. 285.

Gynandrosporous; vegetative cells evidently capitellate, $8-11 \times 25-65\mu$; oogonium $28-35 \times 23-38\mu$; oospore $23-34 \times 21-30\mu$; androsporangium $8-9 \times 6-10\mu$; dwarf male $5-6 \times 11-14\mu$.

United States: Michigan; Brazil.

171b. Var. *africanum* Tiffany.

(Pl. LV, figs. 523 and 524.)

1929, p. 74; *Oe. decipiens* Wittr. forma in West and West 1897, p. 5, and in Hirn 1900, p. 267, Pl. XLVI, fig. 284.

Smaller and idioandrosporous; vegetative cell $8-13 \times 25-60\mu$; oogonium $25-32 \times 24-32\mu$; oospore $24-30 \times 23-28\mu$; androsporangium $7-11 \times 9-13\mu$; dwarf male $6-7 \times 9-12\mu$.

Africa.

171c. Var. **bernardense** (Bates) Hirn.

(Pl. LV, fig. 525.)

1900, p. 268, Pl. XLVI, fig. 286; *Oe. bernardense* Bates 1886, p. 313; *Oe. londonense* Wittr. var. *compressum* West 1891, p. 110, Pl. 18, figs. 10-12.

Smaller; gynandrosporous; androsporangium to 8-seriate; vegetative cell $8-12 \times 18-45\mu$; oogonium $25-31 \times 21-32\mu$; oospore $21-29 \times 17-24\mu$; androsporangium $7-11 \times 5-11\mu$; dwarf male $6-7 \times 10-12\mu$.

England, Spain.

The tendency of the *decipiens* group toward capitellate cells reaches a maximum in f. *dissimile*. The others are either not noticeably or only somewhat capitellate. The small size, nannandrous habit, and median operculum are also characteristic. *F. africanum* is idioandrosporous, while f. *bernardense* is gynandrosporous.

172. **Oe. macrospermum** West and West.

(Pl. LV, figs. 527 and 528.)

1897, p. 472, Pl. 7, figs. 6 and 7; Hirn 1900, p. 227, Pl. XXXVIII, fig. 232; *Oe. macrospermum* West and West f. *patagonicum* Borge and Hirn in Borge 1901, p. 9, Pl. 1, fig. 1, and in Hirn 1906, p. 18, Pl. III, fig. 18; Tiffany 1926, p. 105, Pl. VIII, figs. 93 and 94.

Dioecious, nannandrous, gynandrosporous; oogonium 1 (rarely 2), subdepressed to depressed-globose, operculate, division median, narrow; oospore subdepressed-globose, filling the oogonium, spore wall smooth; dwarf male slightly curved or straight, not infrequently at right angles to the filament, on suffultory cell or on oogonium or scattered; antheridium exterior, 1-4; basal cell elongate; androsporangium 1-?; vegetative cell $10-16 \times 30-80\mu$; oogonium $39-46 \times 34-44\mu$; oospore $36-44 \times 32-42\mu$; dwarf male stipe $9-14 \times 16-30\mu$; androsporangium $14-15 \times 10-11\mu$; antheridium $7-10 \times 6-16\mu$.

United States: Iowa, Michigan, Ohio; England, Latvia; Patagonia.

The above description as emended by Tiffany (1926) from material collected in Iowa includes f. *patagonicum* Borge and Hirn (Borge 1901). The species is characterized by its particularly straight vegetative cells, narrow median operculum, and dwarf males scattered and often at right angles to the filament.

173. **Oe. costatum** Transeau nov. sp.

(Pl. LVI, fig. 548.)

Dioecious, nannandrous, gynandrosporous; oogonium 1, depressed globose, with verticillate folds forming distinct pointed projections at the suture, operculate, division narrow, suprmedian; oospore depressed-globose, filling oogonium, spore wall smooth; androsporangium hypogynous; vegetative cell usually capitellate; terminal cell with rounded apex; vegetative cells $12-15 \times 50-70\mu$; oogonium $50-55 \times 40-44\mu$; oospore $48-53 \times 36-42\mu$; androsporangium $12-13 \times 10-12\mu$.

United States: New York (Smithtown Pond and Fish Hatchery Pond, Long Island).

Distinguished by its depressed-globose oogonia with verticillate folds forming pointed projections at the suture. It is larger than *Oe. boreale* below and is further distinguished by the shape and form of the oogonium.

174. *Oe. boreale* Hirn.

(Pl. LVI, fig. 542.)

1900, p. 275, Pl. XLVII, fig. 300.

Dioecious, nannandrous; oogonium 1 (rarely 2), pyriform, wall thick and lamellose, medianly plicate, 16–19 verticillate folds, operculate, division supramedian; oospore subglobose (or subpyriform-globose), inflating oogonium, spore wall smooth; vegetative cell capitellate; dwarf male broadly obovoid, unicellular, on oogonium; vegetative cell $18-22 \times 55-150\mu$; oogonium $58-65 \times 67-78\mu$; oospore $48-53 \times 48-53\mu$; dwarf male $14-16 \times 15-20\mu$.

Finland.

Cf. *Oe. costatum*, above.

175. *Oe. mirandrium* Skuja.

(Pl. LIV, fig. 514.)

1927, p. 101, Pl. II, figs. 12 and 12a.

Dioecious, nannandrous, (?) idioandrosporous; oogonium 1 (rarely 2), subpyriform globose, operculate, division supramedian, or nearly superior; oospore globose, quite or not filling oogonium, spore wall smooth; dwarf male unicellular, ovoid, on oogonium; terminal cell obtuse; basal cell elongate; vegetative cell $13-20 \times 28-90\mu$; suffultory cell $14-24 \times 35-90\mu$; oogonium $40-43 \times 38-43\mu$; oospore $35-42 \times 35-42\mu$; dwarf male $9-26 \times 11-35\mu$.

Latvia.

Characterized by the supramedian operculum, unicellular dwarf male, and smaller suffultory cell. It bears some resemblance to the macrandrous *Oe. pringsheimii* (No. 95.)

176. *Oe. contortum* Hallas.

(Pl. XLVIII, fig. 464.)

1905, p. 399, fig. 2; Hirn 1906, p. 11, Pl. IV, fig. 25.

Dioecious, nannandrous, gynandrosporous; oogonium 1 (rarely 2), depressed-globose (rarely pyriform), operculate, division inframedian; oospore depressed (rarely globose), quite filling oogonium, spore wall smooth; androsporangium 1–5, curved, hypogynous or rarely epigynous or scattered; dwarf male obovoid, unicellular, on oogonium; vegetative cell $5-7 \times 20-60\mu$; oogonium $23-35 \times 16-35\mu$; oospore $16-28 \times 12-21\mu$; androsporangium $5-7 \times 7-14\mu$; dwarf male $4 \times 13\mu$.

Denmark.

The twisted appearance of the filaments, the inframedian operculum, and the non-filled oogonia are characteristic. It should be remarked that Hallas' original figure which Hirn copies gives every evidence of an inframedian "pore." The description however indicates an operculate oogonium. The curved filaments are similar to those of the monoecious *Oe. curvum* (No. 2). It should be compared with *Oe. platygynum* (No. 167).

177. *Oe. schmidlei* Gutwinski.

(Pl. LVIII, fig. 572.)

1896, p. 2, Pl. 1, fig. 2; Hirn 1900, p. 228, Pl. XXXVIII, fig. 234.

Dioecious, nannandrous; oogonium 1, obovoid-globose, operculate, division superior (? narrow); oospore globose, not filling oogonium, spore wall pitted, rather thick; dwarf male a little curved, on suffultory cell, antheridium exterior, 1?; vegetative cell $10 \times 25-90\mu$; suffultory cell $22 \times 44\mu$; oogonium $29 \times 32-37\mu$; oospore $26 \times 26\mu$; dwarf male stipe $5 \times 27\mu$; antheridium $5-6 \times 5-6\mu$.

Austria; British Columbia.

Gutwinski's figure leaves room for doubt as to the exact nature of the oospore wall. Hirn (1900) regards it as "membrana porifera" which I have interpreted as pitted. The record from British Columbia by Mr. G. H. Wailes gave no added data. In general appearance it resembles the larger and smooth spored *Oe. hians* (No. 180).

178. *Oe. longatum* Kuetzing.

(Pl. LVIII, fig. 563.)

1853, p. 11, Pl. 33, fig. 6; Wittr. 1874, p. 38; Nordst. 1877, p. 28; *Oe. crispulum* Wittr. and Nordst. in Wittr. 1872, p. 5; Hirn 1900, p. 239, Pl. XL, fig. 248; Heering 1914, p. 182, fig. 249; Tiffany 1926, p. 105, Pl. IX, fig. 106; P. B. A. No. 82.

Dioecious, nannandrous; oogonium 1-3, ovoid to ellipsoid, operculate, division superior; oospore ellipsoid about filling the oogonium, spore wall usually smooth, basal cell elongate; terminal cell obtuse; dwarf male on the oogonium; antheridium exterior, 1-?, curved; vegetative cells $4-7 \times 10-35\mu$; oogonium $16-18 \times 21-25\mu$; oospore $15-17 \times 17-19\mu$; dwarf male stipe $5-6 \times 10-15\mu$; antheridium $4-5 \times 5-6\mu$.

United States: Iowa, Massachusetts, Pennsylvania; Austria, England, Finland, Germany, Sweden; British Columbia.

Cf. *Oe. rugulosum*, below.

179. *Oe. rugulosum* Nordstedt.

(Pl. LVIII, fig. 560.)

1877, p. 28, Pl. 3, figs. 12 and 13; *Oe. sp.* Borge 1896, p. 5, Pl. 1, fig. 4; Hirn 1900, p. 241, Pl. XL, figs. 249 and 250; Heering 1914, p. 182, fig. 250; Transeau 1917, p. 231; Tiffany 1926, p. 105, Pl. IX, fig. 96.

Dioecious, nannandrous; oogonium 1-2, obovoid or obovoid-ellipsoid, operculate, division superior; oospore ellipsoid, nearly filling oogonium

(or rarely globose-ellipsoid, not filling oogonium) spore wall sometimes crenulate; dwarf male on or near oogonium; antheridium exterior, 1-?, curved; vegetative cell $4-8 \times 10-35\mu$; oogonium $16-20 \times 22-29\mu$; oospore $15-18 \times 19-23\mu$; antheridium $4-5 \times 5-6\mu$; dwarf male $5-7 \times 11-14\mu$.

United States: Iowa, Michigan, New York; Denmark, France, Sweden; Australia.

179a. Form **minutum** (Hansgirg) Hirn.

(Pl. LVIII, fig. 561.)

1900, p. 242, Pl. XL, fig. 251; *Oe. crispulum* Witttr. and Nordst. var. *minutum* Hansg. 1886, p. 44.

Smaller; median spore wall distinctly crenulate (Transeau); vegetative cell $3-7 \times 12-40\mu$; oogonium $14-18 \times 18-24\mu$; oospore $13-17 \times 15-20\mu$.

United States: Illinois; Austria.

179b. Form **rotundatum** Hirn.

(Pl. LVIII, fig. 562.)

1900, p. 242, Pl. XL, fig. 252.

Oogonium a little larger and shorter; oospore globose-ellipsoid; vegetative cell $5-8 \times 15-31\mu$; oogonium $19-23 \times 20-26\mu$; oospore $18-22 \times 18-23\mu$; dwarf male stipe $5-6 \times 11-15\mu$; antheridium $4-6 \times 4-6\mu$.

Germany.

This species is near *Oe. longatum* (No. 178) and is separable largely on the basis of the form of the oogonium and relative length of the oospore. Sometimes the oospore wall of *Oe. rugulosum* is crenulate. The latter is characteristic of f. *minutum*, which is smaller and has more obovoid oogonia. *F. rotundatum* has globose-ellipsoid oospores and slightly larger oogonia.

180. **Oe. hians** Nordstedt and Hirn.

(Pl. LVIII, figs. 570 and 571.)

In Hirn 1900, p. 227, Pl. XXXVIII, fig. 233.

Dioecious, nannandrous, gynandrosporous; oogonium 1-2, subovoid or globose, operculate, division superior, wide; oospore globose, quite filling oogonium, spore wall smooth, thick, and often lamellose; suffultory cell tumid; androsporangium 1-2, subepigynous; basal cell elongate; terminal cell apically obtuse; dwarf male curved, on suffultory cell, antheridium exterior, unicellular; vegetative cell $9-15 \times 37-145\mu$; suffultory cell $26-33 \times 40-80\mu$; oogonium $37-43 \times 45-60\mu$; oospore $33-40 \times 33-40\mu$; androsporangium $11-12 \times 15-18\mu$; dwarf male stipe $7-9 \times 32-35\mu$; antheridium $6-8 \times 5-6\mu$.

United States: Iowa; Brazil.

Characterized by its swollen suffultory cells, wide operculum, and tendency toward capitellate vegetative cells.

181. *Oe. hoersholmiense* Hallas.

(Pl. LVII, fig. 559.)

1905, p. 400, fig. 5; Hirn 1906, p. 15, Pl. IV, fig. 21.

Dioecious, nannandrous, gynandrosporous; oogonium 1 (rarely 2), ellipsoid, operculate, division superior; oospore ellipsoid or globose, not filling oogonium, spore wall smooth; androsporangium 1-5, often hypogynous or scattered; basal cell elongate; terminal cell obtuse; dwarf male curved, on oogonium, antheridium exterior, 1-2; vegetative cell $7-12 \times 44-160\mu$; oogonium $30-36 \times 56-68\mu$; oospore $26-32 \times 32-45\mu$; androsporangium $7-12 \times 16-21\mu$; dwarf male stipe $9-12 \times 30-35\mu$; antheridium $7-8 \times 10-11\mu$; basal cell $3-14 \times 16-50\mu$.

Denmark.

This species is perhaps near *Oe. ciliatum* (No. 185) from which it differs in its smaller size, longer vegetative cell, and obtuse terminal cell. The figure of Hallas would seem to indicate an operculum in a supreme position, although the description records "superior." I have included it in both positions in the key.

182. *Oe. laetevirens* Wittrock.

(Pl. LVII, fig. 558.)

In Zeller 1876, p. 427; Hirn 1900, p. 239, Pl. XL, fig. 247.

Dioecious, nannandrous; oogonium 1 (rarely 2), subobovoid-globose, operculate, division superior; oospore globose or subdepressed-globose, filling oogonium or enlarging it, spore wall smooth; dwarf male on oogonium, stipe a little curved, antheridium exterior, 1-?; vegetative cell $12-13 \times 18-32\mu$; oogonium $30-34 \times 32-36\mu$; oospore $28-32 \times 27-29\mu$; dwarf male stipe $7-8 \times 20-21\mu$; antheridium $6 \times 5-7\mu$.

Brazil.

Cf. *Oe. macrandrium* (No. 184).183. *Oe. pluviale* Nordstedt.

(Pl. XL, figs. 392 and 393.)

In Rabenh. Alg. Eur. No. 2257, 1871; Wittr. 1872, p. 7; *Oe. intermedium* Kuetz. in Rabenh. Alg. Sachs. No. 828, 1859; *Oe. montagnei* Fior.-Mazz. var. *submarinum* Wittr. in Wittr. and Nordst. Alg. Exs. No. 905, 1889; *Oe. fonticola* Al. Br. in Wolle 1887, p. 93, Pl. 75, figs. 4-6; (?) *Vesiculifera dissiliens* Hass. 1845, p. 202, Pl. 50, fig. 7; *Oe. diplandrium* Juranyi 1873, p. 27, Pl. 1-3; (?) *Oe. fonticola* Al. Br. var. *flavescens* Hansg. 1891, p. 305; *Oe. montagnei* Fior.-Mazz. var. *saxicolum* Wittr. 1876, p. 50, Pl. 13, figs. 29-31; Hirn 1900, p. 280, Pl. XLVIII, fig. 311; P. B. A. No. 1190.

Dioecious, nannandrous, idioandrosporous; oogonium 1 (very rarely 2-3), obovoid-globose or subglobose, operculate, division superior; oospore subglobose or subellipsoid-globose, almost filling oogonium, spore wall smooth; vegetative cell varying much in the same plant; basal cell elongate; terminal cell obtuse; androsporangium to 10-seriate; dwarf male broadly obovoid, unicellular, on oogonium; vegetative cell $22-29 \times 20-55\mu$; androsporangial vegetative cell $18-27 \times 18-27\mu$; oogonium $34-45 \times 34-50\mu$; oospore $32-40 \times 31-43\mu$; androsporangium $17-25 \times 6-13\mu$; dwarf male $10 \times 14-15\mu$.

United States: California; Austria, England, France, Spain, Germany, Hungary, Italy, Ireland, Switzerland, Sweden; Siam.

This species is the *Oe. diplandrium* Juranyi rather widely copied in morphological texts. It is characterized by its short vegetative cells, clustered unicellular dwarf males, and compact and numerous antheridia. It bears some resemblance to the imperfectly known *Oe. fonticola* (No. 208).

184. *Oe. macrandrium* Wittrock.

(Pl. LVII, fig. 549.)

1870, p. 130, Pl. 1, figs. 3-5; Hirn 1900, p. 233, Pl. XXXIX, fig. 240; Collins 1909, p. 258; Heering 1914, p. 180, fig. 248; Tiffany 1926, p. 106, Pl. IX, fig. 97.

Dioecious, nannandrous; oogonium 1-4, globose-ovoid, operculate, division superior; oospore globose, rarely ovoid-globose, not completely filling the oogonium, spore wall smooth; terminal cell obtuse or very shortly apiculate; dwarf male on or near the oogonium, stipe much curved, sometimes 2-3 celled; antheridium 1-7; vegetative cells $15-20 \times 45-100\mu$; oogonium $36-42 \times 43-54\mu$; oospore $31-37 \times 33-39\mu$; dwarf male stipe $12-13 \times 24-33\mu$; antheridium $9-10 \times 7-10\mu$.

United States: Ohio, Iowa, Massachusetts, Pennsylvania; England, Finland, France, Germany, Denmark, Norway, Sweden, Switzerland.

184a. Form *acuminatum* Hirn.

(Pl. LVII, fig. 550.)

1900, p. 234, Pl. XXXIX, fig. 241.

Dwarf male apically broadly acute; vegetative cell $15-19 \times 30-80\mu$; oogonium $37-45 \times 43-55\mu$; oospore $35-40 \times 36-45\mu$; dwarf male stipe $11-15 \times 24-35\mu$; antheridium $9-13 \times 9-13\mu$.

Sweden.

184b. Form *aemulans* Hirn.

(Pl. LVII, fig. 551.)

1906, p. 43; *Oe. macrandrium* Wittr. var. *aemulans* Hirn 1900, p. 235, Pl. XXXIX, fig. 242; (?) *Oe. lundense* Wittr. in Wolle 1887, p. 79, Pl. 77, figs. 9 and 10.

Vegetative cell $10-15 \times 22-90\mu$; oogonium $28-40 \times 33-45\mu$; oospore $26-36 \times 26-36\mu$; dwarf male stipe $9-12 \times 20-25\mu$; antheridium $8-9 \times 8-10\mu$.

United States: Pennsylvania, California, Michigan, Illinois, Mississippi; Brazil.

184c. Form *lundense* (Wittrock) Hirn.

(Pl. LVII, fig. 552.)

1906, p. 42, Pl. III, fig. 19; *Oe. lundense* Wittr. 1872, p. 4, and 1874, p. 23; *Oe. lundense* Wittr. in Hirn 1900, p. 237, Pl. XL, fig. 245; *Oe. fioniae* Hallas 1905, p. 401, fig. 6.

Oogonium subobovoid-globose; oospore wall sometimes thick; vegetative cell $13-21 \times 13-50\mu$; oogonium $31-40 (-45) \times 34-43\mu$; oospore

29-36 (-40) $\times$ 29-36 (-40) μ ; dwarf male stipe 10-15 $\times$ (21-) 24-32 μ ; antheridium 8-10 $\times$ 6-9 μ .

Finland, France, Denmark, Sweden.

184d. Var. **propinquum** (Wittrock) Hirn.

(Pl. LVII, fig. 553.)

1906, p. 42, Pl. IV, fig. 20; *Oe. propinquum* Wittr. 1870, p. 129, and in Hirn 1900, p. 236, Pl. XL, fig. 243; *Oe. eremitum* Hallas 1905, p. 403, fig. 8.

Smaller; oogonium 1-3; vegetative cell 8-14 $\times$ 18-70 μ ; oogonium 28-36 $\times$ 28-45 μ ; oospore 24-31 $\times$ 24-31 μ ; dwarf male stipe 11-14 $\times$ 20-35 μ ; antheridium 7-9 (-12) $\times$ 6-8 μ .

United States: Illinois, Alabama; Austria, France, Sweden, Denmark, Latvia.

184e. Var. **hohenackerii** (Wittrock) Tiffany.

(Pl. LVII, figs. 554-556.)

1929, p. 75; *Oe. hohenackerii* Wittr. 1874, p. 23, and in Hirn 1900, p. 238, Pl. XL, fig. 246; *Oe. tumidulum* in Hohenacker Alg. Exs. No. 404.

Vegetative cell 12-15 $\times$ 24-45 μ ; oogonium 29-33 $\times$ 30-35 μ ; oospore 27-31 $\times$ 28-31 μ ; androsporangium 11-13 $\times$ 10-14 μ ; dwarf male stipe 9-14 $\times$ 18-24 μ ; antheridium 5-6 $\times$ 5-8 μ .

United States: Michigan, Ohio; India.

184f. Var. **scrobiculatum** Ackley.

(Pl. LVII, fig. 557.)

1929, p. 305, Pl. XXXVI, figs. 17-19.

Oogonium globose to subglobose; median layer of oospore wall scrobiculate; vegetative cell 9-11 $\times$ 27-60 μ ; oogonium 32-35 $\times$ 30-35 μ ; oospore 31-33 $\times$ 31-32 μ ; dwarf male stipe 12 $\times$ 10-18 μ ; antheridium 8 $\times$ 6-10 μ .

United States: Michigan (Ox-bow of river near Holland).

The *macrandrium* group is perhaps one of the most interesting and variable in the genus. The species has rather an unfortunate name to be nannandrous. The group is readily recognized by its frequently seriate oogonia, its curved and twisted dwarf males resting on or near the oogonia and frequently clustered, and its relatively short vegetative cells. *Oe. lactevirens* (No. 182) is separable from the group largely on its smaller size. *F. lundense* has very short vegetative cells, and those of var. *hohenackerii* are slightly longer. Var. *scrobiculatum* is the only member of the group with scrobiculate oospores. The remaining forms and varieties are separable largely on the basis of size.

185. *Oe. ciliatum* (Hassall) Pringsheim.

(Pl. LIX, fig. 583.)

Vesiculifera ciliata Hass. 1845, p. 202, Pl. 52, fig. 2; *Oe. ciliatum* Pringsh. 1856, p. 227, Pl. I, figs. 1-10; Pringsh. 1858, p. 70, Pl. 4, figs. 1-14, and Pl. 5, fig. 8; *Oe. piliferum* Auerswald in Rabenh. Alg. Sachs. 1855; Hirn 1900, p. 243, Pl. XLI, fig. 253; Heering 1914, p. 183, fig. 252; Tiffany 1926, p. 106, Pl. IX, figs. 98 and 99.

Dioecious, nannandrous, gynandrosporous; oogonium 1-7, ovoid, to ovoid-ellipsoid (rarely nearly ellipsoid), operculate, division superior or nearly supreme; oospore ovoid to subellipsoid, rarely globose, nearly filling oogonium, spore wall smooth; androsporangium to 8-seriate, often subepigynous; basal cell elongate; terminal cell setiform; dwarf male curved, on oogonium; antheridium exterior, 1; vegetative cell $14-24 \times 35-92\mu$; oogonium $43-50 \times 55-72\mu$; oospore $40-47 \times 44-57\mu$; androsporangium $14-20 \times 10-20\mu$; dwarf male stipe $10-15 \times 24-31\mu$; antheridium $8-10 \times 10-11\mu$.

United States: Iowa, Ohio; England, Finland, Germany, Spain, Sweden, Latvia; British Columbia.

Characterized by its setiform terminal cells and ovoid to ovoid-ellipsoid oogonia. The division of the oogonium is quite often practically supreme rather than superior. It is near *Oe. wabashense* (No. 192); the latter has capitellate vegetative cells, supreme operculum, and smaller oogonia.

186. *Oe. spectabile* Hirn.

(Pl. XLI, figs. 403-405.)

1900, p. 284, Pl. XLIX, fig. 317.

Dioecious, nannandrous, idioandrosporous; oogonium 1, obovoid-ellipsoid or obovoid, operculate, division superior; oospore ellipsoid, nearly filling oogonium, spore wall smooth; androsporangium to 16-seriate, the androsporangial part of the filament often curved; basal cell elongate; dwarf male broadly obovoid, unicellular, on oogonium; vegetative cell $20-32 \times 25-90\mu$; oogonium $42-50 \times 63-72\mu$; oospore $40-48 \times 55-62\mu$; androsporangium $19-27 \times 11-19\mu$; dwarf male $11-14 \times 16-21\mu$.

United States: Mississippi; Australia.

Characterized by the wide variation in cell diameter, the unicellular dwarf male, and the pronounced curve of the terminal part of the androsporangial filament. It is similar to *Oe. implexum* (below) which is smaller and has relatively more slender vegetative cells.

187. *Oe. implexum* Hirn.

(Pl. XLI, figs. 400 and 401.)

1900, p. 283, Pl. XLIX, fig. 316.

Dioecious, nannandrous, gynandrosporous; oogonium 1-3, subobovoid-ellipsoid, operculate, division superior; oospore ellipsoid, filling oogonium, spore wall smooth; androsporangium to 20-seriate

(filament containing androsporangia curved); dwarf male broadly obovoid, unicellular, on oogonium; vegetative cell $16-20 \times 33-100\mu$; oogonium $38-45 \times 50-75\mu$; oospore $36-42 \times 47-60\mu$; androsporangium $16-20 \times 13-25\mu$; dwarf male $12-14 \times 13-15\mu$.

Australia.

Cf. *Oe. spectabile*, above.

188. *Oe. monile* Berkeley and Harvey.

(Pl. LVIII, figs. 564 and 565.)

In Hooker 1860, p. 342, Pl. 196, fig. B; Wittr. 1874, p. 40; (?) *Oe. birmanicum* Wittr. in Moebius 1892, p. 430, fig. 9B; Hirn 1900, p. 229, Pl. XXXVIII and XXXIX, fig. 235.

Dioecious, nannandrous; oogonium 1-8, subovoid or subglobose, operculate, division superior, very broad; oospore globose or subglobose, quite filling oogonium (except in length), spore wall of three layers: outer layer smooth, middle layer scrobiculate, inner layer smooth; vegetative cell sometimes capitellate; basal cell elongate; terminal cell obtuse; dwarf male a little curved, near oogonium, antheridium exterior, 1-?; vegetative cell $9-15 \times 50-160\mu$; suffultory cell $21-29 \times 40-57\mu$; oogonium $30-39 \times 30-56\mu$; oospore $28-38 \times 28-38\mu$; dwarf male stipe $9 \times 25\mu$; antheridium $7 \times 7\mu$.

United States: Massachusetts, Michigan; Australia; Brazil.

188a. Form *borgei* Hirn.

(Pl. LVIII, fig. 567.)

1900, p. 230, Pl. XXXIX, fig. 237.

Vegetative cell $14-20 \times 45-100\mu$; suffultory cell $35-38 \times 45-74\mu$; oogonium $40-50 \times 40-60\mu$; oospore $39-45 \times 39-48\mu$; dwarf male stipe $13-14 \times 36-40\mu$; antheridium $9-11 \times 8-10\mu$.

Australia.

188b. Var. *eminens* Hirn.

(Pl. LVIII, figs. 568 and 569.)

1900, p. 231, Pl. XXXIX, fig. 238.

Much larger; gynandrosporous, androsporangium subepigynous; dwarf male on suffultory cell, antheridium 1-2; vegetative cell $18-25 \times 65-150\mu$; suffultory cell $38-41 \times 48-72\mu$; oogonium $56-68 \times 55-74\mu$; oospore $53-60 \times 50-58\mu$; androsporangium $19-23 \times 15-18\mu$; dwarf male stipe $18-19 \times 40-48\mu$; antheridium $10-13 \times 12-14\mu$.

Australia.

188c. Form *victoriense* G. S. West.

(Pl. LVIII, fig. 566.)

1909, p. 45, fig. 6 A and B.

Dwarf male a little longer and more narrow, on suffultory cell; vegetative cell $12-14 \times 60-96\mu$; suffultory cell $27-28 \times 68-83\mu$; oogonium $42-45 \times 42-52\mu$; oospore $40-42 \times 40-42\mu$; dwarf male stipe $6-8 \times 38-44\mu$.

Victoria, Africa.

The species is characterized by its scrobiculate oospore, wide operculum, enlarged suffultory cell, and usually many-seriate oogonia. Var. *eminens* is much larger and f. *borgei* is considerably larger than the type; f. *victoriense* is near f. *borgei*, differing in the size of the oospore.

189. *Oe. obtruncatum* Wittrock.

(Pl. LIX, fig. 577.)

1874, p. 41; Hirn 1900, p. 284, Pl. XLIX, fig. 318.

Dioecious, nannandrous, gynandrosporous; oogonium 1-6, ellipsoid, or globose-ellipsoid, operculate, division supreme, operculum small and deciduous; oospore of same form as oogonium, nearly filling it, spore wall smooth; vegetative cell evidently capitellate; basal cell elongate; terminal cell (often the oogonium) apically obtuse; dwarf male oblong-pyriform, curved, unicellular, on oogonium; vegetative cell $18-22 \times 56-110\mu$; oogonium $45-55 \times 56-68\mu$; oospore $43-53 \times 52-66\mu$; androsporangium $20 \times 24\mu$; dwarf male stipe $17-20 \times 36-40\mu$; antheridium $6-8 \times 4-6\mu$.

United States: Ohio, Illinois; England; Brazil; India; Australia.

189a. Var. *completum* Hirn.

(Pl. LIX, figs. 579-582.)

1900, p. 285, Pl. L, fig. 319.

Oogonium larger; androsporangium 1-5, hypogynous or subepigynous or epigynous; terminal cell sometimes setiform; vegetative cell $18-22 \times 36-150\mu$; oogonium $55-58 \times 63-75\mu$; oospore $53-56 \times 61-73\mu$; androsporangium $20-22 \times 19-22\mu$.

Bengal.

189b. Var. *ellipsoideum* Wittrock.

(Pl. LIX, fig. 578.)

1878, p. 141; Hirn 1900, p. 286, Pl. L, fig. 320.

Oogonium longer, ellipsoid, 1-2, terminal; vegetative cell $17-23 \times 54-110\mu$; oogonium $42-54 \times 66-75\mu$.

Venezuela.

Oe. obtruncatum is characterized by the usually capitellate vegetative cells, supreme position of the operculum, unicellular nannandrium, and gynandrosporous habit. The two varieties *completum* and *ellipsoideum* are separable largely on the basis of differently sized oogonia. The species is much larger than the gynandrosporous *Oe. wabashense* (No. 192). It is perhaps nearest in general appearance to *Oe. praticolum* (No. 190) and *Oe. supremum* (No. 191), both of which are idioandrosporous and have larger oogonia.

190. *Oe. praticolum* Transeau.

(Pl. LX, figs. 586-588.)

1914, p. 298, Pl. XXIX, figs. 1-5; Tiffany 1926, p. 107, Pl. VII, figs. 82-84.

Dioecious, nannandrous, idioandrosporous; oogonium 1-7, ellipsoid to globose-ellipsoid, often terminal, sometimes scattered; wall sometimes rather thick, operculate, division supreme, lid small and deciduous; oospore ellipsoid to globose-ellipsoid, quite filling the oogonium, wall smooth, androsporangium 1-20; vegetative cell slightly capitellate; basal cell usually elongate; terminal cell apiculate or extended into a long hyaline seta; antheridium exterior, 1; vegetative cell, female 16-26×65-130 μ ; vegetative cell of androsporangial filament 14-22×56-110 μ ; oogonium 48-60×62-74 μ ; oospore 46-58×60-72 μ ; androsporangium 20-24×18-22 μ ; dwarf male stipe 8-14×21-28 μ ; antheridium 6-10×7-12 μ .

United States: Illinois, Ohio, Iowa.

Cf. *Oe. obtruncatum* (No. 189).191. *Oe. supremum* Tiffany.

(Pl. LIX, figs. 573-576.)

1924, p. 185, Pl. I, figs. 3 and 4, Pl. II, figs. 4 and 5; 1926, p. 107, Pl. VIII, figs. 90 and 91.

Dioecious, nannandrous, idioandrosporous; oogonium 1-4, globose or ellipsoid-globose, often terminal, operculate, division at the upper extremity of oogonium, lid often deciduous; oospore globose or ellipsoid-globose, filling oogonium, spore wall smooth; androsporangium 1-6; vegetative cell distinctly capitellate; basal cell elongate; dwarf male broadly obovoid, unicellular, on oogonium; vegetative cell 24-32×60-132 μ ; oogonium 66-78×72-90 μ ; oospore 60-66×66-84 μ ; androsporangium 26-28×30-40 μ ; dwarf male 20-24×24-26 μ ; basal cell 28-32×90-110 μ .

United States: Iowa.

Cf. *Oe. obtruncatum* (No. 189).192. *Oe. wabashense* Tiffany.

(Pl. LX, figs. 584 and 585.)

1927, p. 203, Pl. IX, figs. 3-5.

Dioecious, nannandrous, gynandrosporous; oogonium 2-3, ellipsoid or ovoid, often terminal, operculate, division supreme, lid often deciduous; oospore ellipsoid or ovoid (rarely globose), filling oogonium or not, spore wall smooth; androsporangium 1-3, subepigynous; vegetative cell capitellate; basal cell elongate; dwarf male on oogonium; antheridium 1-?; vegetative cell 12-20×36-64 μ ; oogonium 36-42×44-60 μ ; oospore 34-38×40-55 μ ; androsporangium 12-16×10-20 μ ; dwarf male stipe 12-16×24-40 μ ; antheridium 7-12×6-10 μ ; basal cell 18-20×40-64 μ .

United States: Indiana, Ohio.

Cf. *Oe. ciliatum* (No. 185) and *Oe. obtruncatum* (No. 189).

193. *Oe. tentoriale* Nordstedt and Hirn.

(Pl. LXI, fig. 597.)

In Hirn 1900, p. 248, Pl. XLII, fig. 260.

Dioecious, nannandrous; oogonium 1, terminal, broadly ellipsoid or obovoid-globose, division supreme, lid small, deciduous, oogonium wall longitudinally ribbed on the inner surface, ribs often anastomosing; oospore quite filling oogonium, outer layer of the spore wall with 40–45 longitudinal, very finely crenulate ribs, closely filling in between the ribs of the oogonium, connected by delicate transverse lines, inner layer smooth; basal cell elongate; terminal cell obtuse; dwarf male suberect, on suffultory cell, antheridium exterior 1–?; vegetative cell 20–33 (–37) $\times$ 60–225 μ ; oogonium 66–82 $\times$ 73–88 μ ; dwarf male stipe 10–15 $\times$ 37–48 μ ; antheridium 9–12 $\times$ 7 μ .

United States: Illinois; Brazil.

Cf. *Oe. acrosporum* (No. 195).194. *Oe. michiganense* Tiffany.

(Pl. LXI, figs. 594–596.)

1927, p. 205, Pl. IX, figs. 10–12.

Dioecious, nannandrous, gynandrosporous; oogonium 1–7, globose to ellipsoid-globose (rarely subglobose), operculate, division at the upper extremity of the oogonium; oospore globose, filling the oogonium or not, outer layer of spore wall smooth, middle layer with 12–24 crenulate, sometimes anastomosing, longitudinal ribs, inner layer smooth; suffultory cell enlarged; vegetative cell broadly capitellate; basal cell elongate; terminal cell, occasionally oogonium, apically obtuse; dwarf male, a little curved, on suffultory cell, antheridium interior; vegetative cell 12–24 $\times$ 80–160 μ ; suffultory cell 32–48 $\times$ 64–80 μ ; oogonium 50–64 $\times$ 50–80 μ ; oospores 44–60 $\times$ 44–60 μ ; androsporangium 16–20 $\times$ 16–20 μ ; dwarf male 14–20 $\times$ 40–56 μ ; basal cell 18–20 $\times$ 70–100 μ .

United States: Michigan.

The only nannandrous member of the genus possessing a capitellate vegetative cell, a supreme operculum, and longitudinally ribbed oospores.

195. *Oe. acrosporum* De Bary.

(Pl. LX, fig. 589.)

1854, p. 47 *et seq.*, Pl. 3, figs. 1–12; *Oe. acrosporum* De Bary f. *connectens* Wittr. *in* Wittr. and Nordst. Exs. No. 1101, 1893, *in* P. B. A. No. 409, 1898, *in* Hirn 1900, p. 245, Pl. XLI, fig. 255; Hirn 1900, p. 244, Pl. XLI, fig. 254; Collins 1909, p. 259; Heering 1914, p. 183, fig. 253; Tiffany 1926, p. 107, Pl. IX, fig. 103.

Dioecious, nannandrous, gynandrosporous (or idioandrosporous); oogonium 1, terminal, ellipsoid, operculate, division supreme, operculum small and deciduous; wall of oogonium with longitudinal, sometimes anastomosing, ridges on the inner surface; oospore quite filling the oogonium, outer layer of spore wall with 23–30 longitudinal ridges, closely fitting in between the ridges of the oogonium, inner smooth; basal cell elongate; terminal cell obtuse; androsporangia 1–2, hypo-

gynous; dwarf male curved, on the suffultory cell, stipe sometimes 2-3 celled, upper cells long, antheridium exterior, 1-2; vegetative cell $12-21 \times 40-125\mu$; suffultory cell $16-25 \times 26-76\mu$; oogonium $35-48 \times 50-63\mu$; androsporangium $16-21 \times 12-15\mu$; dwarf male stipe, upper cell $6-8 \times 55-71\mu$; lower cell $9-12 \times 30-38\mu$; antheridium $6-8 \times 9-15\mu$

United States: Ohio, Illinois, Iowa, Michigan, New York, Massachusetts; Austria, England, Finland, France, Germany, Norway, Sweden; Brazil; Manchuria; Madagascar; British Columbia.

195a. Form **boreale** (Wolle) Hirn.

(Pl. LX, fig. 590.)

1900, p. 245, Pl. XLI, fig. 256; *Oe. acrosporum* De Bary var. *boreale* Wolle 1887, p. 84, Pl. 79, figs. 10 and 11.

Filaments few-celled; vegetative cell $14-16 \times 45-80\mu$.

United States: Pennsylvania, New Jersey.

It is very doubtful if this form of the genus is tenable. The plant described by Wolle seems to have had fewer cells to the filament than the species proper. From data at hand regarding other species, it appears this variability in the number of the cells is largely a matter of environmental conditions. There are some species in the genus, however, that are never found except with very few cells to a filament. The form is retained until more definite data can be secured regarding its variations. If it is normally few celled, Wolle's varietal position must be maintained.

195b. Var. **bathmidosporum** (Nordstedt) Hirn.

(Pl. LX, fig. 592.)

1900, p. 246, Pl. XLII, fig. 259; *Oe. bathmidosporum* Nordst. 1878, p. 179; *Oe. acrosporum* De Bary in Gutw. 1897, p. 8; *Oe. acrosporum* De Bary in P. B. A. No. 163, 1896.

Smaller; ribs fewer, 11-17, evidently crenulate; dwarf male stipe unicellular; vegetative cell $12-17 \times 35-125\mu$; suffultory cell $15-22 \times 40-110\mu$; oogonium $30-40 \times 40-54\mu$; dwarf male stipe $9-11 \times 30-34\mu$; antheridium $8-10 \times 9-12\mu$.

United States: Massachusetts; Austria, Sweden; Brazil.

195c. Var. **floridense** Wolle.

(Pl. LX, fig. 593.)

1887, p. 83, Pl. 85, figs. 1 and 2; Hirn 1900, p. 246, Pl. XLI, fig. 258.

Smaller with elongate vegetative cell and more tumid suffultory cell; dwarf male stipe 2-3 celled; vegetative cell $7-8 \times 36-85\mu$; oogonium $33-35 \times 45-50\mu$.

United States: Florida.

The only record is that of Wolle.

195d. Var. **majusculum** Nordstedt.

(Pl. LX, fig. 591.)

1878a, p. 21, Pl. 2, figs. 17-18; Hirn 1900, p. 246, Pl. XLI, fig. 257.

Smaller; vegetative cell $14-21 \times 50-165\mu$; suffultory cell $18-28 \times 42-65\mu$; oogonium $44-56 \times 54-70\mu$.

Australia.

Oe. acrosporum is readily identified by its single terminal oogonium and by the dovetailing of the longitudinal ridges of the inner surface of the oogonium and outer surface of the oospore. The varieties and form (see note under each) are separable largely on the basis of number of ribs and size of cells. The species is easily distinguished from *Oe. tentoriale* (No. 193) by its smaller number of longitudinal ribs on the oospore, by its more slender cells, and differently shaped oogonia.

SPECIES IMPERFECTLY KNOWN.

The following members of the genus are recorded with varying degrees of certainty regarding their validity. A few are unquestionably readily recognized even though a complete description is not available or possible at the present status of our knowledge. Some are very near other species, differing in one or more characters, but lacking sufficient completeness of description to warrant definite placement in the genus. Others are included purely as a matter of record. Wherever possible, the species is included in the key in its probably proper position, with the idea of calling it to the attention of the student of the genus and thereby aid in its identification if collected. Appropriate remarks occur in connection with the description of each species.

196. *Oe. tapeinosporum* Wittrock.

(Pl. XXXIV, fig. 314.)

1874, p. 36; 1878, p. 140; De Toni 1889, p. 76; *Oe. tapeinosporum* var. *Angolense* West and West 1897, p. 5; Hirn 1900, p. 297, Pl. XXIII, fig. 117; Heering 1914, p. 224, fig. 344; Tiffany 1926, p. 96, Pl. VII, fig. 81.

Oogonium 1-2, depressed-globose, operculate, division median, distinct; oospore depressed-globose, not (rarely nearly) filling oogonium, spore wall smooth; basal cell subhemispherical; terminal cell obtuse; vegetative cell (2-) $3-5 \times 10-40\mu$; oogonium $15-19 \times (14-) 17-23\mu$; oospore $13-16 \times 8-14\mu$; basal cell $12-14 \times 5-7\mu$.

United States: Iowa; England, Germany; Columbia; Africa; India.

This species is rather readily distinguished by its conspicuous operculum and small size. No antheridia have been observed thus far. The Iowa material combined the characters of the type and the variety *Angolense*, and the latter thus becomes untenable. The species is perhaps dioecious, macrandrous.

197. *Oe. angustissimum* West and West.

(Pl. XXV, fig. 220.)

1897, p. 6; Hirn 1900, p. 314, Pl. L, fig. 321.

(?) Monoecious; oogonium 2, transversely inflated; oospore transversely elliptical and inflating oogonium, wall smooth; filament irregularly flexed and very narrow; vegetative cell $1.8-2 \times 13-28\mu$; oogonium $9.5 \times 10.5-14.5\mu$; oospore $9.5 \times 6.5\mu$.

Africa.

As far as is known, this is the smallest species of *Oedogonium*. Cf. *Oe. inconspicuum* (No. 198) and *Oe. tapeinosporum* (No. 196).

198. *Oe. inconspicuum* Hirn.

(Pl. XXXIV, fig. 313.)

1895, p. 23, Pl. 1, fig. 8; 1900, p. 296, Pl. XXIII, fig. 116; (?) *Oe. minutissimum* Grun. in Hansgirg 1905, p. 436.

(? Dioecious, macrandrous) Oogonium 1 (rarely 2-3), depressed- or subpyriform-globose, operculate, division median, narrow; oospore depressed-globose, filling the inflated part of oogonium, spore wall smooth; vegetative cell $3-5 \times 20-34\mu$; oogonium $13-18 \times (13-) 17-23\mu$; oospore $12-17 \times 8-12\mu$.

Columbia; Finland, Austria, Sweden.

In common with most of the smaller sized species of the genus, *Oe. inconspicuum* is found attached to other algæ or to submerged macrophytes. Cf. *Oe. angustissimum* above and *Oe. tapeinosporum* (No. 196).

199. *Oe. tenuissimum* Hansgirg.

1888a, p. 398; 1888, p. 222; Hirn 1900, p. 315.

(?) Monoecious; filament irregularly curved; oogonium 1, subpyriform, pore median; oospore globose-ellipsoid, not filling oogonium; vegetative cell $2.3-6 \times 9-25\mu$; oogonium $9-18 \times 14-23\mu$; oospore 15μ in diameter.

Austria.

Characterized by its median pore and elongated oogonium; cf. *Oe. sexangulare* (No. 135) and *Oe. pusillum* (No. 200).

200. *Oe. pusillum* Kirchner.

(Pl. XXXIV, fig. 316.)

1878, p. 59; (?) *Oe. excisum* Wittr. and Lund. in Nordst. 1880, p. 13; *Oe. africanum* Lagerh. 1893, p. 155; *Oe. Klebahnii* Lemm. 1893, p. 509, and 1895, p. 28, figs. 4 and 5; *ibid.* in De Wildemann 1896, p. 66, Pl. 5, figs. 2-7; *Oe. africanum* Lagerh. in Lemm. 1896, p. 502, Pl. 5, figs. 3-20; Hirn 1900, p. 299, Pl. XXIV, fig. 125; Collins 1912, p. 88; Heering 1914, p. 224, fig. 345; Tiffany 1926, p. 95, Pl. VII, fig. 80.

Oogonium 1 (rarely 2), subbiconic-ellipsoid or subbiconic-globose, seen from above circular, margin even, operculate, division wide; oospore ellipsoid or globose, generally constricted at the middle, not quite filling oogonium, spore wall smooth; basal cell subhemispherical; terminal cell obtuse or obtusely conical; vegetative cell $3-6 \times 10-60\mu$; oogonium $14-16 \times 15-25\mu$; oospore $11-13 \times 13-15\mu$; basal cell $7-8 \times 7-8\mu$.

United States: Iowa, Illinois; Austria, France, Germany; Africa; Brazil; Sumatra.

This species is usually very readily recognized by its medianly constricted oospore, and whether eventually found to be monoecious or dioecious, it is a distinct species.

201. *Oe. selandiae* Hallas.

(Pl. XXXVIII, fig. 359.)

1905, p. 496, fig. 14; Hirn 1906, p. 20, Pl. III, fig. 11.

(?) Monoecious; oogonium 1-2, oblong (or angular-oblong), operculate, division superior; oospore oblong-ellipsoid or obovoid, completely filling oogonium or not; vegetative cell $3-5 \times 13-40\mu$; oogonium $9-14 \times 27-30\mu$; oospore $7-12 \times 16-25\mu$.

Denmark.

To be compared with *Oe. gracillimum* (No. 114) and with *Oe. rugulosum* (No. 179).

202. *Oe. sancti thomæ* Wittrock and Cleve.

(Pl. XXXVIII, fig. 360.)

In Witt. 1874, p. 40; 1878, p. 141; Hirn 1900, p. 304, Pl. XXIX, fig. 173; Collins 1909, p. 265.

Oogonium 1-3, pyriform, operculate, division superior; oospore pyriform-obovoid, not quite filling oogonium, spore wall smooth; basal cell subhemispherical; terminal cell very slender, subhyaline; vegetative cell $7-15 \times 16-88\mu$; oogonium $28-33 \times 36-50\mu$; oospore $25-30 \times 28-35\mu$; basal cell $14-23 \times 8-12\mu$.

United States: Mississippi, Ohio; West Indies (St. Thomas).

Material from Mississippi and Ohio have been referred to this species, although antheridia were not observed. It is perhaps dioecious, macrandrous.

203. *Oe. fusus* Hallas.

(Pl. XXXIV, fig. 311.)

1905, p. 407, fig. 15; Hirn 1906, p. 14, Pl. IV, fig. 26.

(?) Dioecious; oogonium 1, ellipsoid, operculate, division median; oospore globose, not filling oogonium, spore wall smooth; vegetative cell $2-3 \times 10-23\mu$; oogonium $14-15 \times 31-35\mu$; oospore $12 \times 12\mu$.

Denmark.

Characterized by its small size and ellipsoid oogonium with globose oospore. Cf. *Oe. inconspicuum* (No. 198) and *Oe. pusillum* (No. 200).

204. *Oe. rhodosporum* (Welwitsch) Wittrock.

(Pl. XL, figs. 388 and 389.)

1874, p. 34; De Toni 1889, p. 72; Hirn 1900, p. 305, Pl. XXXI, fig. 193.

Dioecious (? macrandrous or nannandrous); oogonium 1-3, obovoid to globose obovoid, operculate, division superior; oospore same form as oogonium, nearly or quite filling oogonium, spore wall smooth; antheridium (? androsporangium) 1-6; vegetative cell $16-23 \times 20-66\mu$; oogonium $35-44 \times 45-54\mu$; oospore $33-41 \times 38-48\mu$; antheridium (? androsporangium) $16-19 \times 9-17\mu$.

France, Portugal.

To be compared with *Oe. pluviale* (No. 183) and with the two following species.

205. *Oe. vesicatum* (Lyngbye) Wittrock.

(Pl. XL, figs. 386 and 387.)

Conferva vesicata Lyng. 1819, p. 140, Pl. 47, fig. D1; *Oe. vesicatum* Wittrock 1874, p. 39; Hirn 1900, p. 306, Pl. XXXI, fig. 194.

Dioecious, oogonium 1, obovoid or globose-obovoid, operculate, division superior; oospore same form as oogonium which it nearly fills, spore wall smooth; antheridium (? androsporangium) 1-4; vegetative cell, female $17-23 \times 22-74\mu$; male (? androsporangial) $16-21 \times 28-63\mu$; oogonium $40-45 \times 49-60\mu$; oospore $35-42 \times 40-48\mu$; antheridium (? androsporangium) $16-18 \times 11-16\mu$.

Denmark.

Cf. *Oe. rhodosporum* above and *Oe. montagnei* below.

206. *Oe. montagnei* Fiorini-Mazzanti.

(Pl. XL, figs. 390 and 391.)

1860, p. 259, Pl. 1, figs. 1, 2 and 5; Wittrock 1874, p. 41; Hirn 1900, p. 307, Pl. XXXI, fig. 195.

Oogonium 1-2, obovoid or globose-obovoid, operculate, division superior; oospore of same form as oogonium, filling it or nearly so, spore wall smooth; vegetative cell $18-26 \times 20-110\mu$; oogonium $40-52 \times 48-65\mu$; oospore $38-47 \times 43-52\mu$.

Italy.

Cf. *Oe. rhodosporum* (No. 204) and *Oe. vesicatum* (No. 205).

207. *Oe. flexuosum* Hirn.

(Pl. XL, fig. 394.)

1900, p. 313, Pl. XLVIII, fig. 310.

Oogonium 1, obovoid or obovoid-globose, operculate, division superior; oospore same form as oogonium and filling it, spore wall smooth; filament irregularly curved; vegetative cell $15-18 \times 12-32\mu$; oogonium $29-34 \times 29-43\mu$; oospore $27-32 \times 27-37\mu$.

Ireland.

Characterized by its curved filaments and short cells. Cf. *Oe. fonticola* (No. 208) and *Oe. implexum* (No. 187).

208. *Oe. fonticola* Al. Braun.

(Pl. XLI, fig. 402.)

In Kuetz. 1849, p. 368, and 1858, p. 13, Pl. 40, fig. 2; *Oe. rhodosporum* (Welw.) Wittr. 1876, p. 47; Hirn 1900, p. 313, Pl. XLIX, fig. 315; Heering 1914, p. 225, fig. 341; Tiffany 1926, p. 106.

Dioecious; oogonium 1-2, obovoid or globose-obovoid, operculate, division superior; oospore obovoid-ellipsoid or subglobose, nearly or quite filling oogonium, spore wall smooth; vegetative cell $16-32 \times 12-70\mu$; oogonium $36-43$ (-45) $\times 43-56\mu$; oospore $34-41$ (-43) $\times 40-49\mu$.

United States: Iowa, Indiana, Mississippi; Austria, Italy, Spain, Germany, England.

Cf. *Oe. implexum* (No. 187), *Oe. flexuosum* (No. 207), and *Oe. pluviale* (No. 183).

West (1912) records the presence of very small apparently male filaments of *Oe. fonticola*, indicating a condition intermediate between "truly dioecious macrandrous species and those with large nannandria." If these are really male filaments, the species becomes dioecious, macrandrous. Until the data are more complete, *Oe. fonticola* must remain in the imperfectly known division of the genus.

209. *Oe. oryzæ* Wittrock.

(Pl. XXI, fig. 190.)

1876, p. 51, Pl. 13, figs. 32 and 33; Hirn 1900, p. 294, Pl. XXII, fig. 113.

Oogonium 1-2, scarcely tumid, subobovoid or subcylindric, pore superior; oospore of same form as oogonium, nearly filling it or sometimes enlarging it, spore wall smooth; basal cell elongate; terminal cell acuminate; vegetative cell $24-39 \times 36-120\mu$; suffultory cell $45 \times 56-90\mu$; single or upper series of oogonium $45-55 \times 65-95\mu$; lower series of oogonium $43-53 \times 45-57\mu$; oospore $41-51 \times (44-) 60-80\mu$.

United States: Mississippi; Italy.

209a. Var. **seriosporum** (Lagerheim) Hirn.

(Pl. XXI, figs. 191 and 192.)

1900, p. 294, Pl. XXII, fig. 114; *Oe. seriusporum* Lagerh. 1888, p. 590.

More variable, oogonium sometimes 2-5, lower series of oogonium subcylindric or cylindric-globose; vegetative cell 24-42 (-54) $\times$ 38-120 μ ; oogonium, single or upper series, 48-60 $\times$ 60-99 μ , lower series 42-54 $\times$ (35-) 43-52 μ ; oospore 40-56 $\times$ 40-75 μ .

Germany.

Characterized by the variation in the size of the oogonia and the vegetative cells on the same plant, the species is somewhat near *Oe. mexicanum* (No. 44). The variety is even more variable than the species, and its oogonia are more seriate.

210. **Oe. virceburgense** Hirn.

(Pl. XXXVI, fig. 336.)

1896, p. 8; 1900, p. 301, Pl. XXIV, fig. 128; 1906, p. 54.

Oogonium 1-8, subpyriform- to subdepressed-globose, operculate, division a little above median; oospore subdepressed- or depressed-globose, quite filling oogonium, spore wall smooth; vegetative cell often broadly capitellate; basal cell half round; vegetative cell 4-6 $\times$ 13-45 μ ; oogonium 15-20 $\times$ 15-23 μ ; oospore 14-19 $\times$ 14-17 μ .

Austria, Germany, Hungary.

To be compared with the slightly larger *Oe. spurium* (below) and with *Oe. petri* (No. 90).

211. **Oe. spurium** Hirn.

(Pl. XXXVI, fig. 333.)

1900, p. 301, Pl. XXIV, fig. 131.

Oogonium single, subdepressed- or depressed-globose, operculate, division supramedian; oospore of the same form as oogonium, nearly or completely filling it, spore wall smooth; vegetative cell capitellate; basal cell elongate; terminal cell apically obtuse or truncate-obtuse; vegetative cell 9-13 $\times$ 20-55 μ ; oogonium 26-29 $\times$ 23-28 μ ; oospore 24-28 $\times$ 21-24 μ .

United States: Ohio; Brazil.

Cf. *Oe. virceburgense* above. It also has considerable resemblance to the longer celled filaments of *Oe. sphaerandrium* (No. 105).

212. **Oe. velatum** Hallas.

(Pl. XLI, figs. 398 and 399.)

1905, p. 405, fig. 11; Hirn 1906, p. 23, Pl. IV, fig. 27.

Oogonium single, ellipsoid-globose, operculate, division superior; oospore globose, not filling oogonium, middle layer of spore wall echinate; vegetative cell 3-20 $\times$ 25-400 μ ; oogonium 43 $\times$ 68 μ ; oospore 35 $\times$ 35 μ .

Denmark.

The cell diameter is extremely variable. The echinate appearance of the oospore is questioned by Hirn (1906), who thinks the oospore may be parasitized.

213. *Oe. pilosporum* West.

(Pl. XL, fig. 395.)

1891, p. 109, Pl. 18, fig. 3; Hirn 1900, p. 318, Pl. L, fig. 325.

Oogonium single, oblong-ellipsoid, somewhat narrowed at the upper extremity; oospore subglobose, spore wall thick and outer layer densely pilose, the hairs short; vegetative cell $11-12 \times 55-70\mu$; oogonium $23 \times 48\mu$; oospore (without hairs) $19 \times 21\mu$; oospore wall 2μ thick.

Spain.

The above dimensions and description are copied gingerly by Hirn (1900). From West's figure one is inclined to agree with Hirn that the "oospore" may be nothing more than a fungal parasite within the oogonium.

214. *Oe. hoehnei* Borge.

(Pl. XXVIII, fig. 251.)

1925, p. 12, Pl. I, fig. 11.

(?) Dioecious, macrandrous; oogonium single, ellipsoid, pore superior; oospore globose, not filling oogonium in length, outer spore wall smooth, middle wall areolate; vegetative cell $43-44 \times 148-240\mu$; oogonium $66-72 \times 99-108\mu$; oospore $63-65 \times 63-65\mu$.

Paraguay.

This species has spore markings similar to those of *Oe. areolatum* (No. 59).

215. *Oe. giganteum* Kuetzing; Wittrock.

(Pl. XXIX, fig. 260.)

Kuetz. 1845, p. 200; 1853, p. 12, Pl. 37, fig. 2; Wittr. 1874, p. 42; Hirn 1900, p. 295, Pl. XXIII, fig. 115.

(?) Dioecious, macrandrous; oogonium 1, cylindric-obovoid, pore superior; oospore cylindric-ellipsoid or subellipsoid, nearly filling oogonium, spore wall in three layers: outer smooth, middle layer with 25-30 longitudinal rows of pits, inner layer smooth; vegetative cell $30-50 \times 60-225\mu$; suffultory cell $40-60 \times 60-210\mu$; oogonium $53-69 \times 67-106\mu$; oospore $51-65 \times 65-103\mu$.

United States: Ohio; Austria, England, Denmark, Germany, Sweden.

This species is rather readily identified by its large size and by the longitudinal rows of distinct pits in the middle layer of the oospore wall. It is separable from *Oe. capense* (No. 216) by its larger size. The arrangement of pits and general appearance are somewhat similar to the nannandrous *Oe. concatenatum*

(No. 151). It is doubtless a distinct species, no matter what its final description may be.

216. **Oe. capense** Nordstedt and Hirn.

(Pl. XXIX, fig. 257.)

In Hirn 1900, p. 293, Pl. XV, fig. 88.

Oogonium 1-3, obovoid to ellipsoid, pore superior; oospore of same form as oogonium and filling it, spore wall in three layers: outer smooth, middle pitted, with pits in 25-30 longitudinal rows, inner layer smooth; basal cell elongate; vegetative cell $13-17 \times 40-100\mu$; oogonium $35-38 \times 53-58\mu$; oospore $33-35 \times 46-48\mu$.

Africa.

This species is perhaps dioecious, macrandrous, although antheridia are unreported. Cf. *Oe. giganteum* above.

217. **Oe. moniliforme** Wittrock.

(Pl. XII, fig. 114.)

1874, p. 40; Hirn 1900, p. 288, Pl. V, fig. 28.

Oogonium 1-5, pyriform to globose-obovoid or subglose, pore supramedian; oospore globose or subdepressed-globose, not filling oogonium, middle layer of spore wall scrobiculate; terminal cell apically obtuse; vegetative cell $9-11 \times 30-54\mu$; oogonium $23-28 \times 28-35\mu$; oospore $22-27 \times 22-26\mu$.

United States: Michigan, Ohio; Sweden.

The small size of this species together with its definitely scrobiculate spore wall makes it readily identifiable, even though antheridia have not been so far observed.

218. **Oe. inerme** Hirn.

(Pl. XI, fig. 110.)

1900, p. 287, Pl. II, fig. 10.

(?) Dioecious; macrandrous; oogonium single, subdepressed- or subpyriform-globose, pore median, rimiform and narrow; oospore depressed- or subdepressed-globose, not filling oogonium, spore wall smooth; vegetative cell $12-14 \times 56-96\mu$; oogonium $37-39 \times 34-45\mu$; oospore $33-38 \times 28-32\mu$.

England, France, Germany.

218a. Var. **mentiens** Hirn.

(Pl. XI, fig. 111.)

1900, p. 287, Pl. II, fig. 11.

Smaller; oogonium depressed- to pyriform-globose; oospore sometimes nearly globose; vegetative cell $9-11 \times 37-90\mu$; oogonium $28-32 \times 33-45\mu$; oospore $27-29 \times 25-28\mu$.

France.

The species is to be compared with *Oe. sociale* (No. 6), *Oe. rufescens* (No. 4), and *Oe. laeve* (No. 1). Var. *mentiens* is smaller.

219. *Oe. sol* Hirn.

(Pl. XXXIII, fig. 286.)

1900, p. 303, Pl. XXVIII, fig. 164.

Oogonium single, globose, operculate, division superior; oospore globose, completely filling oogonium, spore wall of three layers: outer smooth, middle with 35-45 granulate ribs sometimes anastomose, inner smooth; vegetative cell $13-15 \times 65-110\mu$; oogonium $46-55 \times 46-55\mu$; oospore $44-53 \times 44-53\mu$.

Brazil.

If found to be dioecious, macrandrous, it is near *Oe. tumidulum* (No. 73) and should perhaps be considered a variety of it.

220. *Oe. urceolatum* Nordstedt and Hirn.

(Pl. XXVII, figs. 238 and 239.)

In Hirn 1900, p. 293, Pl. XIV, fig. 81.

Oogonium 1, obpyriform (rarely oblong-ellipsoid or subellipsoid), pore superior; oospore globose-ellipsoid or ellipsoid, not filling oogonium lengthwise, inflating lower part of oogonium, spore wall in three layers: outer smooth, middle with 15-20 entire, longitudinal ribs, sometimes anastomose and often spirally curved, inner layer smooth; basal cell elongate; vegetative cell $24-30 \times 120-210\mu$; oogonium $58-70 \times 100-125\mu$; oospore $54-60 \times 58-70\mu$.

Brazil.

To be compared with the monoecious *Oe. paulense* (No. 52) and with the dioecious *Oe. boscii* (No. 55).

221. *Oe. londinense* Wittrock.

1874, p. 39; Cooke 1884, p. 170, Pl. 65, fig. 4; Hirn 1900, p. 317.

Oogonium 1-2, globose, operculate, division median; oospore globose, quite filling oogonium; (?) antheridium 1-2, hypogynous; vegetative cell $10-15 \times 15-75\mu$; oogonium $33-35 \times 33-43\mu$; oospore $27-32 \times 27-32\mu$; (?) antheridium $12 \times 10-11\mu$.

England; (?) New Jersey.

This species is very uncertain and Cooke's figure helps none. It has very much the appearance of *Oe. areschougii* (No. 160) except for Wittrock's suspicion that it is monoecious. Quite likely Wittrock was wrong, and the "antheridia" are in reality androsporangia.

222. *Oe. consociatum* Collins and Hervey.

(Pl. XL, fig. 396.)

1917, p. 37, Pl. I, figs. 1-4; Collins 1918, p. 65; P. B. A. No. 2068.

Dioecious, ? macrandrous; oogonium single, globose to depressed-globose, operculate, division median to superior, narrow but distinct; oospore globose to depressed-globose, filling oogonium, spore wall smooth; basal cell depressed-globose; vegetative cell, more or less

distinctly clavate, $6-12 (-20) \times 6-50\mu$; oogonium $28-26 \times 28\mu$; oospore $26 \times 24-26\mu$; basal cell $20-24 \times 12-16\mu$.

Bermuda.

Characterized by its frequently short and clavate vegetative cells and depressed-globose basal cells. Its general appearance is near that of *Oe. vesicatum* (No. 205), *Oe. montagnei* (No. 206), and *Oe. pluviale* (No. 183). Its resemblance to *Oe. inversum* (No. 77), noted by Collins and Hervey, seems to the writer quite far-fetched. The so-called "stellate clusters" of young plants may be merely the development of numerous zoospores accidentally lodged in one mass.

223. *Oe. lageniforme* Hirn.

(Pl. XX, fig. 187.)

1900, p. 291, Pl. XIII, fig. 68.

(?) Dioecious, macrandrous; oogonium 1, obpyriform (in section sometimes trinodulose), pore superior; oospore globose to ellipsoid, occupying the lower part of the oogonium, spore wall smooth; vegetative cell $11-13 \times 45-90\mu$; oogonium $33-36 \times 48-63\mu$; oospore $29-31 \times 31-38\mu$.

Brazil.

In spite of the incomplete description this species is readily recognized by its inferiorly inflated oogonium and its relatively smaller oospore. It bears some resemblance to the monoecious *Oe. pseudoboscii* (No. 27).

224. *Oe. poecilosporum* Nordstedt and Hirn.

(Pl. XXXIV, fig. 305.)

In Hirn 1900, p. 298, Pl. XXIII, fig. 124.

Oogonium 1-2, ellipsoid to depressed-globose (sometimes globose or subglobose), operculate, division median, broad; oospore of same form as oogonium, which it nearly fills, spore wall smooth; terminal cell apically obtuse; vegetative cell $6-8 \times 20-47\mu$; oogonium $24-28 \times 25-38\mu$; oospore $23-26 \times 21-28\mu$.

Africa.

Compare with *Oe. gunnii* (No. 101) and *Oe. psaegetosporum* (No. 83).

225. *Oe. inflatum* Hallas.

(Pl. XXI, figs. 193 and 194.)

1905, p. 408, fig. 16; Hirn 1906, p. 16, Pl. I, fig. 1.

(?) Dioecious; oogonium 1-2, often terminal, subglobose or ellipsoid-globose, poriferous, pore variable between suprmedian and nearly

superior; oospore globose, not filling oogonium, spore wall smooth; basal cell elongate; vegetative cell of considerable variation in diameter, the terminal cell often narrow and produced into a long hyaline seta; vegetative cell $6-27 \times 12-480\mu$; oogonium $37-49 \times 54-76\mu$; oospore $28-35 \times 28-35\mu$.

Denmark.

Cf. *Oe. tyrolicum* (No. 14) and *Oe. urbicum* (No. 11). The chief characteristics of *Oe. inflatum* are the variable position of the pore, the considerable variation in size of the vegetative cell, and the long, narrow hyaline setæ.

226. *Oe. warmingianum* Wittrock.

(Pl. XV, fig. 148.)

1878, p. 140; Hirn 1900, p. 290, Pl. XII, fig. 65.

Oogonium single, ellipsoid-ovoid, pore superior; oospore ellipsoid-globose, not filling oogonium in length, spore wall smooth; (? antheridium subepigynous); vegetative cell $8-9 \times 32-63\mu$; oogonium $35 \times 53\mu$; oospore $30 \times 33\mu$.

Brazil.

It is impossible to determine with any degree of certainty the position of this species in the genus because of the very inadequate description.

227. *Oe. calvum* Wittrock.

(Pl. XXXIV, figs. 309 and 310.)

1874, p. 37; *Oe. vesicatum* Link in Rabenhorst Alg. Europe 1873; Hirn 1900, p. 316, Pl. L, fig. 323.

Oogonium 1-5, subdepressed-globose, operculate, division median; vegetative cell $7-9 \times 22-40\mu$; oogonium $27-30 \times 25-30\mu$.

India.

Cf. *Oe. poecilosporum* (No. 224), *Oe. latiusculum* (No. 81), and *Oe. pratense* (No. 80).

228. *Oe. inclusum* Hirn.

(Pl. XL, fig. 397.)

1895, p. 21, Pl. 1, fig. 4; 1900, p. 318, Pl. L, fig. 324; Fremy and Meslin 1926, p. 124, fig. 8.

Oogonium 1, suboblong-ellipsoid, wall thickened; oospore ellipsoid or suboblong-ellipsoid, not completing oogonium or sometimes enlarging it, spore wall slightly rugulose; basal cell elongate; vegetative cell $8-12 \times 33-150\mu$; oogonium $24-30 \times 48-55\mu$; oospore $18-23 \times 38-48\mu$.

Finland, France.

229. *Oe. plicatum* Wittrock.

(Pl. LVI, fig. 535.)

1874, p. 38; Hirn 1900, p. 310, Pl. XLVII, fig. 296; *Oe. rothii* Breb. in Rabenh. Alg. Eur. 2347, 1873.

Oogonium 1-3, subpyriform-globose, medianly plicate, with 12-17 longitudinal ridges, operculate, division supramedian; oospore subdepressed- to subpyriform-globose, nearly filling oogonium, spore wall smooth; terminal cell apically obtuse; vegetative cell $7-11 \times 16-65\mu$; oogonium $23-28 \times 24-29\mu$; oospore $21-25 \times 21-25\mu$.

India.

To be compared with *Oe. oelandicum* (No. 164), *Oe. megaporum* (No. 165), and *Oe. crenulatum* (No. 233).

230. *Oe. lagerstedtii* Wittrock.

(Pl. LVI, figs. 536-538.)

1874, p. 38; Hirn 1900, p. 311, Pl. XLVII, fig. 306.

Oogonium 1-4, depressed-obovoid, medianly plicate, with 7-11 longitudinal ridges, operculate, division inframedian; oospore depressed-globose or subglobose, not filling oogonium, spore wall smooth; vegetative cell $5-8 \times 10-48\mu$; oogonium $16-23 \times 13-17\mu$; oospore $13-16 \times 11-14\mu$.

Finland, Sweden.

Cf. *Oe. platygynum* (No. 167).231. *Oe. uleanum* Hirn.

(Pl. LV, fig. 529.)

1900, p. 311, Pl. XLVIII, fig. 308; *Oe. sp.* Moebius 1895, p. 174, Pl. 2, figs. 11 and 12.

Oogonium 1-2, subpyriform, medianly plicate, with 7-9 longitudinal ridges, operculate, division inframedian; oospore globose or subglobose, not filling oogonium, spore wall smooth; vegetative cell $8-10 \times 40-80\mu$; oogonium $23-29 \times 24-39\mu$; oospore $18-22 \times 18-25\mu$.

Brazil.

Cf. *Oe. platygynum* (No. 167) and *Oe. lagerstedtii* above.232. *Oe. pulchrum* Nordstedt and Hirn.

(Pl. LVI, fig. 541.)

In Hirn 1900, p. 312, Pl. XLVIII, fig. 309.

(?) Dioecious-nannandrous; oogonium 1, depressed-pyriform, medianly plicate, with 7-10 longitudinal ridges, operculate, division inframedian; oospore globose or subglobose, not filling oogonium, spore wall smooth; (?) androsporangium 1-4, subepigynous or subhypogynous or rarely hypogynous; vegetative cell capitellate; terminal cell (often an oogonium) apically obtuse, vegetative cell $6-9 \times 20-52\mu$; suffultory cell $9-12 \times 15-36\mu$; oogonium $23-26 \times 19-26\mu$; oospore $18-23 \times 16-22\mu$; (?) androsporangium $7-8 \times 7-8\mu$.

Brazil.

Cf. *Oe. bahusiense* (No. 166) and *Oe. platygynum* (No. 167).

233. *Oe. crenulatum* Wittrock.

(Pl. LVI, figs. 532 and 533.)

1876, p. 49, Pl. 13, figs. 25-27; Hirn 1900, p. 309, Pl. XLVII, fig. 294.

Oogonium 1, broadly pyriform-globose or transversely ellipsoid, medianly plicate, with about 13 longitudinal ridges, operculate, division median, narrow but distinct; oospore depressed-globose, inflating oogonium, spore wall smooth; vegetative cell $6-8 \times 30-55\mu$; oogonium $25-27 \times 26-28\mu$; oospore $22-23 \times 15-17\mu$.

Italy.

233a. Var. *gracilius* (Nordstedt) Hirn.

(Pl. LVI, fig. 534.)

1900, p. 309, Pl. XLVII, fig. 295; *Oe. crenulatum* Wittr. f. *gracilior* Nordst. 1888, p. 12.

Smaller; vegetative cell $4-5 \times 13-40\mu$; oogonium $16-18 \times 16-21\mu$; oospore $15-16 \times 12-13\mu$.

New Zealand.

Cf. *Oe. plicatulum* (No. 229), from which it is distinguished by its median position of the operculum.

234. *Oe. saxatile* Hansgiring.

1901, p. 1; Hirn 1906, p. 20.

In 1888 Hansgiring described *Oe. rufescens* var. *saxatile* as having an orange-red color and with these dimensions: vegetative cell $7-9(-12) \times 15-70\mu$; oogonium $18-24 \times 24-36\mu$. Hirn (1900) considered this as a synonym of *Oe. rufescens* (No. 4). In 1901 Hansgiring, apparently feeling the variety to be distinct, gave it specific rank as *Oe. saxatile*, adding nothing to its description. It was found in several localities in Austria in a vegetative state on moist rock. *Oe. saxatile* may be distinct, but its description is at present too incomplete to warrant its retention as a distinct species, except as a matter of record. Vegetative cells red in color have been noted in a very few species of *Bulbochaete*, e. g., *B. sanguinea* (No. 50).

235. *Oe. reticulatum* West and West.

1902, p. 129; Hirn 1906, p. 19.

Vegetative cell elongate; oogonium ovoid-globose; oospore globose, not filling oogonium, spore wall areolate; vegetative cell $7-8 \times 62-100\mu$; oogonium $37 \times 41\mu$; oospore $25 \times 25\mu$.

Ceylon.

To be compared with *Oe. arcyosporum* (No. 58) and *Oe. areolatum* (No. 59).

236. *Oe. reinschii* Roy.

(Pl. XLI, fig. 406.)

In Cooke 1884, p. 160, Pl. 57, fig. 23; *Cymatonema* sp. Reinsch 1875, p. xi, Pl. 6, fig. 1 (incorrectly *Cymatopleura*, p. 77); *Oe. sterile* Hansg. 1888, p. 43 and 260; Hirn 1900, p. 319, Pl. L, fig. 326; Lewis and Taylor 1928, p. 193, text figs. 3 and 4.

Vegetative cell subhexagonal or subellipsoid, sometimes nearly cylindrical; basal cell subhemispherical; apical cell obtuse; vegetative cell 6-9 (-11) $\times$ 8-24 μ ; basal cell 8-9 $\times$ 5-6 μ .

United States: Michigan, Florida, New York, Massachusetts; Brazil, Paraguay; Austria, England, France, Germany, Scotland, Sweden.

This species, though as yet never found in fruit, is readily recognized by its peculiar vegetative cells. Among the typically hexagonal and ellipsoid cells nearly always can be found regularly cylindrical cells.

TABLE III.

Alphabetical list of synonyms. The numbers indicate the species of *BULBOCHAETE* in the text considered synonymous, or under which they are discussed. If the number is preceded by a question mark, the synonym is considered very doubtful and cannot be definitely authenticated.

BULBOCHAETE	No.	BULBOCHAETE	No.
<i>anomala</i> Pringsh.....	?46	<i>pygmaea</i> Wittr.....	34
<i>brebissonii</i> Kuetz. f.....	? 9a	<i>pygmaea</i> a. <i>major</i> Pringsh.....	38a
<i>canbyii</i> Wood.....	?23	<i>pygmaea</i> b. <i>minor</i> Pringsh.....	34
<i>crenulata</i> Pringsh. v. <i>plena</i> Wittr.....	11	<i>rectangularis</i> Wittr. v. <i>lund-</i> <i>dellii</i> Wittr.....	36
<i>crenulata</i> Pringsh. v. <i>supra-</i> <i>mediana</i> Wittr.....	12	<i>rectangularis</i> Wittr. v. <i>tenuis</i> Wittr.....	35
<i>dumosa</i> Wood.....	38a	<i>rectangularis</i> Wittr. v. <i>nor-</i> <i>vegica</i> Wittr.....	35
<i>elachistandria</i> Wittr.....	?12	<i>reticulata</i> Nordst.....	46a
<i>elachistandria</i> Wittr.....	?14	<i>reticulata</i> Nordst. v. <i>minor</i> Lemm.....	38a
<i>ellipsozona</i> West.....	?27	<i>rhadinospora</i> Wittr.....	?36
<i>gracilis</i> Pringsh.....	6b	<i>setigera</i> Agardh.....	23
<i>ignota</i> Wood.....	?37	<i>setigera</i> (Roth) Agardh.....	27
<i>intermedia</i> Pringsh.....	20	<i>setigera</i> v. <i>canbyii</i> Wood.....	?23
<i>intermedia</i> DeBary f. <i>ameri-</i> <i>cana</i> Hirn.....	12	<i>setigera</i> (Roth) Agardh v. <i>punctulata</i> Nordst.....	26
<i>intermedia</i> DeBary f. <i>supra-</i> <i>mediana</i> (Wittr.) Hirn.....	12	<i>speciosa</i> Wittr.....	46
<i>intermedia</i> DeBary v. <i>depressa</i> Hirn.....	12a	<i>subsimplex</i> Wittr.....	38a
<i>minor</i> Al. Br.....	41a	<i>tenuis</i> (Wittr.) Hirn v. <i>nor-</i> <i>vegica</i> (Wittr.) Hirn.....	35
<i>minor</i> Al. Br.....	44	<i>tumida</i> Wittr.....	9
<i>mirabilis</i> Wittr. v. <i>lapponica</i> Wittr. and Lund.....	6	<i>varians</i> Wittr. v. <i>antiqua</i> Nordst.....	40a
<i>mirabilis</i> Wittr. v. <i>immersa</i> Wittr.....	6a	<i>varians</i> Wittr. v. <i>alpina</i> Wittr. and Lund.....	38a
<i>nana</i> Wittr.....	? 2		
<i>nana</i> Wittr. v. <i>subbasispora</i> Wittr.....	1		
<i>pachyderma</i> Reinsch.....	?46		
<i>pringsheimiana</i> Archer.....	?46		
		CONFERVA	
		<i>setigera</i> Roth.....	23

TABLE IV.

Alphabetical list of synonyms for the genus OEDOGONIUM. The numbers indicate the species or varieties in the text considered synonymous, or under which they are discussed. If the number is preceded by a question mark, the synonym is considered very doubtful and cannot be definitely authenticated.

OEDOGONIUM	No.	OEDOGONIUM	No.
<i>acrosporum</i> f. <i>connectens</i> Wittr.	195	<i>crispum</i> v. <i>uruguayense</i> f. <i>pro-</i>	
<i>acrosporum</i> v. <i>boreale</i> Wolle...	195a	<i>prium</i> Hirn.	109e
<i>africanum</i> Lagerh.	200	<i>crispum</i> f. <i>vernale</i> (Hass.	
<i>alternans</i> Kirch.	113	Wittr.) Hirn.	109
<i>amplum</i> Magn. and Wille.	49a	<i>cryptosporum</i> v. <i>vulgare</i> f. <i>abbrev-</i>	
<i>angulosum</i> Hallas.	135a	<i>iata</i> Gutw.	169
<i>apiculatum</i> Wolle.	? 57	<i>cryptosporum</i> v. <i>subdepressum</i>	
<i>apophysatum</i> Al. Br.	138	Wittr.	169
<i>apophysatum</i> Al. Br.	150	<i>cyathigerum</i> v. <i>ornatum</i> Wittr.	149b
<i>apophysatum</i> Al. Br.	151	<i>cyathigerum</i> v. <i>rumelica</i> Istan.	149
<i>apophysatum</i> Pringsh.	151	<i>debaryanum</i> Chmiel.	37a
<i>archerianum</i> Cooke.	? 53b	<i>delicatulum</i> Kuetz.	? 168a
<i>archavaletæ</i> Wittr.	142a	<i>diandronites</i> Carter.	? 25
<i>areschougii</i> f. <i>major</i> Collins.	160b	<i>dioicum</i> Petrov.	37
<i>bathmidosporum</i> Nordst.	195b	<i>echinatum</i> Wittr.	147
<i>bernardense</i> Bates.	171c	<i>eremitum</i> Hallas.	184d
<i>berolinense</i> Wittr.	154	<i>euganeorum</i> Wittr.	95b
<i>birmanicum</i> Wittr.	138	<i>exiguum</i> Elfv.	4a
<i>birmanicum</i> Wittr.	? 188	<i>fionia</i> Hallas.	181c
<i>borisianum</i> Wittr.	138	<i>flavescens</i> Wittr.	133
<i>calcareum</i> v. <i>gaditanum</i> Lewin.	132	<i>flavescens</i> v. <i>gynandrosporum</i>	
<i>calosporum</i> Hirn.	57c	Hirn.	133
<i>candollei</i> (Le Cl.) Breb.	24	<i>fonticola</i> v. <i>flavescens</i> Hansg.	? 183
<i>capillare</i> Kuetz.	35	<i>gemelliparum</i> Pringsh.	45a
<i>capillare</i> v. <i>flavescens</i> .	35	<i>gemelliparum</i> v. <i>majus</i> Wittr.	48
<i>capilliforme</i> v. <i>australe</i> f.		<i>grande</i> v. <i>aequatoriale</i> f. <i>hor-</i>	
<i>diversum</i> Hirn.	37d	<i>tense</i> Wittr.	45b
<i>capilliforme</i> v. <i>australe</i> f.		<i>grande</i> f. <i>robusta</i> Hirn.	45d
<i>uberosporum</i> Hirn.	37c	<i>grande</i> v. <i>robustum</i> (Hirn)	
<i>carbonicum</i> Wittr.	15c	Tiffany.	45d
<i>cardiacum</i> Wittr.	15	<i>hafniense</i> Hallas.	132a
<i>calaractum</i> Wolle.	142c	<i>hexagonum</i> Kuetz.	135
<i>ciliare</i> De Not.	95b	<i>hispanicum</i> Lewin.	109
<i>ciliatum</i> Pringsh.	185	<i>hohenackerii</i> Wittr.	184e
<i>cleveanum</i> v. <i>arvensis</i> Istv.	154	<i>hormosporum</i> West.	149e
<i>cleveanum</i> f. <i>exoticum</i> Hirn.	154	<i>huillense</i> West.	118
<i>concatenatum</i> Wittr.	151	<i>hutchinsiae</i> Wittr.	151a
<i>concatenatum</i> f. <i>luxurians</i> Breb.	150	<i>insignis</i> v. <i>minus</i> Hirn.	74a
<i>concatenatum</i> v. <i>lagenarioides</i>		<i>intermedium</i> Kuetz.	183
Filar.	151	<i>inversum</i> v. <i>subclusum</i> Wittr.	77
<i>condensatum</i> Hallas.	144a	<i>inversum</i> f. <i>subclusum</i> (Wittr.)	
<i>crassipellitum</i> West.	142b	Hirn.	77
<i>crassum</i> Wittr.	49	<i>kjellmanii</i> Wittr.	54
<i>crenulatum</i> f. <i>gracilior</i> Nordst.	233a	<i>klebahnii</i> Lemm.	200
<i>crispulum</i> Wittr. and Nordst.	178	<i>lagerheimii</i> Wittr.	109c
<i>crispulum</i> v. <i>minutum</i> Hansg.	179a	<i>landsboroughi</i> Wittr.	48
<i>crispum</i> v. <i>elongatum</i> Wittr.	109	<i>landsboroughi</i> v. <i>gemelliparum</i>	
<i>crispum</i> f. <i>gracilis</i> Wittr.	109e	(Pringsh.) Wittr.	45a
<i>crispum</i> v. <i>granulosum</i> Nordst.	109a	<i>landsboroughi</i> v. <i>robustum</i> Wittr.	48a

TABLE IV—(Continued).

OEDOGONIUM	No.		OEDOGONIUM	No.
<i>londinense v. compressum West.</i>	171c		<i>rufescens subsp. lundellii f. elongatum Hirn.....</i>	4b
<i>longicolle v. senegalense f. afghanicum Schaarsch.....</i>	166a		<i>seriosporum Lagerh.....</i>	209a
<i>lorentzii Magn. and Wille.....</i>	37b		<i>setigerum Vaup.....</i>	138
<i>lundellii Wittr.....</i>	4b		<i>spletsbergense Wittr..... ?</i>	3a
<i>lundense Wittr.....</i>	184c		<i>sphaericum Hallas.....</i>	116b
<i>lundense Wittr.....</i>	?184b		<i>spinosperrnum Reinsch.....</i>	?136
<i>macrandrium v. aemulans Hirn.....</i>	184b		<i>spiromanulatnm Schm.....</i>	84
<i>mactroapremum f. palagonieum Borge and Hirn.....</i>	172		<i>stagnale Kuetz.....</i>	35
<i>naximum West.....</i>	41a		<i>stagnale Kuetz.....</i>	35a
<i>monandronites Carter.....</i>	? 25		<i>stagnale v. variabilis Lewin.....</i>	40
<i>montagnei v. submarine Wittr.....</i>	183		<i>sterile Hansg.....</i>	236
<i>montagnei v. saxicolnm Wittr....</i>	183		<i>subpisanum Lewin.....</i>	99
<i>monticchii Fior-Mazz.....</i>	77		<i>subcapitellatnm Hirn.....</i>	105
<i>neglectum Hirn.....</i>	27		<i>tapeinosporum v. angolense West and West.....</i>	196
<i>nordstedtii Wittr.....</i>	95b		<i>trichosporum Itzigsh..... ?</i>	9
<i>oblongum v. majus Nordst.....</i>	116a		<i>tumidulum Pringsh.....</i>	11
<i>obruncatum v. oblatum Tilden..</i>	?160b		<i>tumidulum (Roth).....</i>	29
<i>ochroleucum Kuetz.....</i>	6		<i>tumidulum Kuetz.....</i>	48
<i>oelandicum v. subpyriforme Wittr.....</i>	165		<i>tmdulatnm Al. Br.....</i>	119
<i>ornatum Hirn.....</i>	149		<i>tmdulatnmv. interrupte-incisum Schrod.....</i>	119
<i>pachydermatosporum Nordst....</i>	95		<i>tmdulatnm v. moebuisii Schmid..</i>	119
<i>pachydermatosporum Nordst....</i>	95b		<i>tmdulatnm v. senegalense Nordst.....</i>	119b
<i>paludosum Wittr.....</i>	53		<i>urbicum v. oviforme Lewin.....</i>	? 25
<i>paludosum (Hass.) Kuetz.....</i>	26		<i>vacherii Braun.....</i>	30
<i>piliferum Wittr.....</i>	86		<i>vacherii Al. Br.; Wittr. f. insulare Hirn.....</i>	25
<i>piliferum Auersw.....</i>	185		<i>vernale Wittr.....</i>	109
<i>platygynm f. major West.....</i>	167		<i>vernale Wittr.....</i>	12
<i>polymorphum Wolle.....</i>	10		<i>vesicatnm Link.....</i>	171
<i>princeps Wittr.....</i>	38		<i>vesicatnm Link.....</i>	227
<i>pringheimiana Archer.....</i>	? 53b		<i>vesicatnm Wittr.....</i>	205
<i>pringheimi v. nordstedtii f. pachydermatosporum (Nordst.) Hirn.....</i>	95		<i>vesicatnm v. flavesceus Kuetz... </i>	109b
<i>pringheimii v. nordstedtii f. euganeorum (Wittr.) Hirn....</i>	95b		<i>vesicatnm v. fuscesceus Dupray..</i>	135
<i>pringheimii v. varians Nordst..</i>	95b		<i>wittrockianum Hirn.....</i>	145a
<i>propinquum Wittr.....</i>	184d		<i>wollcanum v. insigne Nordst....</i>	150b
<i>punctato-striatum v. minor Wittr.....</i>	84			
<i>pyrrulum v. obesum Wittr.....</i>	108		ANDROGYNIA	No.
<i>rhodosporum (Welw.) Wittr.....</i>	208		<i>echinata Wood.....</i>	147
<i>rivulare Al. Br.....</i>	42		<i>hunatii Wood.....</i>	152
<i>rvolare v. major Wolle.....</i>	49a		<i>mirabilis Wood.....</i>	?138
<i>rostcllatum Pringsh.....</i>	109		<i>multispora Wood.....</i>	143
<i>rostcllatum Pringsh.....</i>	109b			
<i>rotkui Breb.....</i>	4b		CONFERRA	No.
<i>rotkui Pringsh.....</i>	169		<i>capillare L.....</i>	35
<i>rotkui f. major West.....</i>	171		<i>tumidula Eng. Bot.....</i>	73
<i>rufescens f. exiguum (Elv.) Hirn..</i>	4a		<i>tmdulata Breb.....</i>	119
<i>rufescens subsp. lundellii (Wittr.) Hirn.....</i>	4b		<i>vesicata Ag.....</i>	55
			<i>vesicata Lyng.....</i>	205

TABLE IV—(Continued).

CYMATONEMA	No.	VESICULIFERA	No.
<i>confervaceum</i> Kuetz.....	119	<i>aurea</i> Hass.....	138
		<i>borissii</i> Hass.....	138
		<i>candollei</i> Hass.....	?109
		<i>capillaris</i> Hass.....	38
PRINGSHEIMIA	No.	<i>cardiaca</i> Hass.....	15
<i>inaequalis</i> Wood.....	? 15	<i>ciliata</i> Hass.....	185
		<i>compressa</i> Hass.....	? 5
		<i>concatenata</i> Hass.....	151
		<i>crispa</i> Hass.....	109
		<i>crassa</i> Hass.....	? 49
PROLIFERA	No.	<i>dissiliens</i> Hass.....	?183
<i>borisiana</i> Le Cl.....	138	<i>flavescens</i> Hass.....	?133
<i>boscii</i> Le Cl.....	? 55	<i>landsboroughi</i> Hass.....	48
<i>rivularis</i> Le Cl.....	? 42	<i>paludosa</i> Hass.....	? 53
<i>rothii</i> Le Cl.....	?169	<i>princeps</i> Hass.....	38
<i>vaucherii</i> Le Cl.....	? 25	<i>pulchella</i> Hass.....	15b
		<i>vernalis</i> Hass.....	?109

BIBLIOGRAPHY

- Ackley, Alma B. 1929. New species and varieties of Michigan algae. Trans. Amer. Micros. Soc. 48 (3): 302-309. Pls. XXXV and XXXVI, figs. 1-22.
- Agardh, C. A. 1817. Synopsis algarum Scandinaviae, adjecta dispositione universali algarum. pp. i-xi; 1-135. Lund.
- Archer, W. 1866. Description of Bulbochaete Pringsheimiana n. sp. Quar. Jour. Micros. Sci. 6:121-123.
- Bates, F. 1886. Catalogue of the algae of Leicestershire. Flora of Leicestershire, including the Cryptogams, with maps of the county. Leicester Literary and Philos. Soc. 400 pp. 2 col. maps.
- Blackman, F. F., and A. G. Tansley. 1902. A revision of the classification of the green algae. New Phytol. 1:17-24; et seq.
- Borge, O. 1896. Australische Süßwasser-Chlorophyceen. Bihang till K. Svenska Vet.-Akad. Handlingar, Stockholm. 22 (Pt. III, No. 9):1-32. Pl. I-IV, figs. 1-66.
- Borge, O. 1899. Ueber tropische und subtropische Süßwasser-Chlorophyceen. Bihang till K. Svenska Vet.-Akad. Handlingar 24 (12):33 pp. Pl. 2.
- Borge, O. 1901. Süßwasseralgen aus Süd-Patagonien. Bihang till K. Svenska Vet.-Akad. Handlingar 27 (10):1-40. Pl. 1-2.
- Borge, O. 1911. Algologische Notizen 6-7. Botaniska Notiser 1911:197-244. Pl. 2, fig. 10.
- Borge, O. 1913. Beiträge zur Algenflora von Schweden. 2. Botaniska Notiser 1913:1-110. Pl. 1-3. Text fig. 1-2. Ibid. 3. Arkiv für Botanik 18 (10): 1-33. Pl. 1-2. Text fig. 1-2.
- Borge, O. 1925. Die von Dr. F. C. Hoehne während der Expedition Roosevelt-Rondon gesammelten Süßwasseralgen. Archiv für Botanik 19 (17):(1-56). 6 Pls.
- Bory, J. B. de St. Vincent. 1822. Dict. classique d'hist. naturelle. Vol. I, 604 pp. Paris.
- Braun, A. 1849/50. Betrachtungen über die Erscheinung der Verjüngung in der Natur. i-xvi; 1-363 pp. Freiburg.
- Braun, A. 1855. Ueber Chytridium. Abh. K. Akad. Wissensch. zu Berlin 1855: 21-84. Pl. I-VI, 116 figs.
- Carter, H. J. 1858. On specific character, fecundation, and abnormal development in Oedogonium. The Annals and Mag. of Nat. Hist. 1 (Series 3):29-39. Pl. III, figs. 1-16.
- Carter, Nellie. 1926. Freshwater algae from India. Records of the Botanical Survey of India. 9 (4):263-302. Pl. I-II.
- Chmielevsky, V. 1889. Diagnoses algarum novarum. Travaux de la Societe de naturel a l'Universite de Kharkow.
- Cleve, A. 1895. En röd Bulbochaete. Botaniska Notiser 1895:247.
- Collins, F. S. 1908. Oedogonium Huntii rediscovered? Rhodora 10 (112):57-58.
- Collins, F. S. 1909. The green algae of North America. Tufts College Studies 2: 1-480. Pl. I-XVIII. figs. 1-160.
- Collins, F. S. 1912. The green algae of North America (supplementary paper). Tufts College Studies 3: 71-109. Pl. I-II. figs. 1-12.
- Collins, F. S. 1918. The green algae of North America (second supplement). Tufts College Studies 4:1-106. Pl. I-III. figs. 1-28.
- Collins, F. S. 1918a. Notes from the Woods Hole Laboratory—1917. Rhodora 20: 141-143. Pl. 124. figs. 1-5.
- Collins, F. S., and A. B. Hervey. 1917. The algae of Bermuda. Proc. Amer. Acad. 53 (1): 1-195. Pl. I-VI.

- Cooke, M. C. 1884. British fresh-water algae. Vol. I and II. 329 pp. 102 pl. London.
- Cramer, C. 1859. *Oedogonium Pringsheimii* Cramer nova spec. *Hedwigia* 1859 (3): 17-18. Pl. I, fig. C (1-4).
- De Bary, A. 1854. Ueber die Algengattung *Oedogonium* und *Bulbochaete*. Abhandl. d. Senckbergischen Gesellsch. 1:29-104. Pl. II-IV.
- De Bary, A. 1856. Ueber den geschlechtlichen Zeugungsprozess bei den Algen. Berichte über die Verhandl. der Gesellsch. für Beförderung der Naturwissensch. zu Freiburg. 1858: 215-230.
- De Notaris, G. 1868. In *Hedwigia* 1868 (7):120.
- De Toni, G. B. 1889. *Sylloge Algarum omnium hucusque cognitarum*. 1. Chlorophyceae. pp. 15-91.
- De Wildemann, E. 1896. *Materiaux pour la Flore algologique du département de la Meuse* (France). *La Notarisia* 11 (1): 65-73.
- Eichler, B. 1893. *Materyaly do flory wodorostow okolic Miedzyrzecza*. *Pamiętnik Fizyograficzny* 12 (3): 119-136, Pl. 2-4. (1894).
- Filarszky, F. 1899. *Adatok a Pieninek moszatvegetatiojához*. *Matematikai és Természettudományi Közlemények* (1901) 27 (4):721-800.
- Fiorini-Mazzanti, E. 1860. *De novis microphyceis comitissae*. *Atti dell Accademia Pontificia de nuovi Lincei*. 1860:259-260. 6 figs.
- Fremy, P., and R. Meslin. 1926. *Excursion botanique dans la lande de la Meauffe*. *Bull. de la Société Linéenne de Normandie* 9 (7):118-130.
- Fritsch, F. E. 1902. *Algological notes II. The germination of the zoospores in Oedogonium*. *Ann. Bot.* 16: 412-417, text fig. 23.
- Fritsch, F. E. 1902. *The structure and development of the young plants in Oedogonium*. *Ann. Bot.* 16: 467-485.
- Fritsch, F. E. 1904. *Algological notes. No. 5. Some points in the structure of a young Oedogonium*. *Ann. Bot.* 18: 648-653. fig. 61.
- Fritsch, F. E. 1912. *Freshwater algae of the South Orkneys*. *Report Sci. Results of the Scottish Antarctic Exped.* 3: 95-134.
- Fritsch, F. E. 1918. *A contribution to our knowledge of the freshwater algae of Africa*. *Annals of South African Museum* 9: 486-611. figs. 1-43.
- Fritsch, F. E. 1927. *A treatise on the British freshwater algae, by the late G. S. West. New and Revised Ed.* Pp. xviii-534. 207 figs. Cambridge University Press.
- Fritsch, F. E., and F. Rich. 1913. *British freshwater algae in nature*. *Annales de Biologie lacustre* 6: (1-83). Text fig. 1.
- Fritsch, F. E., and F. Rich. 1924. *Contributions to our knowledge of the freshwater algae of Africa*. *Trans. Royal Soc. of S. Africa* 11: 297-398.
- Fritsch, F. E., and E. Stephens. 1921. *Contributions to our knowledge of the freshwater algae of Africa*. *Trans. Royal Soc. of S. Africa* 9: 1-72. figs. 1-29.
- Gutwinski, R. 1896. *De nonnullis algis novis vel minus cognitis*. *Krakow, Akad. Mat.-Przyrod. Rozpr.* (1898) 13: 33-63; *Cracovie Ac. Sci. Bull.* 1896: 342-350.
- Gutwinski, R. 1897. *Materyaly do Flory Głonow Galicyi Res ad Floram Algarum Galiciae Congestae Part. IV. La Nuova Notarisia* 1897: 125-136.
- Hallas, E. 1905. *Nye Arter af Oedogonium fra Danmark*. *Botanisk Tidssk.* 26 (3): 397-410. figs. 1-18.
- Hansgirg, A. 1886. *Prodromus der Algenflora von Böhmen. Part I. Archiv der Naturw. Landesdurchforschung von Böhmen* 5 (6): (1-96). text-figs. 1-45.
- Hansgirg, A. 1888. *Ibid.* 6 (6): (97-288). text figs. 46-124.
- Hansgirg, A. 1888a. *Algae novae aquae dulcis*. *Notarisia* 3: 398-400.
- Hansgirg, A. 1891. *Beiträge zur Kenntnis der Süßwasser-Algen und Bacterien-Flora Böhmens, Steiermarks, der österreichisch-ungarischen Küstenlanden und Bosniens*. *Sitzungsber. d. K. Bohmischen Gesellsch. d. Wissensch. Math.-Naturw.* (1892).

- Hansgirg, A. 1892. Beiträge zur Kenntnis der Süßwasseralgen und Bakterien-Flora von Tirol und Böhmen. Ibid. 1892 (10): 105-156.
- Hansgirg, A. 1893. Prodrömus der Algenflora von Böhmen. Archiv der Naturw. Landesd. von Böhmen 8 (4): 209-212.
- Hansgirg, A. 1901. Ein Nachtrag zu meinen Prodrömus der Algenflora von Böhmen. Oesterreichische botan. Zeitschrift 51:96.
- Hansgirg, A. 1905. Grundzüge der Algenflora von Niederösterreich. Beihefte zum Bot. Centralbl. 18 (3): 417-522.
- Hardy, A. D. 1906. The fresh-water algae of Victoria. The Victorian Naturalist 23: 18-22, et seq.
- Hassall, A. H. 1842. Observations on a new group, genus and subgenus of fresh-water Confervae. Annals and Magazine of Nat. Hist. 10: 336-344; 385-395.
- Hassall, A. H. 1843. Description of British freshwater Confervae. Annals and Mag. of Nat. Hist. 11: 428-437.
- Hassall, A. H. 1845. A history of the British freshwater algae. Vol. I. 462 pp. Vol. II. 24 pp. 103 pl.
- Heering, W. 1914. Ulotrichales, Microsporales, Oedogoniales in Pascher, A. Die Süßwasserflora Deutschlands, Österreichs und der Schweiz. 6 (Chlorophyceae III): 250 pp. 385 pl. Gustav Fischer, Jena.
- Hirn, K. E. 1895. Verzeichnis finnlandischer Oedogoniaceen. Acta Soc. pro Fauna et Flora Fennica 11 (6): 1-24. 1 Pl.
- Hirn, K. E. 1896. Algologische Notizen. Öfversigt af Finska Vet.-Soc. Forhandlingar 38:18-25.
- Hirn, K. E. 1900. Monographie und Iconographie der Oedogoniaceen. Acta Soc. Sci. Fennicae 27:1-394. Pl. I-LXIV. figs. 1-396.
- Hirn, K. E. 1906. Studien ueber Oedogoniaceen. Acta Soc. Sci. Fennicae 34: 1-63. Pl. I-IV. figs. 1-29.
- Hodgetts, W. J. 1921. A study of some of the factors controlling the periodicity of freshwater algae in nature. New Phytol. 20: 150-164; 195-227; et seq.
- Hooker, J. D. 1860. Flora Tasmaniae. Vol. II. The Botany of the Antarctic voyage of H. M. Discovery Ships Erebus and Terror in the years 1839-43.
- Istvanfi, G. 1887. Diagnoses praeviae algarum novarum in Hungaria observatarum Notarisia 2: 234-242.
- Istvanfi, G. 1890. Rumeliai Algak. Frivaldszky Imre gyujtesebol. Termeszettarjai Fuzetek. 13: 67-77.
- Iwanoff, L. 1901. Observations on the aquatic flora of a lake district. From the Bologovsky Biol. Station of the Imper. Soc. of Naturalists of St. Petersburg. St. Petersburg. (Russian).
- Juranyi, L. 1873. Beitrag zur Morphologie der Oedogonien. Pringsheim's Jahr. fur wissenschaft. Botanik. 9: 1-35. Pl. I-III.
- Kirchner, O. 1878. Kryptogamenflora von Schlesien. Im Namen der Schlesische Gesellschaft für Vaterland. 2 (1)
- Klebahn, H. 1892. Studien über Zygoten. II. Befruchtung von Oedogonium Boscii. Pring. Jahrbücher wissenschaft. Bot. 24: 235-267. Pl. 3.
- Kraskowitz, G. 1905. Ein Beitrag zur Kenntnis der Zellteilungsvorgänge bei Oedogonium. Sitzber. Kais. Akad. Wiss. Wien. M. N. Kl. 114 (1): 1-38. 3 Pl. 11 text figs.
- Kuetzing, F. T. 1843. Phycologia generalis. i-xxxii; 1-485. 80 Col. Pl. Leipzig.
- Kuetzing, F. T. 1845. Phycologia germanica. pp. 1-240, Nordhausen.
- Kuetzing, F. T. 1849. Species algarum. Pp. 1-922. Leipzig.
- Kuetzing, F. T. 1853. Tabulae phycologicae. Vol. III. Nordhausen. 78 pp.
- Kuetzing, F. T. 1854. Tabulae phycologicae oder Abbildungen der Tange. Vol. IV. pp. 1-23.
- Lagerheim, G. 1890. Contribuciones a la flora algologica Ecuador. I-II. Anales Univ. de Quito No. 27:31.

- Lagerheim, G. 1893. Chlorophyceen aus Abessinien und Kordofan. *La Nuova Notarisia* 1893: 153-166.
- Le Clerc, L. 1817. Sur la fructification du genre prolifere de M. Vaucher. *Paris, Mus. Hist. Nat. Mem.* 3: 462-478.
- Lemmermann, E. 1893. Versuch einer Algenflora der Umgegend von Bremen. *Abhandl. des Naturwiss. Vereins zu Bremen* 12: 497-550.
- Lemmermann, E. 1895. Verzeichniss der in der Umgegend von Plön gesammelten Algen. *Forschungsh. Biol. Stät. Plön* 3: 18 pp.
- Lemmermann, E. 1898. Algologische Beiträge. IV-V. *Abhandl. des Naturwiss. Vereins zu Bremen* 14 (3): 501-524. Pl. V, figs. 3-11.
- Lemmermann, E. 1909. Algologische Beiträge. XI. *Oedogonium cardiacum* var. *minor* Lemm. nov. var. *Archiv für Hydrobiol. und Planktonk.* 4: 191-192.
- Lemmermann, E. 1910. Beiträge zur Kenntniss der Planktonalgen. *Archiv für Hydrobiol. und Planktonk.* 5: 291-338.
- Lewin, M. 1888. Ueber spanische Süßwasseralgen. *Bihang till K. Svenska Vet.-Akad. Handlingar, Stockholm.* 14 (Pt. III, No. 1): 1-24. Pl. I-III, figs. 1-56.
- Lewis, I. F., and W. R. Taylor. 1928. Notes from the Woods Hole Laboratory—1928. *Rhodora* 30: 193-198.
- Link, H. F. 1820. Epistola de Algis aquaticis in genera disponendis in Nees von Esenback, C. G. H. *Horae physicae berolinenses.* pp. 1-8, pl. 1.
- Linnaeus, C. 1753. *Species Plantarum* Vol. II. pp. 561-1200.
- Lyngbye, H. C. 1819. *Tentamen hydrophythologiae danicae.* *Hafniae* 4: i-xxxii; 1-248. Pl. 1-70.
- Moebius, M. 1894. Australische Süßwasseralgen II. *Abhandl. der Senckenbergischen naturforsch. Gesellsch.* (1895) 18: 309-350. Pl. I-II. 60 figs.
- Moebius, M. 1895. Ueber einige brasilianische algen. *Hedwigia* 34: 173-176; et seq.
- Nordstedt, O. 1877. Bohusläns Oedogonieer. Öfversigt af K. Vet.-Akad. *Forhandlingar* 34 (4): 21-33. 1 pl.
- Nordstedt, O. 1878. De algis aquae dulcis et de Characeis ex insulis Sandwicensibus a Sv. Berggren 1875 reportatis. *Minneskrift K. Fysiog. Sallsk.* i Lund. 1878: 1-24.
- Nordstedt, O. 1878a. Algologiska smasaker. *Botaniska Notiser* 1878 (6): 176-180.
- Nordstedt, O. 1880. De Algis et Characeis 1-2. *Acta Univ. Lund* 16: 1-20. 1 pl.
- Nordstedt, O. 1883. Tva nya afrikande Bulbochaete-arter. *Botaniska Notiser* 1883: 154-155.
- Nordstedt, O. 1888. Freshwater algae collected by Dr. S. Berggren in New Zealand and Australia. *Kongl. Svenska Vet.-Akad. Handlingar* 22 (8): 1-98. Pl. 1-7.
- Nordstedt, O. 1888a. Einige Characeen bestimmungen. *Hedwigia.* 1888 (7 and 8): 181-196.
- Ohashi, Hiro. 1926. *Oedogonium nebraskensis*, sp. nov. *Bot. Gaz.* 82: 207-214. figs. 1-20.
- Oltmanns, F. 1922. *Morphologie und Biologie der Algen* Vol. I. pp. 331-341. Gustav Fischer, Jena.
- Pascher, A. Über die Zwergmännchen der Oedogoniaceen. *Hedwigia* 46: 265-278.
- Petrovsky, A. 1861. Etudes Algologiques. I. Note sur une nouvelle espece d'Oedogonium. *Bull. de la Soc. Imperiale des Naturalistes de Moscou* 34: 605-611. Pl. XIII. figs. 6-11.
- Poulsen, V. A. 1877. Om Svaermosporens Spiring hos en Art af Slaegten Oedogonium. *Bot. Tidsskrift* 10: 1-15. Pl. 1.
- Pringsheim, N. 1855. Ueber die Befruchtung und Keimung der Algen. *Berichte d. Königl. Akad. d. Wissensch. zu Berlin* 1855: 133-165. 1 Pl. 34 figs.

- Pringsheim, N.** 1856. Untersuchungen ueber Befruchtung und Generationswechsel der Algen. Berichte d. Konigl. Akad. d. Wissensch. zu Berlin 1856: 225-237. 1 Pl. figs. 1-10.
- Pringsheim, N.** 1858. Beiträge zur Morphologie und Systematik der Algen. I. Morphologie der Oedogonien. Pringsheim's Jahrb. für Wissensch. Botanik 1:1-81. Pls. I-VI.
- Printz, H.** 1927. Oedogoniaceæ p. 244 in **Engler and Prantl** Die Natürlichen Pflanzenfamilien Vol. 3. 463 pp. Leipzig.
- Rabenhorst, L.** 1868. Flora Europaea Algarum aquae dulcis et submarinae. 1864-1868: Sec. III. pp. 1-461. Leipzig.
- Reinsch, P.** 1875. Contributiones ad Algologiam et Fungologiam 1: i-xii; 1-103. Leipzig.
- Rich, Florence.** 1925. Further notes on the algae of Leicestershire. Jour. of Bot. 63: 71-78. figs. 1-6.
- Roth, A. G.** 1806. Catalecta botanica. Vol. III. 350 pp. 12 Pls.
- Scharreschmidt, G.** 1884. Notes on Afghanistan algae. Jour. Linn. Soc. of Bot. 21: 241-250. Pl. 5.
- Schaffner, J. H.** 1927. Extraordinary sexual phenomena in plants. Bull. Torr. Club 54: 619-629.
- Scherffel, A.** 1901. Einige Beobachtungen über Oedogonium mit halbkugeliger Fusszelle. Ber. d. deutsch. Bot. Ges. 19: 557-563. Pl. 31.
- Schmidle, W.** 1894. Aus der Chlorophyceen-Flora der Torfstiche zu Virnheim. Flora 78: 42-66. Pl. 7.
- Schmidle, W.** 1896. Süßwasseralgen aus Australien. Flora 82: 297-313. Pl. 9.
- Schmidle, W.** 1901. Einige Algen, welche Prof. Dr. Volkens auf den Carolinen gesammelt hat. Hedwigia 40: 343-349.
- Schroeder, B.** 1897. Die Algen der Versuchsteiche des Schles. Fischereivereins zu Trachenberg Plön Biol. Stät. Forschungsber. 5: 29-66.
- Silfvenius, A. J.** 1903. Zur Kenntnis der Verbreitung finnischer Chlorophyceen und Cyanophyceen. Meddel. af Soc. Pro Fauna et Flora Fenn. 29: 10-22. Text figs. 1-3.
- Skuja, H.** 1927. Vorarbeiten zu einer Algenflora von Lettland. Acta Horti Botanici Univ. Latviensis 2 (2/3): 51-116. Pl. I-II.
- Skvortzow, B. W.** 1926. Über einige Süßwasseralgen aus der Nord-Manschurei im Jahre 1916 gesammelt. Archiv für Hydrobiol. 16: 421-436. 8 figs.
- Stahl, E.** 1891. Oedocladium protonema, eine neue Oedogoniaceen-Gattung. Pringsheim's Jahrb. für Wissensch. (1892) 23 (3): 339-348. Pls. 16, 17.
- Strasburger, Ed.** 1880. Ueber Zellbildung und Zelltheilung. Zeitschr. f. Naturwiss. 13 (2): 50.
- Taylor, W. R.** 1928. Alpine algal vegetation of the mountains of British Columbia. Proc. Acad. Nat. Sci. Philadelphia 80: 45-114.
- Tiffany, L. H.** 1921. New forms of Oedogonium. Ohio Jour. of Sci. 21: 272-275. Pl. 1. figs. A-F.
- Tiffany, L. H.** 1924. A physiological study of growth and reproduction among certain green algae. The Ohio Jour. of Sci. 24 (2): 65-100.
- Tiffany, L. H.** 1924. Some new forms of Spirogyra and Oedogonium. Ohio Jour. of Sci. 24: 180-187. Pls. I-III. 20 figs.
- Tiffany, L. H.** 1926. The filamentous algae of northwestern Iowa, with special reference to the Oedogoniaceae. Trans. Amer. Micros. Soc. 45 (2): 69-132. 16 pls. 179 figs.
- Tiffany, L. H.** 1927. New species and varieties of Chlorophyceae. Bot. Gaz. 83 (2): 202-206. Pl. IX, figs. 1-12.
- Tiffany, L. H.** 1928. The algal genus Bulbochaete. Trans. Amer. Micros. Soc. 47 (2): 121-177. Pls. XIV-XXIII. figs. 1-99.
- Tiffany, L. H.** 1929. A key to the species, varieties, and forms of the algal genus Oedogonium. Ohio Jour. of Sci. 29 (2): 62-80.

- Tiffany, L. H., and E. N. Transeau.** 1927. *Oedogonium* periodicity in the North Central States. *Trans. Amer. Micros. Soc.* 46 (3): 166-174.
- Tilden, Josephine E.** 1898. Observations on some West American thermal algæ. *Bot. Gaz.* 25 (1): 89-105. Pl. VIII-X. 43 figs.
- Transeau, E. N.** 1913. Annotated list of the algæ of Eastern Illinois. *Trans. Ill. Acad. of Sci.* 6: 69-89.
- Transeau, E. N.** 1914. New species of green algæ. *Amer. Jour. of Bot.* 1: 289-301. Pl. XXV-XXIX. 44 figs.
- Transeau, E. N.** 1916. The periodicity of freshwater algæ. *Amer. Jour. of Bot.* 3: 121-133.
- Transeau, E. N.** 1917. The algæ of Michigan. *Ohio Jour. of Bot.* 17: 217-232.
- Transeau, E. N.** 1918. A new species and a new variety of algæ from Oneida Lake, New York. *Tech. Pub. No. 9, N. Y. State Coll. of Forestry* 18: 237-238. Pl. A.
- Transeau, E. N., and Hanford Tiffany.** 1919. New Oedogoniaceæ. *Ohio Jour. of Sci.* 19: 240-243. Pl. XIV. figs. a-i.
- Tuttle, A. H.** 1910. Mitosis in *Oedogonium*. *Jour. Exp. Zool.* 9: 143-157. figs. 1-18.
- Van Wisselingh, C.** 1908. Über Karyokinese bei *Oedogonium*. *Bot. Centralbl.* 23: 157-190. Pl. 13-16.
- Vaucher, J. P.** 1803. *Histoire des conferves d'eau douce.* pp. i-xv; 1-285. Geneva.
- Vaupell, C.** 1859. Sur la reproduction et la fecondation d'une espece du genre *Oedogonium*. *Ann. Sci. Nat.* 11: 192-204.
- West, G. S.** 1899. The alga-flora of Cambridgeshire. *Jour. of Bot.* 37: 49-58; 106-116; et seq. Pl. 394-396.
- West, G. S.** 1904. *The British freshwater algæ.* 372 pp. 166 figs. Cambridge Univ. Press.
- West, G. S.** 1904a. West Indian freshwater algæ. *Jour. of Bot.* 42: 281-294. Pl. 464.
- West, G. S.** 1907. Report on the freshwater algæ, including phytoplankton, of the Third Tanganyika Expedition conducted by Dr. W. A. Cunningham, 1904-1905. *Jour. Linn. Soc. London* 38: 81-197. Pl. 2-10.
- West, G. S.** 1909. The algæ of the Yan Yean Reservoir, Victoria. *Jour. Linn. Soc.* 39 (269): 1-88. Pl. 1-6.
- West, G. S.** 1909a. The algæ of the Birket Qarum, Egypt. *Jour. of Bot.* 47: 237-244. Pl. 498.
- West, G. S.** 1912. Freshwater algæ. *Annals of S. African Museum* 9: 61-90.
- West, G. S.** 1912a. Algological notes. X. *Jour. of Bot.* 50: 321-325. Fig. 1.
- West, G. S.** 1914. A contribution to our knowledge of freshwater algæ from Columbia. *Mem. d. l. Soc. Neuchâtel d. Sc. Nat.* 5: 1013-1051. Text figs. 1-3. Pl. XXI-XXIII. figs. 1-71.
- West, G. S.** 1916. *Algæ* 1. 475 pp. figs. 1-270. Cambridge Univ. Press.
- West, G. S., and F. E. Fritsch.** 1927. A treatise on the British freshwater algæ. Rev. Ed. See **Fritsch, F. E.,** 1927.
- West, W.** 1891. Contribution to the freshwater algæ of West Ireland. *Jour. Linn. Soc.* 29: 103-216. Pl. 18-24.
- West, W.** 1893. Notes on Scotch freshwater algæ. *Jour. of Bot.* 31: 97-104. Pl. 333.
- West, W.** 1912. Freshwater algæ in a biological survey of Clare Island, Ireland. *Proc. Royal Irish Acad.* 31 (1): 1-62.
- West, W., and G. S. West.** 1897. *Welwitsch's African freshwater algæ.* *Jour. of Bot.* 35: 1-7; 33-42; et seq. Pl. 365-369.
- West, W., and G. S. West.** 1901. Freshwater Chlorophyceæ. *Flora of Koh-Chang.* *Bot. Tidskr.* 24: 157-186. Pl. 2-4.
- West, W., and G. S. West.** 1902. A contribution to the freshwater algæ of Ceylon. *Trans. Linn. Soc. of London.* 6 (3): 123-215. Pl. 17-22.

- West, W., and G. S. West. 1902a. A contribution to the freshwater algae of the north of Ireland. Trans. Royal Irish Acad. 32 (Sec. B, 1): 1-100. Pl. I-III.
- West, W., and G. S. West. 1903. Notes on freshwater algae III. Jour. of Bot. 41: 33-41. Pl. 446-448.
- Wille, N. 1880. Bidrag til Kundskaben om Norges Terskvandsalger I. Småalenes Chlorophyllophyceer. Christiania Vid.-Selsk. Forhandl. 1880 (11): 1-72. Pl. I-II.
- Wille, N. 1884. Bidrag til Sydamerikas Algflora I-III. Bihang till K. Svenska Vet.-Akad. Handlingar 8: 1-64. Pl. 1-3.
- Wille, N. 1887. Algologische Mitteilungen. Pring. Jahrbücher 18: 425-518. Pl. 16-19.
- Wittrock, V. B. 1870. Dispositio Oedogoniacearum Suecicarum. Öfversigt af Kongl. Vetenskaps.-Akad. Forhandl. 1870 (3): 119-144. Pl. 1.
- Wittrock, V. B. 1872. Oedogoniaceae novae, in Suecia lectae. Botaniska Notiser 1872 (1): 1-8. Pl. 1. figs. 1-8.
- Wittrock, V. B. 1872a. Om Gotlands och Olands sotvattensalger. Bihang till K. Svenska Vet.-Akad. Handl. 1 (1): 1-72. Pl. I-IV. 46 figs.
- Wittrock, V. B. 1874. Prodrömus Monographiae Oedogoniarum. Nova Acta Regiae Societatis Scientiarum Upsala. 9 (3): 1-64. Pl. I. 25 figs.
- Wittrock, V. B. 1876. in Nordstedt and Wittrock. Desmidiaceae et Oedogoniaceae ab O. Nordstedt in Italia et Tyrolia collectae. Öfversigt af K. Vet.-Akad. Forhandl. 1876 (6): 25-56. Pl. XII & XIII, figs. 1-33.
- Wittrock, V. B. 1878. Oedogoniaceae Americanae hucusque cognitae. Botaniska Notiser 1878 (5): 133-145.
- Wolle, F. 1877. Freshwater algae II. Bull. Torr. Bot. Club 6: 137-141.
- Wolle, F. 1880. Freshwater algae IV. Bull. Torr. Club 7: 43-48.
- Wolle, F. 1887. Freshwater algae of the United States. Vols. 1 and 2. Bethlehem, Pa.
- Wood, H. C. 1869. On Oedogonium Huntii. Proc. Amer. Phil. Soc. 1865-1868. 10 (73): 333-335.
- Wood, H. C. 1872. A contribution to the freshwater algae of North America. Smithsonian. Contrib. to Knowledge No. 241. 262 pp. 21 Pl.
- Woronichin, N. N. 1923. Algae nonnullae novae e Caucaso I. Notulae Syst. ex Inst. Crypt. Horti Bot. Petropolitani, Petrograd. 2: 97-100.
- Zeller, G. 1873. Algae collected by Mr. S. Kurz in Arracan and British Burma. Jour. Asiatic Soc. of Bengal 42 (2): 175-193.
- Zeller, G. 1873a. In Hedwigia 1873 (12): 189.
- Zeller, G. 1876. Algae Brasilienses circa Rio de Janeiro a Dr. A. Glaziou. Warming's Symbolae ad Flor. Bras. Central Cognos. 22: 426-432. Vidensk. Medd. Nat. (1876).

EXSICCATAE

- Areschoug, J. E. *Algae Scandinavicae exsiccatae*. Upsala.
- Collins, F. S., I. Holden, and W. A. Setchell. *Phycotheca Boreali-Americana*. (P. B. A.). Malden, Massachusetts.
- Desmazieres, J. B. H. J. *Plantes cryptogames de France*. Lille.
- Farlow, W. G., C. L. Anderson, and D. C. Eaton. *Algae Am. Bor. Exs.* Cambridge.
- Rabenhorst, L. *Die Algen Sachsens resp. Europa's*. Dresden.
- Roumeguere, C. *Les Algues fluviales et terrestres de France*. Toulouse.
- Tilden, Josephine E. *American Algae*. Minneapolis.
- Wittrock, V. B., and O. Nordstedt. *Algae aquae dulcis exs.* Upsala and Stockholm.
- Wittrock, V. B., O. Nordstedt, and G. Lagerheim. *Algae aquae dulcis exs.* Lund.

INDEX*

BULBOCHAETE

	Page		Page
<i>affinis</i> Hirn.....	48	<i>megastoma</i> Wittr. and Lund.....	31
<i>alabamensis</i> Transeau and Brown...	40	<i>minor</i> Al. Br.....	47
<i>angulosa</i> Wittr. and Lund.....	36	<i>v. germanica</i> Hirn.....	47
<i>f. picta</i> Hirn.....	37	<i>minuta</i> West and West.....	33
<i>anomala</i> Pringsh.....	48	<i>mirabilis</i> Wittr.....	32
		<i>f. immersa</i> (Wittr.) Hirn.....	32
<i>basispora</i> Wittr. and Lund.....	31	<i>v. gracilis</i> (Pringsh.) Hirn.....	32
<i>borealis</i> Wittr.....	37	<i>monile</i> Wittr. and Lund.....	30
<i>brebissonii</i> Kuetz.....	33	<i>v. robusta</i> Hirn.....	30
<i>v. minor</i> Woron.....	34	<i>nana</i> Wittr.....	30
<i>brevifulta</i> Wittr.....	50	<i>nordstedtii</i> Wittr.....	37
<i>bullardi</i> Trans. and Tiff.....	47	<i>f. suberecta</i> Collins.....	37
		<i>v. minor</i> Woron.....	37
<i>congener</i> Hirn.....	42	<i>obliqua</i> Lund.....	39
<i>crassa</i> Pringsh.....	41	<i>polyandria</i> Cleve.....	38
<i>crassiuscula</i> Nordst.....	41	<i>f. notabilis</i> Hirn.....	38
<i>crenulata</i> Pringsh.....	34	<i>punctulata</i> (Nord.) Hirn.....	41
<i>debaryana</i> Wittr. and Lund.....	31	<i>pygmaea</i> Pringsh. Wittr.....	43
<i>denticulata</i> Wittr.....	48	<i>pyrulum</i> Lund.....	43
<i>diamesandria</i> Nordst. and Hirn.....	38	<i>quadrata</i> Wittr.....	42
<i>dispar</i> Wittr.....	40	<i>rectangularis</i> Wittr.....	45
<i>v. ripartiana</i> Wittr.....	41	<i>v. hiloensis</i> Nordst.....	45
<i>doliiformis</i> Borge.....	43	<i>repanda</i> Wittr.....	44
<i>elachistandria</i> Wittr.....	39	<i>rhadinospora</i> Wittr.....	46
<i>elator</i> Pringsh.....	32	<i>f. antiqua</i> (Nord.) Hirn.....	46
<i>f. pumila</i> Hirn.....	33	<i>v. litoralis</i> Hirn.....	47
<i>v. scrobiculata</i> Tiffany.....	33	<i>sanguinea</i> Hansg.....	50
<i>furberæ</i> Collins.....	34	<i>sessilis</i> Wittr.....	39
<i>gigantea</i> Pringsh.....	38	<i>f. glabra</i> Hirn.....	39
<i>horrida</i> Nordst.....	50	<i>setigera</i> (Roth) Agard.....	40
<i>imperialis</i> Wittr.....	49	<i>spirogranulata</i> West and West.....	49
<i>v. regalis</i> Wittr.....	49	<i>subintermedia</i> Elfv.....	36
<i>insignis</i> Pringsh.....	48	<i>tenuis</i> (Wittr.) Hirn.....	43
<i>v. reticulata</i> (Nord.) Hirn.....	49	<i>v. norvegica</i> (Wittr.) Hirn.....	44
<i>intermedia</i> De Bary.....	35	<i>valida</i> Wittr.....	42
<i>v. depressa</i> Wittr.....	35	<i>varians</i> Wittr.....	45
<i>lagoensis</i> Wittr.....	46	<i>v. hawaiiensis</i> Nordst.....	46
		<i>v. subsimplex</i> (Wittr.) Hirn.....	45

OEDOCLADIUM

<i>albemarlensis</i> Lewis..... 51	<i>*media</i> Lewis..... 52
<i>*hazenii</i> Lewis..... 52	<i>protonema</i> Stahl..... 51

*New species, varieties, and forms or new combinations are indicated by an asterisk.

OEDOGONIUM

	Page		Page
<i>acmandrium</i> Elfv.....	103	<i>concatenatum</i> (Hass.) Wittr.....	136
<i>acrosporum</i> De Bary.....	157	<i>f. hulchinsiae</i> (Wittr.) Hirn.....	136
<i>f. boreale</i> (Wolle) Hirn.....	158	<i>v. rectangulare</i> Rich.....	137
<i>v. b. nidosporum</i> (Nord.) Hirn.....	158	<i>confertum</i> Hirn.....	141
<i>v. floridense</i> Wolle.....	158	<i>consociatum</i> Collins and Hervey.....	167
<i>v. majusculum</i> Nordst.....	159	<i>contortum</i> Hallas.....	147
<i>ahlstrandii</i> Wittr.....	115	<i>*costatum</i> Transeau.....	146
<i>alternans</i> Wittr. and Lund.....	128	<i>crassiusculum</i> Wittr.....	129
<i>americanum</i> Transeau.....	97	<i>v. arechavaleta</i> (Wtr.) Hirn.....	130
<i>angustissimum</i> West and West.....	160	<i>v. idioandrosporum</i> Nor. & Wittr.....	130
<i>anomalum</i> Hirn.....	83	<i>*v. cataractum</i> (Wolle) Tiff.....	130
<i>arcysporum</i> Nord. and Hirn.....	94	<i>crassum</i> (Hass.) Wittr.....	88
<i>areolatum</i> Lagerh.....	94	<i>f. amplum</i> (M. and W.) Hirn.....	88
<i>areschougii</i> Wittr.....	140	<i>v. longum</i> Transeau.....	88
<i>v. americanum</i> Tiffany.....	141	<i>v. sublumidum</i> Hirn.....	88
<i>f. robustum</i> Hirn.....	141	<i>crenulatocostatum</i> Wittr.....	92
<i>argenteum</i> Hirn.....	96	<i>f. cylindricum</i> Hirn.....	93
<i>*f. michiganese</i> Tiffany.....	97	<i>v. aureum</i> Tilden.....	93
<i>armigerum</i> Hirn.....	133	<i>v. longiarticulatum</i> Hansg.....	93
<i>aster</i> Wittr.....	120	<i>crenulatum</i> Wittr.....	171
<i>australianum</i> Hirn.....	99	<i>v. gracilius</i> (Nordst.) Hirn.....	171
<i>autumnale</i> Wittr.....	112	<i>crispum</i> (Hass.) Wittr.....	113
<i>bahusiense</i> Nordst.....	142	<i>f. granulosum</i> (Nordst.) Hirn.....	113
<i>bengalense</i> Hirn.....	139	<i>f. inflatum</i> Hirn.....	113
<i>biforme</i> Nordst.....	83	<i>v. gracilescens</i> Wittr.....	114
<i>bohemicum</i> Hirn.....	105	<i>v. hawaiiense</i> Nordst.....	114
<i>boreale</i> Hirn.....	147	<i>v. uruguayense</i> Magnus & Wille.....	114
<i>borisianum</i> (Le Cl.) Wittr.....	128	<i>cryptoporum</i> Wittr.....	65
<i>v. westii</i> Tiffany and Brown.....	128	<i>v. vulgare</i> Wittr.....	66
<i>boscii</i> (Le Cl.) Wittr.....	91	<i>curtum</i> Wittr. and Lund.....	74
<i>f. dispar</i> Hirn.....	91	<i>curvum</i> Pringsh.....	65
<i>v. notabile</i> Lemm.....	92	<i>cyathigerum</i> Wittr.....	133
<i>v. occidentale</i> Hirn.....	92	<i>f. americanum</i> Wolle.....	134
<i>brasilense</i> Borge.....	140	<i>f. latum</i> (Wittr.) Hirn.....	134
<i>braunii</i> Kuetz.; Pringsh.....	125	<i>f. perfectum</i> Hirn.....	134
<i>v. hafniense</i> (Hallas) Hirn.....	125	<i>v. ellipticum</i> Magn. and Wille.....	134
<i>v. zehneri</i> Tiffany.....	125	<i>v. hormosporum</i> (West) Hirn.....	135
<i>calcareum</i> Cleve.....	67	<i>cymatosporum</i> Wittr. & Nordst.....	67
<i>calvum</i> Wittr.....	169	<i>danicum</i> Hallas.....	145
<i>capense</i> Nordst and Hirn.....	166	<i>decipiens</i> Wittr.....	145
<i>capillare</i> (L.) Kuetz.....	80	<i>f. dissimile</i> Hirn.....	145
<i>f. stagnale</i> (Ktz.; Wittr.) Hirn.....	80	<i>v. africanum</i> Tiffany.....	145
<i>capilliforme</i> Kuetz.; Wittr.....	81	<i>v. bernardense</i> (Bates) Hirn.....	146
<i>f. debaryanum</i> (Chmiel.) Hirn.....	81	<i>depressum</i> Pringsh.....	124
<i>f. lorentzii</i> (M. and W.) Hirn.....	81	<i>dictyosporum</i> Wittr.....	94
<i>v. australe</i> Wittr.....	82	<i>f. westii</i> Tiffany.....	94
<i>v. diversum</i> (Hirn) Tiffany.....	82	<i>dioicum</i> Carter.....	107
<i>capitellatum</i> Wittr.....	104	<i>donnellii</i> Wolle; Hirn.....	132
<i>cardiacum</i> (Hass.) Wittr.....	70	<i>v. wittrockianum</i> Hirn.....	132
<i>f. interjectum</i> Hirn.....	71	<i>echinatum</i> (Wood) Wittr.....	133
<i>f. pulchellum</i> (Hass.) Hirn.....	71	<i>echinospermum</i> Al. Br.....	121
<i>v. carbonicum</i> Wittr.....	71	<i>v. horridum</i> Hirn.....	122
<i>v. minus</i> Lemm.....	71	<i>elegans</i> West and West.....	138
<i>ciliatum</i> (Hass.) Pringsh.....	153	<i>excisum</i> Wittr. and Lund.....	100
<i>clavatum</i> Hallas.....	141	<i>exocostatum</i> Tiffany.....	89
<i>cleveanum</i> Wittr.....	138	<i>exspirale</i> Tiffany.....	122
<i>f. exoticum</i> Hirn.....	138		

	Page
<i>fabulosum</i> Hirn.....	83
<i>v. columbianum</i> G. S. West.....	84
<i>v. maximum</i> (West) Hirn.....	83
<i>v. punctatum</i> Lemm.....	84
<i>flavescens</i> (Hass.) Wittr.....	125
<i>flexuosum</i> Hirn.....	163
<i>fonticola</i> Al. Braun.....	163
<i>foveolatum</i> Wittr.....	95
<i>fragile</i> Wittr.....	75
<i>v. abyssinicum</i> Hirn.....	75
<i>*v. robustum</i> (West) Tiffany.....	75
<i>franklinianum</i> Wittr.....	71
<i>fusus</i> Hallas.....	162
<i>gallicum</i> Hirn.....	124
<i>geniculatum</i> Hirn.....	78
<i>giganteum</i> Kuetz.; Wittr.....	165
<i>glabrum</i> Hallas.....	72
<i>globosum</i> Nordst.....	74
<i>gracillimum</i> Wittr. and Lund.....	116
<i>f. majus</i> West and West.....	116
<i>grande</i> Kuetz.; Wittr.....	85
<i>f. gemelliparum</i> (Pringsh.) Hirn.....	85
<i>v. aequatoriale</i> Wittr.....	85
<i>v. angustum</i> Hirn.....	86
<i>v. majus</i> Hansg.....	86
<i>gunnii</i> Wittr.....	110
<i>hians</i> Nordst. and Hirn.....	149
<i>hircii</i> Gutwinski.....	73
<i>v. africanum</i> G. S. West.....	73
<i>hispidum</i> Nordst.....	137
<i>hoehnei</i> Borge.....	165
<i>hoersholmiense</i> Hallas.....	150
<i>howardii</i> G. S. West.....	101
<i>v. minus</i> Tiffany.....	102
<i>humili</i> Wood.....	137
<i>hystericum</i> Trans. and Tiffany.....	120
<i>hystrix</i> Wittr.....	120
<i>v. canadense</i> Tiffany.....	121
<i>v. subglobosum</i> Wittr.....	121
<i>illinoisense</i> Transeau.....	123
<i>implexum</i> Hirn.....	153
<i>inclusum</i> Hirn.....	169
<i>inconspicuum</i> Hirn.....	160
<i>indicum</i> Hirn.....	139
<i>inermis</i> Hirn.....	166
<i>v. mentiens</i> Hirn.....	166
<i>infimum</i> Tiffany.....	101
<i>inflatum</i> Hallas.....	168
<i>insigne</i> Hirn.....	100
<i>intermedium</i> Wittr.....	72
<i>v. fennicum</i> Tiffany.....	73
<i>inversum</i> Wittr.....	100
<i>iowense</i> Tiffany.....	106
<i>irregularis</i> Wittr.....	131
<i>v. condensatum</i> (Hallas) Hirn.....	132
<i>itzigsohnii</i> De Bary.....	117
<i>f. heteromorphum</i> Hirn.....	118
<i>v. minus</i> West.....	118
<i>kirchneri</i> Wittr.....	116
<i>kitutæ</i> G. S. West.....	129

	Page
<i>kjellmanii</i> Wittr.; Hirn.....	89
<i>kurzii</i> Zeller.....	79
<i>laetevirens</i> Wittr.....	150
<i>laeve</i> Wittr.....	65
<i>lageniforme</i> Hirn.....	168
<i>lagerstedtii</i> Wittr.....	170
<i>landsboroughi</i> (Hass.) Wittr.....	87
<i>v. norvegicum</i> Wittr.....	87
<i>latiusculum</i> Tiffany.....	102
<i>lautumnarium</i> Wittr.....	72
<i>leiopleurum</i> Nordst. and Hirn.....	91
<i>lindmanianum</i> Wittr.....	133
<i>londinense</i> Wittr.....	167
<i>longatum</i> Kuetz.....	148
<i>longicolle</i> Nordst.....	144
<i>v. senegalense</i> Nordst.....	144
<i>loricatum</i> Hirn.....	111
<i>macrandrium</i> Wittr.....	151
<i>f. acuminatum</i> Hirn.....	151
<i>f. aemulans</i> Hirn.....	151
<i>f. lundense</i> (Wittr.) Hirn.....	151
<i>v. propinquum</i> (Wittr.) Hirn.....	152
<i>v. hokenackerii</i> (Wittr.) Tiffany.....	152
<i>v. scrobiculatum</i> Ackley.....	152
<i>macrospermum</i> West and West.....	146
<i>magnusii</i> Wittr.....	68
<i>mammiferum</i> Wittr.; Nordst.....	118
<i>manschuricum</i> Skvortzow.....	126
<i>margaritifera</i> Nordst. and Hirn.....	92
<i>martinicense</i> Hirn.....	79
<i>megaporum</i> Wittr.....	142
<i>mexicanum</i> Wittr.....	85
<i>michiganense</i> Tiffany.....	157
<i>minus</i> Wittr.....	103
<i>mirandrium</i> Skuja.....	147
<i>mitratum</i> Hirn.....	105
<i>monile</i> Berk. & Harv.; Wittr.....	154
<i>f. borgei</i> Hirn.....	154
<i>v. eminens</i> Hirn.....	154
<i>f. victoriense</i> G. S. West.....	154
<i>moniliforme</i> Wittr.....	166
<i>montagnei</i> Fiorini-Mazzanti; Wittr.....	162
<i>multisporum</i> Wood.....	131
<i>v. magnum</i> Ackley.....	131
<i>nanum</i> Wittr.; Tiffany.....	109
<i>nebraskense</i> Ohashi.....	119
<i>nobile</i> Wittr.....	99
<i>v. minus</i> Hirn.....	100
<i>nodulosum</i> Wittr.....	108
<i>v. commune</i> Hirn.....	108
<i>obesum</i> (Wittr.) Hirn.....	112
<i>oblongellum</i> Kirch.....	116
<i>oblongum</i> Wittr.....	117
<i>f. majus</i> (Nordst.) Hirn.....	117
<i>f. sphaericum</i> (Hallas) Hirn.....	117
<i>oboviforme</i> Wittr.....	86
<i>obsoletum</i> Wittr.....	69
<i>obtruncatum</i> Wittr.....	155
<i>v. completum</i> Hirn.....	155
<i>v. ellipsoideum</i> Wittr.....	155

	Page		Page
<i>oelandicum</i> Wittr.; Hirn.....	142	<i>rupestre</i> Hirn.....	114
<i>f. minus</i> Borge.....	142	<i>f. pseudautumnale</i> Hirn.....	115
<i>oryzæ</i> Wittr.....	163	<i>sancti thomæ</i> Wittr. and Cleve.....	161
<i>v. seriosporum</i> (Lager.) Hirn.....	163	<i>saxatile</i> Hansg.....	171
<i>oviforme</i> (Lewin) Hirn.....	78	<i>schmidlei</i> Gutwinski.....	148
<i>pachyandrium</i> Wittr.....	84	<i>scrobiculatum</i> Wittr.....	95
<i>pachydermum</i> Wittr. and Lund.....	115	<i>selandiæ</i> Hallas.....	161
<i>paludosum</i> (Hass.) Wittr.....	90	<i>semiapertum</i> Nordst. and Hirn.....	124
<i>v. americanum</i> Nordst.....	90	<i>sexangulare</i> Cleve.....	126
<i>v. parvisporum</i> Hirn.....	90	<i>v. angulosum</i> (Hallas) Hirn.....	126
<i>paucocostatum</i> Transeau.....	98	<i>v. majus</i> Wille.....	127
<i>v. gracilis</i> Tiffany.....	98	<i>silvaticum</i> Hallas.....	127
<i>paulense</i> Nordst. and Hirn.....	90	<i>simplex</i> Hirn.....	111
<i>perspicuum</i> Hirn.....	141	<i>sociale</i> Wittr.....	67
<i>petri</i> Wittr.....	105	<i>sodiroanum</i> Lagerheim.....	77
<i>pilosporum</i> West.....	165	<i>sol</i> Hirn.....	167
<i>pisanum</i> Wittr.....	109	<i>spectabile</i> Hirn.....	153
<i>v. gracilis</i> Trans. and Tiffany.....	110	<i>sphaerandrium</i> Wittr. and Lund.....	112
<i>pitthophoræ</i> Wittr.....	112	<i>spirale</i> Hirn.....	123
<i>plagiostomum</i> Wittr.....	80	<i>v. acutum</i> G. S. West.....	123
<i>v. gracilius</i> Wittr.....	81	<i>v. latviense</i> Tiffany.....	123
<i>platygynum</i> Wittr.....	143	<i>spurium</i> Hirn.....	164
<i>f. obtusum</i> Hirn.....	143	<i>stellatum</i> Wittr.....	127
<i>v. continuum</i> Nordst.....	143	<i>suboctangulare</i> G. S. West.....	79
<i>v. novæ zelandiæ</i> Hirn.....	144	<i>subrectum</i> Hirn.....	87
<i>plicatulum</i> Wittr.....	170	<i>suecicum</i> Wittr.....	68
<i>plusiosporum</i> Wittr.....	70	<i>f. australe</i> G. S. West.....	68
<i>pluviale</i> Nordst.....	150	<i>supremum</i> Tiffany.....	156
<i>poecilosporum</i> Nordst. and Hirn.....	168	<i>tapeinosporum</i> Wittr.....	159
<i>porrectum</i> Nordst. and Hirn.....	109	<i>taphrosporum</i> Nordst. and Hirn.....	97
<i>por tense</i> Transeau.....	102	<i>tentoriale</i> Nordst. and Hirn.....	157
<i>praticolum</i> Transeau.....	156	<i>tenuissimum</i> Hansg.....	160
<i>princeps</i> (Hass.) Wittr.....	82	<i>tiffanii</i> Ackley.....	95
<i>pringsheimii</i> Cramer; Wittr.....	107	<i>trioicum</i> Woronichin.....	106
<i>v. abbreviatum</i> Hirn.....	107	<i>tumidulum</i> (Kuetz.) Wittr.....	99
<i>v. nordstedtii</i> Wittr.....	108	<i>tyrolicum</i> Wittr.....	70
<i>psaegmatosporum</i> Nordst.....	103	<i>uleanum</i> Hirn.....	170
<i>pseudacrosporum</i> Wittr.....	98	<i>undulatum</i> (Breb.) Al. Br.; Wittr.....	118
<i>pseudoboscii</i> Hirn.....	76	<i>v. americanum</i> Transeau.....	119
<i>pulchrum</i> Nordst. and Hirn.....	170	<i>f. senegalense</i> (Nordst.) Hirn.....	119
<i>punctatostriatum</i> De Bary.....	104	<i>upsaliense</i> Wittr.....	77
<i>punctatum</i> Wittr.....	96	<i>v. fennicum</i> Hirn.....	78
<i>pungens</i> Hirn.....	122	<i>urbicum</i> Wittr.....	69
<i>pusillum</i> Kirchner.....	161	<i>urceolatum</i> Nordst. and Hirn.....	167
<i>pyriforme</i> Wittr.....	110	<i>varians</i> Wittr. and Lund.....	69
<i>pyrulum</i> Wittr.....	111	<i>vaucherii</i> (Le Cl.) Al. Br.; Wittr.....	76
<i>v. amplius</i> Taylor.....	111	<i>velatum</i> Hallas.....	164
<i>quadratum</i> Hallas.....	105	<i>verrucosum</i> Hallas.....	95
<i>reinschii</i> Roy.....	172	<i>vesicatum</i> (Lyngbye) Wittr.....	162
<i>reticulatum</i> West and West.....	171	<i>victoriense</i> G. S. West.....	129
<i>rhodosporum</i> (Welwitsch) Wittr.....	162	<i>virceburgense</i> Hirn.....	164
<i>richterianum</i> Lemm.....	76	<i>wabashense</i> Tiffany.....	156
<i>rigidum</i> Hirn.....	139	<i>warmingianum</i> Wittr.....	169
<i>riculare</i> (Le. Cl.) Al. Br.....	84	<i>welwitschii</i> West.....	106
<i>rothii</i> (Le. Cl.) Pringsheim.....	144	<i>wolleanum</i> Wittr.....	135
<i>rufescens</i> Wittr.....	66	<i>f. insigne</i> (Nordst.) Hirn.....	136
<i>*v. exiguum</i> (Elfv.) Tiffany.....	66	<i>v. concinnum</i> Hirn.....	135
<i>v. lundellii</i> (Wittr.) Tiffany.....	66	<i>wyliei</i> Tiffany.....	97
<i>rugulosum</i> Nordst.....	148	<i>zigzag</i> Cleve.....	74
<i>f. minutum</i> (Hansg.) Hirn.....	149		
<i>f. rotundatum</i> Hirn.....	149		

PLATES

PLATE I.

	Page
Fig. 1. <i>Bulbochaete elatior</i> Pringsheim. (Redrawn from Hirn).....	32
Fig. 2. <i>B. elatior</i> f. <i>pumila</i> Hirn. (Redrawn from Hirn).....	33
Fig. 3. <i>B. elatior</i> v. <i>scrobiculata</i> Tiffany.....	33
Fig. 4. <i>B. crenulata</i> Pringsheim. (Redrawn from Hirn).....	34
Fig. 5. <i>B. intermedia</i> De Bary.....	35
Fig. 6. <i>B. intermedia</i> v. <i>depressa</i> Wittrock.....	35
Fig. 7. <i>B. brebissonii</i> Kuetzing. (After Hirn).....	33
Fig. 8. <i>B. furberae</i> Collins. (After Lewis, in Collins 1918a).....	34
Fig. 9. <i>B. minuta</i> West and West. (After G. S. West).....	33

	Page
Fig. 10. <i>Bulbochaete borealis</i> Wittrock. (Redrawn from Hirn).....	37
Fig. 11. <i>B. polyandria</i> Cleve, with oogonia.....	38
Fig. 12. <i>B. polyandria</i> Cleve, with androsporangia. (After Wittrock).....	38
Fig. 13. <i>B. polyandria</i> f. <i>notabilis</i> Hirn. (Redrawn from Hirn).....	38
Fig. 14. <i>B. angulosa</i> Wittrock and Lundell. (Redrawn from Hirn).....	36
Fig. 15. <i>B. subintermedia</i> Elfving. (Redrawn from Hirn).....	36
Fig. 16. <i>B. diamesandria</i> Nordstedt and Hirn. (Redrawn from Hirn).....	38
Fig. 17. <i>B. nordstedtii</i> Wittrock.....	37
Fig. 18. <i>B. nordstedtii</i> f. <i>suberecta</i> Collins.....	37

	Page
Fig. 19. <i>Bulbochaete sessilis</i> Wittrock. (Redrawn from Hirn).....	39
Fig. 20. <i>B. sessilis</i> f. <i>glabra</i> Hirn. (Redrawn from Hirn).....	39
Fig. 21. <i>B. elachistandria</i> Wittrock. (Redrawn from Pringsheim).....	39
Fig. 22. <i>B. quadrata</i> Wittrock. (Redrawn from Hirn).....	42
Fig. 23. <i>B. dispar</i> Wittrock. (Redrawn from Hirn).....	40
Figs. 24 and 25. <i>B. dispar</i> v. <i>ripartiana</i> Wittrock. (After Hirn).....	41
Fig. 26. <i>B. obliqua</i> Lundell. (After Hirn).....	39

	Page
Fig. 27. <i>Bulbochaete valida</i> Wittrock. (Redrawn from Hirn).....	42
Fig. 28. <i>B. crassiuscula</i> Nordstedt, with androsporangia. (After Nordstedt) ..	41
Fig. 29. <i>B. crassiuscula</i> Nordstedt, with oogonia. (After Hirn).....	41
Fig. 30. <i>B. pyrulum</i> Lundell. (Redrawn from Hirn).....	43
Fig. 31. <i>B. punctulata</i> (Nordstedt) Hirn. (After Nordstedt).....	41
Fig. 32. <i>B. crassa</i> Pringsheim.....	41

	Page
Figs. 33 and 34. <i>Bulbochaete congener</i> Hirn.....	42
Figs. 35 and 36. <i>B. gigantea</i> Pringsheim.....	38
Fig. 37. <i>B. setigera</i> (Roth) Agardh.....	40
Fig. 38. <i>B. alabamensis</i> Transeau and Brown.....	40

Page

- Figs. 39 and 40. *Bulbochaete doliiformis* Borge. (Redrawn from Borge)..... 43
 Fig. 41. *B. pygmaea* Pringsheim..... 43
 Fig. 42. *B. horrida* Nordstedt. (Redrawn from Hirn)..... 50
 Fig. 43. *B. rectangularis* Wittrock..... 45
 Fig. 44. *B. rectangularis* v. *hiloensis* Nordstedt. (After Hirn)..... 45
 Fig. 45. *B. anomala* Pringsheim. (After Pringsheim)..... 48
 Fig. 46. *B. lagoensis* Wittrock. (Redrawn from Hirn)..... 46
 Fig. 47. *B. repanda* Wittrock..... 44
 Fig. 48. *B. varians* Wittrock..... 45
 Fig. 49. *B. varians* v. *subsimplax* (Wittrock) Hirn..... 45
 Fig. 50. *B. varians* v. *hawaiiensis* Nordstedt..... 46
 Fig. 51. *B. denticulata* Wittrock. (Redrawn from Hirn)..... 48

	Page
Figs. 52 and 53. <i>Bulbochaete bullardi</i> Transeau and Tiffany.....	47
Fig. 54. <i>B. minor</i> Al. Braun; Wittrock.....	47
Fig. 55. <i>B. minor</i> v. <i>germanica</i> Hirn. (After Hirn).....	47
Fig. 56. <i>B. brevifulta</i> Wittrock. (Redrawn from Hirn).....	50
Figs. 57 and 58. <i>B. spirogranulata</i> West and West. (Redrawn from G. S. West).....	49
Fig. 59. <i>B. tenuis</i> (Wittrock) Hirn. (Redrawn from Hirn).....	43
Fig. 60. <i>B. tenuis</i> v. <i>norvegica</i> (Wittrock) Hirn. (Redrawn from Hirn).....	44
Fig. 61. <i>B. affinis</i> Hirn. (Redrawn from Hirn).....	48

	Page
Fig. 62. <i>Bulbochaete insignis</i> Pringsheim.....	48
Fig. 63. <i>B. insignis</i> v. <i>reticulata</i> (Nordstedt) Hirn. (After Hirn).....	49
Fig. 64. <i>B. imperialis</i> Wittrock. (Redrawn from Hirn).....	49
Fig. 65. <i>B. imperialis</i> v. <i>regalis</i> Wittrock. (Redrawn from Hirn).....	49
Fig. 66. <i>B. rhadinospora</i> Wittrock. (After Hirn).....	46
Fig. 67. <i>B. rhadinospora</i> f. <i>antiqua</i> (Nordstedt) Hirn. (After Hirn).....	46
Fig. 68. <i>B. rhadinospora</i> v. <i>litoralis</i> Hirn. (After Hirn).....	47

	Page
Fig. 69. <i>Bulbochaete mirabilis</i> Wittrock. (Redrawn from Hirn).....	32
Fig. 70. <i>B. mirabilis</i> f. <i>immersa</i> (Wittrock) Hirn. (Redrawn from Hirn).....	32
Fig. 71. <i>B. mirabilis</i> v. <i>gracilis</i> (Pringsheim) Hirn. (Redrawn from Hirn)....	32
Fig. 72. <i>B. monile</i> Wittrock and Lundell. (Redrawn from Hirn).....	30
Fig. 73. <i>B. monile</i> v. <i>robusta</i> Hirn. (Redrawn from Hirn).....	30
Fig. 74. <i>B. basispora</i> Wittrock and Lundell.....	31
Fig. 75. <i>B. nana</i> Wittrock. (Redrawn from Hirn).....	30
Fig. 76. <i>B. megastoma</i> Wittrock and Lundell. (After Wittrock).....	31
Fig. 77. <i>B. debaryana</i> Wittrock and Lundell. (Redrawn from Hirn).....	31

Figs. 78-83. Germination of oospore of *B. elachistandria* Wittrock. (After Pringsheim)

Figs. 84-92. Development and germination of zoospores of *B. varians* Wittrock v. *subsimplex* (Wittrock) Hirn.

Figs. 93-98. Development of bristle and first cell division of a germinated zoospore of *B. setigera* (Roth) Agardh. (After Pringsheim).

Fig. 99. A young plant of *B. setigera* (Roth) Agardh showing manner of branching: a, Primary filament developed from zoospore. b, Branch of the first order. c, d, e, Branches of the second, third, and fourth order, respectively. (Modified from Pringsheim).

PLATE XI.

	Page
Fig. 100. <i>Oedogonium laeve</i> Wittr.....	65
Fig. 101. <i>Oe. cryptoporum</i> Wittr.....	65
Fig. 102. <i>Oe. cryptoporum</i> v. <i>vulgare</i> Wittr. (Redrawn from Hirn).....	66
Figs. 103 and 104. <i>Oe. curvum</i> Pringsh.....	65
Fig. 105. <i>Oe. rufescens</i> Wittr. (Redrawn from Hirn).....	66
Fig. 106. <i>Oe. rufescens</i> v. <i>exiguum</i> (Elfv.) Tiffany.....	66
Fig. 107. <i>Oe. rufescens</i> v. <i>lundellii</i> (Wittr.) Tiffany. (Redrawn from Hirn).....	66
Fig. 108. <i>Oe. calcareum</i> Cleve. (Redrawn from Hirn).....	67
Fig. 109. <i>Oe. sociale</i> Wittr. (Redrawn from Hirn).....	67
Fig. 110. <i>Oe. inerme</i> Hirn. (Redrawn from Hirn).....	166
Fig. 111. <i>Oe. inerme</i> v. <i>mentiens</i> Hirn. (Redrawn from Hirn).....	166

	Page
Figs. 112 and 113. <i>Oedogonium cymatosporum</i> Wittr. and Nordst. (Redrawn from Hirn).....	67
Fig. 114. <i>Oe. moniliforme</i> Wittr.....	166
Fig. 115. <i>Oe. magnusii</i> Wittr. (Redrawn from Hirn).....	68
Fig. 116. <i>Oe. suecicum</i> Wittr.....	68
Fig. 117. <i>Oe. suecicum</i> f. <i>australe</i> G. S. West. (Redrawn from West).....	68
Figs. 118 and 119. <i>Oe. urbicum</i> Wittr. (Redrawn from Pringsheim).....	69
Fig. 120. <i>Oe. varians</i> Wittr. and Lund.....	69
Fig. 121. <i>Oe. obsoletum</i> Wittr. (Redrawn from Hirn).....	69
Fig. 122. <i>Oe. plusiosporum</i> Wittr. (Redrawn from Hirn).....	70
Fig. 123. <i>Oe. tyrolicum</i> Wittr.....	70

	Page
Figs. 124 and 125. <i>Oedogonium cardiacum</i> (Hass.) Wittr. (Redrawn from Hirn).....	70
Fig. 126. <i>Oe. cardiacum</i> f. <i>pulchellum</i> (Hass.) Hirn. (Redrawn from Hirn)..	71
Fig. 127. <i>Oe. cardiacum</i> f. <i>interjectum</i> Hirn. (Redrawn from Hirn).....	71
Fig. 128. <i>Oe. cardiacum</i> v. <i>carbonicum</i> Wittr.....	71
Fig. 129. <i>Oe. cardiacum</i> v. <i>minus</i> Lemm.....	71
Fig. 130. <i>Oe. glabrum</i> Hallas. (Redrawn from Hallas).....	72
Fig. 131. <i>Oe. franklinianum</i> Wittr.....	71

	Page
Figs. 132 and 133. <i>Oedogonium lautumiarum</i> Wittr. (Redrawn from Hirn).....	72
Fig. 134. <i>Oe. intermedium</i> Wittr.....	72
Fig. 135. <i>Oe. intermedium</i> v. <i>fennicum</i> Tiffany.....	73
Fig. 136. <i>Oe. hirnii</i> Gutw. (Redrawn from W. and G. S. West).....	73
Fig. 137. <i>Oe. hirnii</i> Gutw. (Redrawn from Gutwinski).....	73
Fig. 138. <i>Oe. hirnii</i> v. <i>africanum</i> G. S. West.....	73
Fig. 139. <i>Oe. globosum</i> Nordst. (Redrawn from Hirn).....	74
Fig. 140. <i>Oe. plagiostomum</i> Wittr. (Redrawn from Hirn).....	80
Figs. 141 and 142. <i>Oe. plagiostomum</i> v. <i>gracilius</i> Wittr.....	81

	Page
Figs. 143 and 144. <i>Oedogonium zigzag</i> Cleve. (Redrawn from Hirn).....	74
Fig. 145. <i>Oe. fragile</i> Wittr.....	75
Fig. 146. <i>Oe. fragile</i> v. <i>robustum</i> (W. and W.) Tiffany. (Redrawn from West).....	75
Fig. 147. <i>Oe. fragile</i> v. <i>abyssinicum</i> Hirn. (Redrawn from Hirn).....	75
Fig. 148. <i>Oe. warmingianum</i> Wittr. (Redrawn from Hirn).....	169
Fig. 149. <i>Oe. curtum</i> Wittr. and Lund. (Redrawn from Hirn).....	74
Figs. 150 and 151. <i>Oe. vaucherii</i> (Le Cl.) A. Br.; Wittr.....	76

	Page
Fig. 152. <i>Oedogonium oviforme</i> (Lewin) Hirn.....	78
Figs. 153 and 154. <i>Oe. sodiroanum</i> Lagerh.....	77
Fig. 155. <i>Oe. geniculatum</i> Hirn.....	78
Fig. 156. <i>Oe. richterianum</i> Lemm. (Redrawn from Hirn).....	76
Fig. 157. <i>Oe. upsaliense</i> Wittr.....	77
Fig. 158. <i>Oe. upsaliense</i> v. <i>fennicum</i> Hirn. (Redrawn from Hirn).....	78

	Page
Fig. 159. <i>Oedogonium martinicense</i> Hirn. (Redrawn from Hirn).....	79
Fig. 160. <i>Oe. suboctangulare</i> West and West. (Redrawn from West).....	79
Figs. 161 and 162. <i>Oe. biforme</i> Nordst. (Redrawn from Hirn).....	83
Fig. 163. <i>Oe. kurzii</i> Zeller. (Redrawn from Hirn).....	79

Figs. 164 and 165. <i>Oedogonium capillare</i> (L.) Kuetz.....	Page 80
Fig. 166. <i>Oe. capillare</i> f. <i>stagnale</i> (Kuetz.; Wittr.) Hirn. (Redrawn from Hirn).....	80
Figs. 167-169. <i>Oe. princeps</i> (Hass.) Wittr.....	82
Figs. 170 and 171. <i>Oe. anomalum</i> Hirn.....	83

	Page
Figs. 172 and 173. <i>Oe. capilliforme</i> Kuetz.; Wittr.....	81
Fig. 174. <i>Oe. capilliforme</i> f. <i>debaryanum</i> (Chmiel.) Hirn.....	81
Fig. 175. <i>Oe. capilliforme</i> f. <i>lorentzii</i> (M. & W.) Hirn. (Redrawn from Hirn) ..	81
Figs. 176 and 177. <i>Oe. capilliforme</i> v. <i>australe</i> Wittr.....	82
Fig. 178. <i>Oe. capilliforme</i> v. <i>diversum</i> (Hirn) Tiffany. (Redrawn from Hirn).....	82

	Page
Figs. 179 and 180. <i>Oedogonium pachyandrium</i> Wittr. (Redrawn from Hirn) ..	84
Figs. 181-183. <i>Oe. subrectum</i> Hirn. (Redrawn from Hirn)	87
Figs. 184 and 185. <i>Oe. oboviforme</i> Wittr. (Redrawn from Hirn)	86
Fig. 186. <i>Oe. pseudoboscii</i> Hirn. (Redrawn from Silfvenius)	76
Fig. 187. <i>Oe. lageniforme</i> Hirn. (Redrawn from Hirn)	168

	Page
Figs. 188 and 189. <i>Oe. mexicanum</i> Wittr. (Redrawn from Hirn).....	85
Fig. 190. <i>Oe. oryzae</i> Wittr. (Redrawn from Hirn).....	163
Figs. 191 and 192. <i>Oe. oryzae</i> v. <i>seriosporum</i> (Lagerh.) Hirn. (Redrawn from Hirn).....	164
Figs. 193 and 194. <i>Oe. inflatum</i> Hallas. (Redrawn from Hallas).....	168

	Page
Figs. 195 and 196. <i>Oedogonium grande</i> Kuetz.; Wittr.....	85
Fig. 197. <i>Oe. grande</i> v. <i>majus</i> Hansg.....	86
Figs. 198 and 199. <i>Oe. grande</i> v. <i>angustum</i> Hirn. (Redrawn from Hirn).....	86
Fig. 200. <i>Oe. grande</i> f. <i>gemelliparum</i> (Pringsh.) Hirn. (Redrawn from Hirn).....	85
Fig. 201. <i>Oe. grande</i> v. <i>aequatoriale</i> Wittr. (Redrawn from Hirn).....	85

PLATE XXIII.

	Page
Figs. 202 and 203. <i>Oedogonium crassum</i> (Hass.) Wittr.....	88
Figs. 204 and 205. <i>Oe. crassum</i> f. <i>amplum</i> (M. & W.) Hirn. (Redrawn from Hirn).....	88
Figs. 206 and 207. <i>Oe. crassum</i> v. <i>subtumidum</i> Hirn. (Redrawn from Hirn).....	88

	Page
Fig. 208. <i>Oedogonium crassum</i> v. <i>longum</i> Transeau.....	88
Figs. 209 and 210. <i>Oe. rivulare</i> (Le CL.) Al. Br. (Redrawn from Hirn).....	84
Figs. 211 and 212. <i>Oe. landsboroughi</i> (Hass.) Wittr.....	87
Figs. 213 and 214. <i>Oe. landsboroughi</i> v. <i>norvegicum</i> Wittr.....	87

Page
Figs. 215 and 216. *Oedogonium fabulosum* Hirn. (Redrawn from Hirn)..... 83
Figs. 217 and 218. *Oe. fabulosum* v. *maximum* (West) Hirn. (Redrawn from West)..... 83
Fig. 219. *Oe. fabulosum* v. *columbianum* G. S. West. (Redrawn from West). 84
Fig. 220. *Oe. angustissimum* West and West. (Redrawn from West).....160

(This plate presents a comparison between the largest and smallest species of *Oedogonium*.)

	Page
Figs. 221 and 222. <i>Oedogonium leiopleurum</i> Nord. and Hirn. (Redrawn from Hirn).....	91
Fig. 223. <i>Oe. kjellmanii</i> Wittr.; Hirn. (Redrawn from Hirn).....	89
Fig. 224. <i>Oe. paulense</i> Nord. and Hirn. (Redrawn from Schmidle).....	90
Figs. 225 and 226. <i>Oe. boscii</i> (Le Cl.) Wittr.....	91
Figs. 227 and 228. <i>Oe. boscii</i> v. <i>occidentale</i> Hirn. (Redrawn from Hirn)....	92
Fig. 229. <i>Oe. boscii</i> v. <i>notabile</i> Lemm. (Redrawn from Hirn).....	92
Fig. 230. <i>Oe. boscii</i> f. <i>dispar</i> Hirn. (Redrawn from Hirn).....	91
Fig. 231. <i>Oe. paludosum</i> (Hass.) Wittr.....	90
Fig. 232. <i>Oe. paludosum</i> v. <i>parvisporum</i> Hirn.....	90

	Page
Fig. 233. <i>Oedogonium paludosum</i> v. <i>americanum</i> Nord. (Redrawn from Hirn).....	90
Fig. 234. <i>Oe. margaritiferum</i> Nord. and Hirn. (Redrawn from Hirn).....	92
Figs. 235-237. <i>Oe. exocostatum</i> Tiffany.....	89
Figs. 238 and 239. <i>Oe. urceolatum</i> Nord. and Hirn. (Redrawn from Hirn)....	167

	Page
Figs. 240 and 241. <i>Oedogonium crenulato-costatum</i> Wittr.....	92
Figs. 242 and 243. <i>Oe. crenulato-costatum</i> v. <i>longi-articulatum</i> Hansg.....	93
Fig. 244. <i>Oe. crenulato-costatum</i> f. <i>cylindricum</i> Hirn.....	93
Fig. 245. <i>Oe. crenulato-costatum</i> v. <i>aureum</i> Tilden. (Redrawn from Hirn)...	93
Fig. 246. <i>Oe. dictyosporum</i> Wittr. (Redrawn from Hirn).....	94
Fig. 247. <i>Oe. arcyosporum</i> Nord. and Hirn. (Redrawn from Hirn).....	94
Figs. 248-250. <i>Oe. areolatum</i> Lagerh. (Redrawn from Hirn).....	94
Fig. 251. <i>Oe. hoeheii</i> Borge. (Redrawn from Borge).....	165

	Page
Fig. 252. <i>Oedogonium foveolatum</i> Wittr. (Redrawn from Hirn).....	95
Figs. 253 and 254. <i>Oe. argenteum</i> Hirn.....	96
Figs. 255 and 256. <i>Oe. punctatum</i> Wittr. (Redrawn from Hirn).....	96
Fig. 257. <i>Oe. capense</i> Nord. and Hirn. (Redrawn from Hirn).....	166
Figs. 258 and 259. <i>Oe. scrobiculatum</i> Wittr. (Redrawn from Hirn).....	95
Fig. 260. <i>Oe. giganteum</i> Kuetz.; Wittr. (Redrawn from Hirn).....	165

Figs. 261-263. <i>Oedogonium tiffanii</i> Ackley.....	Page 95
Figs. 264-267. <i>Oe. wyliei</i> Tiffany.....	97
Figs. 268 and 269. <i>Oe. pseudacrosporum</i> Wittr. (Redrawn from Hirn).....	98

	Page
Figs. 270 and 271. <i>Oedogonium taphrosporum</i> Nord. and Hirn. (Redrawn from Hirn).....	97
Figs. 272-274. <i>Oe. verrucosum</i> Hallas. (Redrawn from Hallas).....	95
Figs. 275 and 276. <i>Oe. americanum</i> Transeau.....	97

	Page
Fig. 277. <i>Oedogonium paucocostatum</i> Transeau.....	98
Fig. 278. <i>Oe. paucocostatum</i> v. <i>gracilis</i> Tiffany.....	98
Fig. 279. <i>Oe. nobile</i> Wittr. (Redrawn from Hirn).....	99
Fig. 280. <i>Oe. nobile</i> v. <i>minus</i> Hirn. (Redrawn from Hirn).....	100
Figs. 281 and 282. <i>Oe. tumidulum</i> (Kuetz.) Wittr. (Redrawn from Hirn)....	99
Figs. 283 and 284. <i>Oe. australianum</i> Hirn. (Redrawn from Hirn).....	99

	Page
Fig. 285. <i>Oedogonium insigne</i> Hirn. (Redrawn from Hirn).....	100
Fig. 286. <i>Oe. sol</i> Hirn. (Redrawn from Hirn).....	167
Figs. 287 and 288. <i>Oe. latiusculum</i> Tiffany.....	102
Figs. 289-292. <i>Oe. infimum</i> Tiffany.....	101
Fig. 293. <i>Oe. howardii</i> G. S. West. (Redrawn from West).....	101
Fig. 294. <i>Oe. howardii</i> v. <i>minus</i> Tiffany.....	102
Figs. 295-297. <i>Oe. inversum</i> Wittr.....	100

	Page
Figs. 298 and 299. <i>Oedogonium pratense</i> Transeau.....	102
Fig. 300. <i>Oe. acmandrium</i> Elfv. (Redrawn from Hirn).....	103
Fig. 301. <i>Oe. psaeagmatosporum</i> Nordst. (Redrawn from Hirn).....	103
Fig. 302. <i>Oe. minus</i> Wittr.....	103
Figs. 303 and 304. <i>Oe. punctatostriatum</i> De Bary. (Redrawn from Hirn).....	104
Fig. 305. <i>Oe. poecilosporum</i> Nordst. and Hirn. (Redrawn from Hirn).....	168
Figs. 306-308. <i>Oe. quadratum</i> Hallas. (Redrawn from Hallas).....	105
Figs. 309 and 310. <i>Oe. calvum</i> Wittr. (Redrawn from Hirn).....	169
Fig. 311. <i>Oe. fusus</i> Hallas. (Redrawn from Hallas).....	162
Fig. 312. <i>Oe. inconspicuum</i> Hirn. (Redrawn from Nordstedt).....	160
Fig. 313. <i>Oe. inconspicuum</i> Hirn.....	160
Fig. 314. <i>Oe. tapeinosporum</i> Wittr.....	159
Fig. 315. <i>Oe. capitellatum</i> Wittr. (Redrawn from Hirn).....	104
Fig. 316. <i>Oe. pusillum</i> Kirchner.....	161
Fig. 317. <i>Oe. excisum</i> Wittr. and Lund. (Redrawn from Wittrock).....	160
Fig. 318. <i>Oe. gunnii</i> Wittr.....	110

	Page
Fig. 319. <i>Oedogonium crispum</i> (Hass.) Wittr.....	113
Fig. 320. <i>Oe. crispum</i> v. <i>hawaiense</i> Nord. (Redrawn from Hirn).....	114
Fig. 321. <i>Oe. crispum</i> v. <i>gracilescens</i> Wittr. (Redrawn from Hirn).....	114
Fig. 322. <i>Oe. crispum</i> f. <i>granulosum</i> (Nord.) Hirn. (Redrawn from Hirn).....	113
Fig. 323. <i>Oe. crispum</i> f. <i>inflatum</i> Hirn. (Redrawn from Hirn).....	113
Fig. 324. <i>Oe. crispum</i> v. <i>uruguayense</i> M. & W. (Redrawn from Hirn).....	114
Figs. 325 and 326. <i>Oe. pringsheimii</i> Cramer; Wittr.....	107
Figs. 327-330. <i>Oe. pringsheimii</i> v. <i>nordstedtii</i> Wittr.....	108
Fig. 331. <i>Oe. pringsheimii</i> v. <i>abbreviatum</i> Hirn. (Redrawn from Hirn).....	107

	Page
Fig. 332. <i>Oedogonium petri</i> Wittr. (Redrawn from Hirn).....	105
Fig. 333. <i>Oe. spurium</i> Hirn. (Redrawn from Hirn).....	164
Figs. 334 and 335. <i>Oe. mitratum</i> Hirn.....	105
Fig. 336. <i>Oe. virceburgense</i> Hirn. (Redrawn from Hirn).....	164
Fig. 337. <i>Oe. sphaerandrium</i> Wittr. and Lund. (a. After Borge. b. After Nordstedt).....	112
Fig. 338. <i>Oe. pithophorae</i> Wittr. (Redrawn from Hirn).....	112
Fig. 339. <i>Oe. loricaum</i> Hirn. (Redrawn from Hirn).....	111
Fig. 340. <i>Oe. simplex</i> Hirn. (Redrawn from Hirn).....	111
Fig. 341. <i>Oe. autumnale</i> Wittr. (Redrawn from Hirn).....	112
Figs. 342 and 343. <i>Oe. nanum</i> Wittr.; Tiffany.....	109
Fig. 344. <i>Oe. bohemicum</i> Hirn.....	105
Fig. 345. <i>Oe. rupestre</i> Hirn. (Redrawn from Hirn).....	114
Fig. 346. <i>Oe. rupestre</i> f. <i>pseudautumnale</i> Hirn. (Redrawn from Hirn).....	115

	Page
Fig. 347. <i>Oedogonium obesum</i> (Wittr.) Hirn. (Redrawn from Hirn).....	112
Fig. 348. <i>Oe. pyriforme</i> Wittr. (Redrawn from Hirn).....	110
Fig. 349. <i>Oe. pyrulum</i> Wittr. (Redrawn from Hirn).....	111
Fig. 350. <i>Oe. pyrulum</i> v. <i>amplius</i> Taylor.....	111
Figs. 351 and 352. <i>Oe. welwitschii</i> West.....	106
Figs. 353-355. <i>Oe. iowense</i> Tiffany.....	106
Figs. 356 and 357. <i>Oe. dioicum</i> Carter. (After Carter).....	107

Fig. 358.	<i>Oedogonium ahlstrandii</i> Wittr.	(Redrawn from Hirn)	Page
Fig. 359.	<i>Oe. selandiae</i> Hallas.	(After Hallas)	115
Fig. 360.	<i>Oe. sancti thomae</i> Wittr. and Cleve		161
Fig. 361 and 362.	<i>Oe. porrectum</i> Nord. and Hirn.	(Redrawn from Hirn)	109
Figs. 363 and 364.	<i>Oe. pisanum</i> Wittr.	(Redrawn from Hirn)	109
Figs. 365 and 366.	<i>Oe. pisanum</i> v. <i>gracilis</i> Trans. and Tiff.		110
Fig. 367.	<i>Oe. oblongum</i> Wittr.		117
Figs. 368 and 369.	<i>Oe. oblongum</i> f. <i>sphaericum</i> (Hallas) Hirn.	(After Hallas)	117
Fig. 370.	<i>Oe. oblongum</i> f. <i>majus</i> (Nord.) Hirn.	(Redrawn from Hirn)	117
Fig. 371.	<i>Oe. kirchneri</i> Wittr.	(Redrawn from Hirn)	116
Fig. 372.	<i>Oe. oblongellum</i> Kirch.	(After W. and G. S. West)	116
Fig. 373.	<i>Oe. gracillimum</i> Wittr. and Lund.		116

	Page
Figs. 374 and 375. <i>Oedogonium nodulosum</i> Wittr. (Redrawn from Hirn).....	108
Fig. 376. <i>Oe. nodulosum</i> v. <i>commune</i> Hirn. (Redrawn from Hirn).....	108
Figs. 377 and 378. <i>Oe. pachydermum</i> Wittr. and Lund. (Redrawn from Hirn).....	115
Figs. 379 and 380. <i>Oe. itzigsohnii</i> De Bary. (Redrawn from Hirn).....	117
Fig. 381. <i>Oe. itzigsohnii</i> f. <i>heteromorphum</i> Hirn. (Redrawn from Hirn).....	118
Fig. 382. <i>Oe. itzigsohnii</i> v. <i>minus</i> West. (Redrawn from West).....	118
Figs. 383-385. <i>Oe. mamiferum</i> Wittr.; Nord.....	118

	Page
Figs. 386 and 387. <i>Oedogonium vesicatum</i> (Lyng.) Wittr. (Redrawn from Hirn).....	162
Figs. 388 and 389. <i>Oe. rhodosporum</i> (Welw.) Wittr. (Redrawn from Hirn).....	162
Figs. 390 and 391. <i>Oe. montagnei</i> Fior.-Mazz. (Redrawn from Hirn).....	162
Figs. 392 and 393. <i>Oe. pluviale</i> Nord. (Redrawn from Hirn).....	150
Fig. 394. <i>Oe. flexuosum</i> Hirn. (Redrawn from Hirn).....	163
Fig. 395. <i>Oe. pilosporum</i> West. (Redrawn from West).....	165
Fig. 396. <i>Oe. consociatum</i> Collins and Hervey. (Redrawn from Collins)....	167
Fig. 397. <i>Oe. inclusum</i> Hirn. (Redrawn from Hirn).....	169

Figs. 398 and 399. *Oedogonium velatum* Hallas. (Redrawn from Hallas).....164
Figs. 400 and 401. *Oe. implexum* Hirn. (Redrawn from Hirn).....153
Fig. 402. *Oe. fonticola* Al. Br. (Redrawn from Hirn).....163
Figs. 403-405. *Oe. spectabile* Hirn. (Redrawn from Hirn).....153
Fig. 406. *Oe. reinschii* Roy.....172

	Page
Fig. 407. <i>Oedogonium undulatum</i> (Breb.) Al. Br.; Wittr. (Redrawn from Hirn).....	118
Fig. 408. <i>Oe. undulatum</i> v. <i>americanum</i> Trans.....	119
Fig. 409. <i>Oe. undulatum</i> f. <i>senegalense</i> (Nord.) Hirn.....	119
Figs. 410-413. <i>Oe. nebraskense</i> Ohashi (slightly on larger scale).....	119

	Page
Fig. 414. <i>Oedogonium hystricinum</i> Transeau and Tiffany.....	120
Fig. 415. <i>Oe. armigerum</i> Hirn. (Redrawn from Hirn).....	133.
Fig. 416. <i>Oe. aster</i> Wittr. (Redrawn from Hirn).....	120
Fig. 417. <i>Oe. hystrix</i> Wittr. (After Wittrock).....	120
Fig. 418. <i>Oe. hystrix</i> v. <i>canadense</i> Tiffany.....	121
Fig. 419. <i>Oe. hystrix</i> v. <i>subglobosum</i> Wittr. (After Wittrock).....	121
Figs. 420 and 421. <i>Oe. echinospermum</i> Al. Br.....	121
Fig. 422. <i>Oe. echinospermum</i> v. <i>horridum</i> Hirn. (After Wolle).....	122

	Page
Figs. 423 and 424. <i>Oedogonium pungens</i> Hirn. (Redrawn from Hirn).....	122
Figs. 425 and 426. <i>Oe. exospirale</i> Tiffany.....	122
Figs. 427 and 428. <i>Oe. spirale</i> Hirn. (Redrawn from Hirn).....	123
Fig. 429. <i>Oe. spirale</i> v. <i>latviense</i> Tiffany.....	123
Fig. 430. <i>Oe. spirale</i> v. <i>acutum</i> West and West. (Redrawn from West).....	123
Figs. 431-433. <i>Oe. illinoisense</i> Transeau. (Redrawn from Transeau).....	123

	Page
Figs. 434 and 435. <i>Oedogonium lindmanianum</i> Wittr. (Redrawn from Hirn).....	133
Fig. 436. <i>Oe. echinatum</i> (Wood) Wittr. Oospore. (After Wood).....	133
Fig. 437. <i>Oe. donnellii</i> Wille; Hirn. (Redrawn from Hirn).....	132
Figs. 438 and 439. <i>Oe. donnellii</i> v. <i>wittrockianum</i> Hirn. (Redrawn from Hirn).....	132
Fig. 440. <i>Oe. hispidum</i> Nordst. (Redrawn from Hirn).....	137
Fig. 441. <i>Oe. stellatum</i> Wittr. (Redrawn from Hirn).....	127

	Page
Fig. 442. <i>Oedogonium manschuricum</i> Skvortzow. (After Skvortzow).....	126
Fig. 443. <i>Oe. cleveanum</i> Wittr. (A, antheridium; S, dwarf male stipe).....	138
Fig. 444. <i>Oe. cleveanum</i> Wittr. (Redrawn from Hirn).....	138
Fig. 445. <i>Oe. semiapertum</i> Nord. and Hirn. (Redrawn from Hirn).....	124
Fig. 446. <i>Oe. depressum</i> Pringsh. (Redrawn from Hirn).....	124
Fig. 447. <i>Oe. irregulare</i> Wittr.	131
Figs. 448 and 449. <i>Oe. irregulare</i> v. <i>condensatum</i> (Hallas) Hirn. (After
Hallas).....	132
Figs. 450 and 451. <i>Oe. multisporum</i> Wood.....	131
Fig. 452. <i>Oe. multisporum</i> v. <i>magnum</i> Ackley.....	131

	Page
Fig. 453. <i>Oedogonium gallicum</i> Hirn. (Redrawn from Hirn).....	124
Fig. 454. <i>Oe. sexangulare</i> Cleve.....	126
Figs. 455 and 456. <i>Oe. sexangulare</i> v. <i>angulosum</i> (Hallas) Hirn. (After Hallas).....	126
Fig. 457. <i>Oe. sexangulare</i> v. <i>majus</i> Wille. (Redrawn from Hirn).....	127
Fig. 458. <i>Oe. braunii</i> Kuetz; Pringsh.....	125
Fig. 459. <i>Oe. braunii</i> v. <i>zehneri</i> Tiffany.....	125
Figs. 460-462. <i>Oe. braunii</i> v. <i>hafniense</i> (Hallas) Hirn. (After Hallas).....	125

	Page
Fig. 463. <i>Oedogonium alternans</i> Wittr. and Lund. (Redrawn from Hirn).....	128
Fig. 464. <i>Oe. contortum</i> Hallas. (After Hallas).....	147
Figs. 465-468. <i>Oe. huntii</i> Wood. (After Wood).....	137
Fig. 469. <i>Oe. borisianum</i> (Le Cl.) Wittr.....	128
Fig. 470. <i>Oe. borisianum</i> v. <i>westii</i> Tiffany and Brown.....	128

	Page
Fig. 471. <i>Oedogonium flavescent</i> (Hass.) Wittr.....	125
Fig. 472. <i>Oe. silvaticum</i> Hallas. (After Hallas).....	127
Fig. 473. <i>Oe. victoriense</i> G. S. West. (After West).....	129
Fig. 474. <i>Oe. crassiusculum</i> Wittr.....	129
Figs. 475 and 476. <i>Oe. crassiusculum</i> v. <i>idioandrospermum</i> Nord. and Wittr.....	130
Fig. 477. <i>Oe. crassiusculum</i> v. <i>cataractum</i> (Wolle) Tiffany.....	130

	Page
Fig. 478. <i>Oedogonium crassiusculum</i> v. <i>archavaletae</i> (Wittr.) Hirn. (Redrawn from Hirn).....	130
Figs. 479 and 480. <i>Oe. wolleanum</i> Wittr. (Redrawn from Hirn).....	135
Fig. 481. <i>Oe. wolleanum</i> v. <i>concinnum</i> Hirn. (Redrawn from Hirn).....	135
Figs. 482 and 483. <i>Oe. wolleanum</i> f. <i>insigne</i> (Nord.) Hirn. (Redrawn from Hirn).....	136

Figs. 484 and 485.	<i>Oedogonium cyathigerum</i> Wittr.	Page
Figs. 486 and 487.	<i>Oe. cyathigerum</i> f. <i>perfectum</i> Hirn. (Redrawn from Hirn)	133
Figs. 488-490.	<i>Oe. cyathigerum</i> f. <i>ornatum</i> (Wittr.) Hirn. (Redrawn from Hirn)	134

	Page
Figs. 491 and 492. <i>Oe. cyathigerum</i> v. <i>hormosporum</i> (West) Hirn. (Redrawn from Hirn).....	135
Fig. 493. <i>Oe. cyathigerum</i> v. <i>ellipticum</i> M. and W. (Redrawn from Hirn)....	134
Fig. 494. <i>Oe. concatenatum</i> (Hass.) Wittr.....	136
Fig. 495. <i>Oe. concatenatum</i> f. <i>hutchinsiae</i> (Wittr.) Hirn.....	136

TIFFANY

50 MICRONS

	Page
Figs. 496 and 497. <i>Oedogonium concatenatum</i> v. <i>rectangulare</i> Rich. (After Rich).....	137
Figs. 498 and 499. <i>Oe. elegans</i> West and West. (After West).....	138
Fig. 500. <i>Oe. areschougii</i> Wittr. (After Wittrock).....	140
Fig. 501. <i>Oe. areschougii</i> v. <i>americanum</i> Tiffany.....	140
Figs. 502 and 503. <i>Oe. areschougii</i> f. <i>robustum</i> Hirn. (Redrawn from Hirn).....	140
Figs. 504 and 505. <i>Oe. bengalense</i> Hirn. (Redrawn from Hirn).....	139
Figs. 506 and 507. <i>Oe. rigidum</i> Hirn. (Redrawn from Hirn).....	139

	Page
Fig. 508. <i>Oedogonium brasiliense</i> Borge. (Redrawn from Hirn).....	140
Fig. 509. <i>Oe. indicum</i> Hirn. (Redrawn from Hirn).....	139
Fig. 510. <i>Oe. confertum</i> Hirn. (Redrawn from Hirn).....	141
Fig. 511. <i>Oe. perspicuum</i> Hirn. (Redrawn from Hirn).....	141
Figs. 512 and 513. <i>Oe. clavatum</i> Hallas. (After Hallas).....	141
Fig. 514. <i>Oe. mirandrium</i> Skuja. (After Skuja).....	147
Fig. 515. <i>Oe. danicum</i> Hallas. (After Hallas).....	145

Page

- Figs. 516 and 517. *Oedogonium longicollae* Nord.; Hirn. (After Nordstedt)...144
 Figs. 518 and 519. *Oe. longicollae* v. *senegalense* Nord. (After Nordstedt)...144
 Fig. 520. *Oe. decipiens* Wittr. (After Wittrock)...145
 Figs. 521 and 522. *Oe. decipiens* f. *dissimile* Hirn.145
 Figs. 523 and 524. *Oe. decipiens* v. *africanum* Tiffany. (Redrawn from Hirn)...145
 Fig. 525. *Oe. decipiens* v. *bernardense* (Bates) Hirn. (Redrawn from Hirn)...146
 Fig. 526. *Oe. rothii* (Le Cl.) Pringsh. (After Pringsheim)...144
 Figs. 527 and 528. *Oe. macrospermum* West and West...146
 Fig. 529. *Oe. uleanum* Hirn. (Redrawn from Hirn)...170
 Fig. 530. *Oe. bahusiense* Nordst. (After Nordstedt)...142
 Fig. 531. *Oe. bahusiense* Nordst., oogonium seen from above...142

	Page
Figs. 532 and 533. <i>Oedogonium crenulatum</i> Witttr. (After Wittrock).....	171
Fig. 534. <i>Oe. crenulatum</i> v. <i>gracilius</i> (Nord.) Hirn. (Redrawn from Hirn)....	171
Fig. 535. <i>Oe. plicatum</i> Witttr. (Redrawn from Hirn).....	170
Figs. 536-538. <i>Oe. lagerstedtii</i> Witttr. (Redrawn from Hirn).....	170
Fig. 539. <i>Oe. oelandicum</i> Witttr.; Hirn.....	142
Fig. 540. <i>Oe. oelandicum</i> f. <i>minus</i> Borge. (Redrawn from Borge).....	142
Fig. 541. <i>Oe. pulchrum</i> Nord. and Hirn. (Redrawn from Hirn).....	170
Fig. 542. <i>Oe. boreale</i> Hirn. (Redrawn from Hirn).....	147
Fig. 543. <i>Oe. megaporum</i> Witttr.....	142
Fig. 544. <i>Oe. platygynum</i> Witttr. (After Wittrock).....	143
Fig. 545. <i>Oe. platygynum</i> f. <i>obtusum</i> Hirn. (Redrawn from Hirn).....	143
Fig. 546. <i>Oe. platygynum</i> v. <i>continuum</i> Nordst. (After Nordstedt).....	143
Fig. 547. <i>Oe. platygynum</i> v. <i>novae zelandiae</i> Hirn. (After Nordstedt).....	144
Fig. 548. <i>Oe. costatum</i> Transeau.....	146

(Figs. 533, 537, 541, 544 show also oögonia in optical section seen from above.)

	Page
Fig. 549. <i>Oedogonium macrandrium</i> Wittr.....	151
Fig. 550. <i>Oe. macrandrium</i> f. <i>acuminatum</i> Hirn. (Redrawn from Hirn).....	151
Fig. 551. <i>Oe. macrandrium</i> f. <i>aemulans</i> Hirn. (Redrawn from Hirn).....	151
Fig. 552. <i>Oe. macrandrium</i> f. <i>lundense</i> (Wittr.) Hirn. (Redrawn from Hirn).....	151
Fig. 553. <i>Oe. macrandrium</i> v. <i>propinquum</i> (Wittr.) Hirn.....	152
Figs. 554-556. <i>Oe. macrandrium</i> v. <i>hohenackerii</i> (Wittr.) Tiffany.....	152
Fig. 557. <i>Oe. macrandrium</i> v. <i>scrobiculatum</i> Ackley. (Redrawn from Ackley).....	152
Fig. 558. <i>Oe. laetevirens</i> Wittr. (Redrawn from Hirn).....	150
Fig. 559. <i>Oe. hoersholmiense</i> Hallas. (Redrawn from Hallas).....	150

	Page
Fig. 560. <i>Oedogonium rugulosum</i> Nordst.	148
Fig. 561. <i>Oe. rugulosum</i> f. <i>minutum</i> (Hansg.) Hirn. (Redrawn from Hirn)	149
Fig. 562. <i>Oe. rugulosum</i> f. <i>rotundatum</i> Hirn. (Redrawn from Hirn)	149
Fig. 563. <i>Oe. longatum</i> Kuetz.	148
Figs. 564 and 565. <i>Oe. monile</i> Berk. and Harv. (Redrawn from Hirn)	154
Fig. 566. <i>Oe. monile</i> f. <i>victoriense</i> G. S. West. (After West)	154
Fig. 567. <i>Oe. monile</i> f. <i>borgei</i> Hirn. (Redrawn from Hirn)	154
Figs. 568 and 569. <i>Oe. monile</i> v. <i>eminens</i> Hirn. (Redrawn from Hirn)	154
Figs. 570 and 571. <i>Oe. hians</i> Nord. and Hirn. (Redrawn from Hirn)	149
Fig. 572. <i>Oe. schmidlei</i> Gutw. (After Gutwinski)	148

	Page
Figs. 573-576. <i>Oedogonium supremum</i> Tiffany.....	156
Fig. 577. <i>Oe. obtruncatum</i> Wittr.....	155
Fig. 578. <i>Oe. obtruncatum</i> v. <i>ellipsoideum</i> Wittr. (Redrawn from Hirn).....	155
Figs. 579-582. <i>Oe. obtruncatum</i> v. <i>completum</i> Hirn. (Redrawn from Hirn).....	155
Fig. 583. <i>Oe. ciliatum</i> (Hass.) Pringsh.....	153

Figs. 584 and 585. <i>Oedogonium wabashense</i> Tiffany.....	Page
Figs. 586-588. <i>Oe. praticolum</i> Transeau.....	156
Fig. 589. <i>Oe. acrosporum</i> De Bary.....	157
Fig. 590. <i>Oe. acrosporum</i> f. <i>boreale</i> (Wolle) Hirn. (After Wolle).....	158
Fig. 591. <i>Oe. acrosporum</i> v. <i>majusculum</i> Nordst. (After Hirn).....	159
Fig. 592. <i>Oe. acrosporum</i> v. <i>bathmidosporum</i> (Nord.) Hirn. (After Hirn)....	158
Fig. 593. <i>Oe. acrosporum</i> v. <i>floridense</i> Wolle. (After Wolle).....	158

	Page
Figs. 594-596. <i>Oedogonium michiganense</i> Tiffany.....	157
Fig. 597. <i>Oe. tentoriale</i> Nordst. and Hirn.....	157
Figs. 598-600. Vegetative cell of <i>Oe. crassiusculum</i> Wittr. v. <i>idioandrosporum</i> N. and W. showing cell division and formation of apical cap.....	130
Figs. 601 and 602. <i>Oedocladium media</i> Lewis. (After Lewis).....	52
Fig. 603. <i>Oedocladium media</i> Lewis, habit sketch. (After Lewis).....	52

	Page
Figs. 604-608. <i>Oedocladium hazenii</i> Lewis, showing habit and reproductive structures. Fig. 606, escape of androspores from androsporangia. (After Lewis).....	52
Figs. 609-611. <i>Oedocladium hazenii</i> Lewis, stages in the germination of a zoospore. (After Lewis).....	52
Fig. 612. <i>Oedocladium hazenii</i> Lewis, sporeling. (After Lewis). (R, rhizoidal cell; G, green cell of vegetative filament).....	52

	Page
Figs. 613-615. <i>Oedocladium albemarlensis</i> Lewis. (After Lewis).....	51
Figs. 616-618. <i>O. albemarlensis</i> Lewis. Stages in cell division and branching.....	51
Fig. 619. <i>Oedocladium protonema</i> Stahl. Habit sketch, showing vegetative and rhizoidal cells, the latter with two resting cells. (Redrawn from Stahl.)	51
Figs. 620 and 621. <i>O. protonema</i> Stahl. Male and female branches. (Redrawn from Stahl).....	51
Fig. 622. <i>O. protonema</i> Stahl. Part of filament with three side branches. (Redrawn from Stahl).....	51
Fig. 623. <i>O. protonema</i> Stahl. Sporeling. (Redrawn from Stahl). (G, cell of green filament; R, rhizoidal cell).....	51

Figs. 624-630. Different positions of operculum in operculate oogonia of Oedogonium. 624, supreme; 625, superior; 626, supramedian; 627, median; 628, inframedian; 629, inferior; 630, infimus (basal).

Figs. 631-635. Different positions of pore in poriferous oogonia of Oedogonium. 631, superior; 632, supramedian; 633 and 634, median; 635, inferior.

Fig. 636. Subhemispherical basal cell of Oedogonium.

Fig. 637. Elongated type of basal cell of Oedogonium.

Fig. 638. Formation of oogonium O and suffultory cell S from a division of a vegetative cell, in Oedogonium.

Figs. 639 and 640. Suffultory cells of Oedogonium.

Figs. 641-645. Formation of oogonium in Bulbochaete. OO, young oogonium; PO, patent oogonium; EO, erect oogonium; US, upper suffultory cell; LS, lower suffultory cell; D, dwarf male with antheridium; A, androsporangium.

Figs. 646 and 647. Development of 4 zoospores from germinating oospore. (After Juranyi.)